



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Variation in the estimated prevalence of multimorbidity: systematic review and meta-analysis of 194 international studies

| | |
|-------------------------------|---|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2021-057017 |
| Article Type: | Original research |
| Date Submitted by the Author: | 04-Sep-2021 |
| Complete List of Authors: | Ho, Iris Szu-Szu; The University of Edinburgh Usher Institute of Population Health Sciences and Informatics Azcoaga-Lorenzo, Amaya; University of St Andrews Bute Medical School Akbari, Ashley; Swansea University Medical School Davies, Jim; University of Oxford, Department of Computer Science Hodgins, Peter; The University of Edinburgh Usher Institute of Population Health Sciences and Informatics Khunti, Kamlesh; University of Leicester College of Life Sciences Kadam, Umesh; University of Leicester College of Life Sciences Lyons, Ronan; Swansea University, Health Data Research UK McCowan, Colin; University of St Andrews, School of Medicine Mercer, Stewart; The University of Edinburgh, Usher Institute Nirantharakumar, Krishnarajah; University of Birmingham Guthrie, Bruce; University of Edinburgh |
| Keywords: | EPIDEMIOLOGY, GENERAL MEDICINE (see Internal Medicine), GERIATRIC MEDICINE |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Variation in the estimated prevalence of multimorbidity: systematic review and meta-analysis of 194 international studies

Iris Szu-Szu Ho,¹ Amaya Azcoaga-Lorenzo,² Ashley Akbari,³ Professor Jim Davies,⁴ Peter Hodgins,¹ Kamlesh Khunti,⁵ Umesh Kadam,⁵ Ronan A Lyons,³ Colin McCowan,² Stewart W Mercer,¹ Krishnarajah Nirantharakumar,⁶ Bruce Guthrie.¹

1. University of Edinburgh, Usher Institute, Medical School, Teviot Place, Edinburgh, EH8 9AG, United Kingdom
2. University of St Andrews, School of Medicine, Medical and Biological Sciences, North Haugh, St Andrews, KY16 9TF, United Kingdom
3. Swansea University, Medical School, Population Health Science, Singleton Park, Swansea, SA2 8PP, United Kingdom
4. University of Oxford, Department of Computer Science, Wolfson Building, Parks Road, Oxford, OX1 3QD, United Kingdom
5. University of Leicester, Leicester General Hospital, Gwendolen Road, Leicester, LE5 4PW, United Kingdom
6. University of Birmingham, Institute of Applied Health Research, Birmingham, B15 2TT, United Kingdom

Corresponding author: Iris Szu-Szu Ho. Email: iris.s.ho@outlook.com

Abstract

Objective. To examine the variation in the estimated prevalence of multimorbidity.

Methods. In this systematic review and meta-analysis, we conducted searches in nine bibliographic databases (PsycINFO, Embase, Global Health, Medline, Scopus, Web of Science, Cochrane Library, CINAHL, and ProQuest Dissertations & Theses Global) for prevalence studies published between database inception and 21 January 2020. Studies reporting the prevalence of multimorbidity (in all age groups and in community, primary care, care home and hospital settings) were included. Studies with an index condition or those that did not include people with no long-term conditions in the denominator were excluded. Retrieved studies were independently reviewed by two reviewers, and relevant data were extracted using pre-designed pro-forma. We used meta-analysis to pool the estimated prevalence of multimorbidity across studies, and used random-effects meta-regression and subgroup analysis to examine the association of heterogeneous prevalence estimates with study and measure characteristics.

Results. 13,807 titles were screened, of which 194 met inclusion criteria for meta-analysis. The pooled prevalence of multimorbidity was 42.7% (95%CI=39.2%-46.2%) with high heterogeneity ($I^2>99\%$). In adjusted meta-regression models, participant mean age and the number of conditions included in a measure accounted for 52.6% of heterogeneity in effect sizes. The estimated prevalence of multimorbidity was significantly higher in studies with older adults and those that included larger numbers of conditions. There was no significant difference in estimated prevalence between low- or middle-income countries (37.8%) and high-income countries (44.3%), or between self-report (40.0%) and administrative/clinical databases (52.7%).

Conclusions. The pooled prevalence of multimorbidity was significantly higher in older populations and when studies included a larger number of baseline conditions. The findings suggest that, to improve study comparability and quality of reporting, future studies should use a common core conditions set for multimorbidity measurement and report multimorbidity prevalence stratified by socio-demographics.

Funding: HDRUK (CFC0110)

Review registration: PROSPERO (CRD42020172409)

Strengths and limitations of this study

- This study used meta-regression to examine the variation of estimated prevalence of multimorbidity and how measure and study characteristics influenced prevalence estimates.
- The use of multiple imputation in this study minimised biased estimates caused by missing values and unbalanced classes and enhanced statistical accuracy.
- The inclusion of studies with various measure and study characteristics enabled a better understanding of the contributing factors of the heterogeneity of multimorbidity prevalence.
- Due to inconsistent reporting of multimorbidity prevalence and data unavailability, the estimated multimorbidity prevalence stratified by sex, ethnicity and socio-economic status could not be explored in this study.

Introduction

Population ageing is a worldwide phenomenon, with the World Health Organization (WHO, 2018) estimating that the proportion of the global population aged 60 and older will double from 12% to 22% between 2015 and 2050 [1]. A key implication of population ageing is that increasing numbers of people will be living with multimorbidity. Multimorbidity, commonly defined as the co-occurrence of two or more long-term conditions [2], adversely affects people’s risk of death, health-related quality of life, functional ability, and mental well-being [3, 4]. Multimorbidity affects all groups of society, but is known to be more common in older people, in women, and in those from low socio-economic backgrounds [5-7]. It poses major challenges to the delivery of care in health systems internationally which are often focused on the management of single diseases and lack appropriate coordination and continuity of care across different sectors [8, 9]. Disparities in health and health and social care could be found at any stage along the continuum of chronic diseases, from prevention to the management of diseases. To understand these disparities among multimorbid populations requires consistently monitoring the populations (e.g. incidence, prevalence, health impact, risk factors and delivery of care) defined by race and ethnicity, gender, age, socio-economic status, physical environment and geographic factors.

Previous systematic reviews have identified issues in the measurement of multimorbidity, related to the choice of chronic conditions counted in measures, the categorisation of conditions and diseases, and the counting or weighting method used [10, 11]. Although weighted measures are often used when the purpose of measurement is to predict future outcomes, a simple count of conditions remains the most commonly-used method for the measurement of multimorbidity, and is optimal for estimating multimorbidity prevalence [12]. However, the estimated prevalence of multimorbidity varies widely in the literature ranging from 3.5% to 100% [13], likely reflecting a combination of varying measures and varying populations studied [14]. This review aimed to examine variation in the estimated prevalence of multimorbidity, including estimated prevalence in different subgroups and associations with study and multimorbidity measure characteristics.

Methods

The systematic review and meta-analysis reported here is part of a larger review which aimed to examine 1) how multimorbidity has been constructed and 2) measured by international studies

(n=566) and 3) variation in the estimated prevalence of multimorbidity across studies. Analysis in relation to the first two registered objectives has been reported [15], and this paper reports the third registered objective.

Inclusion and exclusion criteria

The eligibility criteria for this review were defined based on the CoCoPop framework—Condition, Context, and Population [16]. The condition included in this review is prevalence of multimorbidity. The majority of studies defined multimorbidity as the co-existence of two or more chronic conditions, and used the cut-off to estimate its prevalence in a population of interest. We therefore included studies that used this definition for examining multimorbidity prevalence across international studies. For this analysis, we included studies carried out in the community, primary care, care home and hospitals, and those estimating the prevalence of multimorbidity in the population studied. Studies that did not include a relevant denominator population – for example, only examining patients with an index condition or excluding patients who did not have multimorbidity – were excluded. Qualitative research, studies not published in English, and conference abstracts were also excluded.

Search strategy

The search strategy for this review was developed in collaboration with a specialist medical librarian (Supplementary Table S1). Key terms relevant to multimorbidity and measurement were combined using Boolean logic to identify studies that met the inclusion criteria. We included medical subject headings to provide a sensitive search for relevant literature. Databases included in the search were Ovid interface (PsycINFO, Embase, Global Health, Medline), Scopus, Web of Science, Cochrane Library, EBSCO interface (CINAHL Plus), and ProQuest Dissertations & Theses Global, from inception to 21 January 2020. In addition to the database searches, our secondary search strategy included hand-searching reference lists of retrieved articles and tracked citations to maximise the yield.

Study screening and selection

Articles retrieved from databases were organised using EndNote X9 bibliographic software and Excel, and then were imported to Covidence for screening [17]. Titles, abstracts, and full-texts of retrieved articles were screened against the eligibility criteria by two reviewers. Throughout the review process, any disagreement that arose was resolved through discussion between the two

reviewers (IS-SH and PH), and through the involvement of a third reviewer (BG) if necessary. The study selection process is summarised in Figure 1.

Data extraction

We extracted data on the characteristics of the included studies using pre-designed data extraction pro-forma. The extracted data include 1) authors, 2) publication year, 3) study purpose, 4) method, 5) country, 6), continent, 7) country income (classified as ‘high’ and ‘low or medium’ [combined because of small numbers] allocated based on the World Bank Group at the time of review [18]), 8) study participants, 9) mean age, 10) sample size, 11) number of conditions, 12) setting, 13) data collection method/data source, 14) number of multimorbidity cases, and 15) proportion of multimorbidity (calculated based on item 10 and 14). Data on the estimated prevalence stratified by sex, ethnicity and socio-economic status were fragmented and unavailable in many studies, and thus these could not be retrieved for analyses.

Risk of bias assessment

We used the Effective Public Health Practice Project (EPHPP) quality assessment tool for quantitative studies to assess the risk of bias and the quality of each of the included studies, in terms of 1) selection bias, 2) study design, 3) confounders, 4) blinding, 5) data collection method, 6) withdrawals and dropouts [19]. We assessed also publication bias (rated high if there was selective reporting within studies) and conflict of interest (rated unclear if conflict of interest declaration was not reported). Each study was rated and assigned an overall risk of bias as ‘high’, ‘moderate’, or ‘low’ (please see the details in appendix p26).

Data analysis

We used descriptive statistics to measure frequency distribution, central tendency and variability of all variables. Univariate generalized linear models were used to investigate the association between continuous/count predictor (mean age/number of conditions) and categorical predictors. We summarised the prevalence of multimorbidity using metaprop [20, 21]. The presence of effect size heterogeneity was examined using the Q statistic and I-squared. Significant heterogeneity was identified, so we used subgroup analysis and meta-regression with random-effects models to identify potential moderating factors.

Outlying studies were identified using studentised residuals, leave-one-out analysis and Mahalanobis distance. Studies with studentised residuals that were larger than 2 or 3 and those that contributed to heterogeneity in leave-one-out analyses were scrutinized [22]. Mahalanobis distance was used for pattern recognition and multivariate outlier detection [23]. Study effect sizes were graphically displayed to identify outlying studies and explore subgroup effects (Supplementary Figure S1). In initial analysis of heterogeneity and outliers, 24 studies were found to make a significant contribution to the high level of observed heterogeneity in multimorbidity prevalence and significant changes in the summary effect size. The 24 studies were excluded for one or more of the following reasons: 1) their contribution to high levels of heterogeneity in the leave-one-out test, 2) being identified as an outlying value in the studentised residuals test ($z\text{-score} \geq 2$), 3) their Mahalanobis distance exceeding the chi-squared critical value at a 0.01 significance level, 4) infrequent values in compositional categorical data (e.g. only one study examined prevalence in children). The process of identifying outliers, the rationale for exclusion of each study, and the characteristics of outlying studies are documented in Supplementary Figure S2 and Table S2 and S3. Sensitivity analysis was performed to explore the impact of excluding the 24 studies in meta-analysis.

There was missingness in two predictors, with 37% missingness in the ‘mean age’ of the study population variable (some of which reported it categorically, and thus were treated as missing data) and 6% missingness in the ‘number of conditions’ included in the multimorbidity measure variable. Multiple imputation with 60 imputed datasets and 10 iterations was performed where random forest was used to impute missing data [24, 25]. Following multiple imputation, fraction of missing information (FMI) was computed to quantify the impact of missing data, which ranged from 0.05 to 0.3 indicating that the uncertainty in the values imputed for missing data is small/moderate [26].

A random-effects regression tree approach with ten-fold cross-validation was used to identify subgroups (cut-offs) of the ‘mean age’ and ‘number of conditions’ variables with differential effect sizes [27]. Given considerable variation in the effect sizes, we conducted meta-regression with the restricted maximum likelihood (REML) estimator to examine the possible sources of heterogeneity in effect sizes [20, 21, 28]. As the variable ‘multimorbidity prevalence’ did not follow the normal distribution (positively skewed), we applied logit transformation to the variable for analyses and converted the logits back to odds ratios (e^{logit}) and proportions ($p = e^{\text{logit}} / e^{\text{logit}+1}$) for reporting. For model selection, we refitted the models using maximum likelihood and then

conducted a log-likelihood test to compare the fit of models [29]. A permutation test with 1000 permuted datasets was performed on the final meta-regression model to calculate p value and avoid type 1 error [30]. Subgroup analysis with the REML method was used to estimate the pooled multimorbidity prevalence of subgroups of each variable (age, the number of conditions included in a measure, setting, data source, continent, country income, study risk of bias). Forest-like plots were used to display the effect sizes of included studies [31]. The presence of publication bias was assessed using Egger’s test, which did not find evidence of publication bias [32]. All statistical tests were performed using R version 4.0.4.

Patients and public involvement

No patients were involved in the development of the research question, outcome measures, study design and implementation. Nonetheless, we have previously discussed preliminary review findings and issues relevant to multimorbidity measurement with our patient and public involvement group. We plan to disseminate the review findings to researchers, clinicians, policy makers and public audiences through news media, social media and seminars.

Results

After screening 13,807 titles and abstracts, 218 studies were identified which estimated the prevalence of multimorbidity using a cut-off of ‘two or more’ conditions. Following the removal of 24 outlying studies, 194 studies were included in the meta-analysis (Table 1, Supplementary Table S4). Of the 194 studies, 64 studies were from Europe, 47 from North America, 45 from Asia, 11 from Australasia, 12 from South America, and four from Africa (Table 1 and Figure

2). Seventy-four percent of studies were from high-income countries (n=145) and 25.3% from low- and middle-income countries (LMICs) (one from low-income, nine from lower middle-income, 29 from upper middle-income, and 10 from multiple low- and middle-income countries). The majority of studies (n=147) estimated the prevalence of multimorbidity in community settings, followed by primary care (n=33) and hospital setting (n=14). Prevalence data were collected through either self-report (n=151) or medical records and administrative databases (n=43). In a univariate linear regression (Supplementary Table S5), we found that studies from Europe, database studies and studies conducted in hospital settings were more likely to measure multimorbidity in an older population and included a larger number of conditions in a

multimorbidity measure, compared to those from other continents, self-report studies, and studies conducted in primary care and community settings. In respect to risk of bias in included studies (Supplementary Table S6 and Figure S3), 11.3% were rated as high risk of bias, 84% as moderate risk of bias, and 4.6% as low risk of bias.

The pooled estimate of multimorbidity prevalence across the 194 studies was 42.7% (95%CI 39.2%-46.2%), τ^2 is 1.0 (95%CI 0.9-1.3) with high heterogeneity ($I^2 > 99\%$), and meta-regression was therefore used to examine study characteristics associated with heterogeneity. Mean age ($F=103.1$, $p<0.0001$, $R^2=34.7\%$) and number of conditions ($F=34.4$, $p<0.0001$, $R^2=14.8\%$) were the strongest univariate predictors and positively associated with the estimated prevalence of multimorbidity (Figure 3). Meta-regression tree analysis (Supplementary Figure S4) partitioned the mean age variable into three homogeneous subgroups (aged <59 , aged 59-73, aged ≥ 74) and the number of conditions variable into four homogeneous subgroups (<9 , 9-19, 20-43, ≥ 44). The categorical 'mean age' and 'number of conditions' variables explained 42.7% and 17.3% of the heterogeneity in effect sizes respectively (larger than the original numerical variables). We therefore used the categorical variables identified from the regression trees for meta-analyses.

In univariate meta-regression, primary care studies (pooled multimorbidity prevalence 49.7%, OR 1.5, 95%CI 1.0-2.2) and hospital based studies (pooled multimorbidity prevalence 59.6%, OR 2.2, 95%CI 1.3-3.9) had significantly higher rates of multimorbidity than community-based studies (39.5%) (Table 2). Multimorbidity prevalence was significantly higher in database studies (pooled multimorbidity prevalence 52.7%, OR 1.7, 95%CI 1.2-2.4) than self-report studies (pooled multimorbidity prevalence 40.0%). In the mean age categorical variable, the pooled prevalence estimates of the three subgroups were statistically significantly different from one another, and considerably higher in studies with mean participant age ≥ 74 (pooled multimorbidity prevalence 69.0%, OR 5.7, 95%CI 4.2-7.7) and mean participant age 59-73 (pooled multimorbidity prevalence 50.3%, OR 2.6, 95%CI 2.0-3.3) than those with mean participant age <59 (pooled multimorbidity prevalence 28.2%) (Table 2 and Figure 4). Similar patterns were also found in the number of conditions variable where studies including ≥ 44 conditions in measurement (pooled multimorbidity prevalence 87.6%, OR 15.0, 95%CI 5.9-38.3), 20-43 conditions (pooled multimorbidity prevalence 51.4%, OR 2.2, 95%CI 1.5-3.3), and 9-19 conditions (pooled multimorbidity prevalence 43.2%, OR 1.6, 95%CI 1.2-2.2) yielded higher prevalence estimates than studies including <9 conditions in measurement (pooled multimorbidity prevalence 32.1%)

with a dose-response relationship. The estimated prevalence of multimorbidity was 44.3% in high-income countries compared to 37.8% in low or middle income countries, but the difference was not statistically significantly different (OR 1.3, 95%CI 0.9-1.8). In study risk of bias, no statistically significant difference in pooled prevalence of multimorbidity was found between studies with low, moderate and high risk of bias.

In the adjusted meta-regression model, we found that compared to studies where participant mean age was <59, multimorbidity prevalence remained significantly higher in studies with mean participant age 59-73 (OR 2.5, 95%CI 2.0-3.1) and in studies with mean participant age ≥74 (OR 4.7, 95%CI 3.6-6.2). Compared to measures including <9 conditions, multimorbidity prevalence was higher in measures including ≥44 conditions (OR 7.3, 95%CI 3.5-15.0), 20-43 conditions (OR 2.1, 95%CI 1.5-2.8), and 9-19 conditions (OR 1.6, 95%CI 1.3-2.0). Nonetheless, no difference was found between primary care, community, and hospital settings. Compared to studies from North America, prevalence was lower in studies from Europe (OR 0.6, 95%CI 0.4-0.7), Australasia (OR 0.6, 95%CI 0.4-0.9), or Asia (OR 0.6, 95%CI 0.4-0.7). No significant difference in prevalence estimates between self-report and routine database studies was evident after controlling for study and measure characteristics. The model explained 56.8% of the heterogeneity in multimorbidity prevalence, with the mean age and number of conditions variables providing most explanatory power (52.6% of the heterogeneity).

Sensitivity analysis including the 24 outlying studies (Supplementary Table S7) was similar to primary analysis except for “study setting” variable. The mean participant age and number of conditions variables remained the strongest predictors of multimorbidity prevalence in sensitivity analysis. However, the estimated prevalence in sensitivity analysis (including outlying studies) was much lower in studies including ≥44 conditions in a multimorbidity measure (pooled multimorbidity prevalence 51.6, OR 2.5, 95%CI 1.5-4.0) compared to primary analysis excluding outlying studies (pooled multimorbidity prevalence 87.6, OR 7.3, 95%CI 3.5-15.0). The difference in estimates was mainly attributed to the three outlying studies that included 146, 147 and 259 conditions in a measure respectively but yielded relatively low mean multimorbidity prevalence (mean prevalence 54.3%)[33-35]. In respect to study settings, the pooled prevalence in sensitivity analysis was statistically significantly higher in primary care compared to community in both unadjusted and adjusted models, whereas in primary analysis the difference was not statistically significant after controlling for study and measure characteristics. The higher pooled prevalence

in primary care settings found in sensitivity analysis could be explained by two outlying primary care based studies that had mean participant age of 56 but high estimated prevalence (89.1% and 72.7% respectively) [34, 36].

Discussion

The overall estimate of multimorbidity prevalence in adults across all the included studies was 42.7% (95%CI=39.2%-46.2%), but with very high heterogeneity. More than half of the observed heterogeneity was explained by study mean participant age and the number of conditions included in the multimorbidity measure, with older age and larger number of conditions strongly associated with a higher prevalence of multimorbidity. The difference in estimated prevalence was small between self-report and administrative/clinical databases, and between study settings. No significant difference was found between studies from low- or middle-income and high-income countries, but North American studies had higher estimated prevalence than other continents.

Two prior systematic reviews examined the prevalence of multimorbidity across studies [37, 38]. Fortin et al. (2012) conducted a narrative review of 21 studies and found various operationalisations of multimorbidity and a large variation in the prevalence of multimorbidity, particularly in studies with participants aged 75 and older [37]. Nguyen et al. (2019) meta-analysed the prevalence of multimorbidity across 70 studies from community settings and found that the pooled estimated prevalence was 33.1% with high levels of heterogeneity ($I^2 > 99\%$) [38]. The pooled prevalence of multimorbidity in Nguyen et al study is lower than in this study, likely because we have included studies from primary care and hospital settings (the pooled prevalence of multimorbidity in community-based studies in this analysis was 39.5%). Nguyen et al. (2019) did not carry out a meta-regression, but in narrative analysis comment that the prevalence of multimorbidity appeared higher in older adults and women [38]. Our review findings are consistent with previous literature finding that age is most important determinant of multimorbidity [5, 37-39]. While we did not find a significant difference between low and middle-income and high-income countries, Nguyen et al. in their review showed a statistically significantly higher pooled prevalence in high-income countries (the pooled prevalence from 18 studies was 37%) than low or middle-income countries (the pooled prevalence from 31 studies was 29%). This difference in findings may be due to the inclusion in our review of a larger number of studies from high-income or upper middle-income countries, whereas very few studies were from low-income or lower

middle-income countries. The low number of included studies from low-income countries in this review could be explained by fewer attention paid to this relatively new research field (multimorbidity) in low-income countries and our literature search restricted to English language (proficient language of reviewers). The estimated prevalence of multimorbidity in North America was higher compared to other continents in this study despite older study populations and larger numbers of conditions found in studies from Europe. Possible explanation of the results could be related to over-diagnosis and medicalisation [40].

The strengths of this review are searches conducted in multiple databases, the large number of studies identified and the use of meta-analytic approaches to examine factors associated with heterogeneity of estimated multimorbidity prevalence. We examined and handled outlying studies and missing data (multiple imputation) with rigour and excluded studies that did not take into account ‘healthy’ populations (populations with no long-term conditions) to minimize biased estimates of multimorbidity prevalence. This review has limitations. Sensitivity analysis including all studies had similar findings with two exceptions, namely that sensitivity analysis found: a weaker (but still statistically significant) association with the number of conditions included in the multimorbidity measure than primary analysis; and a statistically significantly higher pooled prevalence in primary care compared to community based studies versus no significant association in primary analysis. Although we examined associations with study characteristics including mean participant age, a limitation is the lack of information in the reviewed studies on prevalence estimates stratified by participant characteristics including sex, ethnicity, and socio-economic status. An additional uncontrolled factor is how studies measured multimorbidity in terms of the type (as opposed to the number) of the conditions included in measures, which varied substantially across studies with too much heterogeneity to model [15]. Last but not least, measurement of multimorbidity is a relatively new research field and its labelling has been used variably. Thus, it is likely that not all relevant studies were identified and included in this review, but we were rigorous in our application of inclusion/exclusion criteria and did not favour adding known papers that did not appear in the search or where excluded through the process.

In spite of the methodological limitations, this review adds to our understanding of how study and measure characteristics can influence the estimated prevalence of multimorbidity. Mean age of the study population and the number of conditions included in the multimorbidity measure were the major factors associated with varying estimated prevalence of multimorbidity. A key implication

is that comparing prevalence between studies requires more stratified estimates of multimorbidity prevalence. We therefore strongly recommend that as well as overall prevalence, future studies should clearly report multimorbidity prevalence stratified by age and sex at a minimum, and ideally by ethnicity and socio-economic status. This will allow readers to capture a more holistic picture of multimorbidity prevalence in the population studied, and allow better comparison of prevalence in different populations, and accurate pooled estimates of prevalence in reviews.

Additionally, the number of conditions included in a measure is strongly associated with estimated multimorbidity prevalence. It would be ideal if studies additionally reported prevalence using a common core set of conditions agreed by consensus. Parallel reporting of the bespoke set chosen for the context and purpose, and a core set would improve comparability of prevalence estimates, and help identify the additional value of any bespoke multimorbidity measures. The lack of any significant difference in estimated prevalence between self-report and clinical/administrative databases in this review suggests that provided careful attention is paid to the number and type of conditions included in measures, exactly how data is collected may be less important.

To conclude, in recent years, there has been an increasing interest in the epidemiology of multimorbidity internationally. This review finds that population characteristics and measurement content are the major factors that influenced prevalence estimates of multimorbidity. Studies with older populations and larger numbers of conditions yielded a higher estimate of multimorbidity prevalence. However, heterogeneity between studies has made comparison of multimorbidity prevalence across studies difficult. To improve comparability and quality of reporting, this review suggests that future studies should use common core condition set for the measurement of multimorbidity and clearly report the prevalence of multimorbidity stratified by socio-demographics.

Contributorship statement

All authors have made substantial contributions: CMC, KN, UK, KK, RAL, JD, AA, AAL and SWM were involved in conception of the work, acquisition of funding, and critically commenting on the manuscript. IS-SH led and BG substantially contributed to the design, analysis, and interpretation of data for the review, and are responsible for the decision to submit the manuscript. IS-SH and PH screened and reviewed retrieved studies. All authors contributed to the edits of the manuscript and had access to the data. The final draft has been approved by all authors.

Competing interests

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: we had financial support from HDRUK for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Funding

This project was funded by Health Data Research UK (HDRUK: CFC0110).

Data sharing statement

Study data are available in supplementary appendix.

Figure legends

Figure 1: Preferred reporting items for systematic reviews and meta-analyses (PRISMA) flow diagram

Figure 2: Country of origin of the included studies estimating the prevalence of multimorbidity (except studies from multiple countries)

Figure 3: Relationship between the prevalence of multimorbidity and mean age or number of conditions (the area of points is proportional to inverse variances)

Figure 4: The distribution of prevalence estimates within the subgroups of mean age and number of conditions (forest-like plot for a large review)

Table 1: Summary of study characteristics (Supplementary Table S8 shows the definition of variables)

| Name of variable | Descriptive statistics (n=194) |
|--|--|
| Prevalence of multimorbidity (%) | Range: 5.4 to 95.6 Median: 38.4 (IQR 28.3 to 57.0) Mean: 43.5 (SD 20.7) Pooled prevalence with the REML estimator: 42.7 (39.2-46.2) |
| Mean age of study population (year) | Range: 32.2-83.8 Median: 62.4 (IQR 49.9 to 72.3) Mean: 61.1 (SD 12.8) |
| No of conditions (count) | Range: 3-75 Median: 14 (IQR 9 to 20) Mean: 17 (SD 10) |
| Country income (count, %) | |
| High income | 145 (74.7%) |
| Low- or Middle-income | 49 (25.3%) |
| Continent (count, %) | |
| Europe | 64 (33.0%) |
| North America | 47 (24.2%) |
| Asia | 45 (23.2%) |
| Australasia | 11 (5.7%) |
| South America | 12 (6.2%) |
| Africa | 4 (2.1%) |
| Multiple continents | 11 (5.7%) |
| Study population (count, %) | |
| Only older people | 63 (32.5%) |
| Middle-aged and older | 46 (23.7%) |
| All adults | 85 (43.8%) |
| Setting (count, %) | |
| Community | 147 (75.8%) |
| Primary care | 33 (17.0%) |
| Hospital | 14 (7.2%) |
| Source (count, %) | |
| Self-report | 151 (77.8%) |
| Database | 43 (22.2%) |
| Risk of bias assessment (count, %) | |
| Low | 9 (4.6%) |
| Moderate | 163 (84.0%) |
| High | 22 (11.3%) |

IQR: Interquartile range. SD: Standard deviation. The percentages were rounded so they do not add to 100%.

Table 2: Output of meta-analytic models (n=194)

| | Pooled prevalence of multimorbidity of each subgroup (% , 95% CI) | Meta-regression Unadjusted Odds Ratio (95% CI) | Meta-regression Adjusted Odds Ratio (95% CI) R ² 56.8% | FMI |
|---------------------------|---|--|--|------|
| Group of mean age | | R ² 42.7% | | |
| <59 | 28.2 (25.4-31.2) | Ref | Ref | Ref |
| 59-73 | 50.3 (45.3-55.3) | 2.6 (2.0-3.3)*** | 2.5 (2.0-3.1)*** | 0.3 |
| ≥74 | 69.0 (62.9-74.5) | 5.7 (4.2-7.7)*** | 4.7 (3.6-6.2)*** | 0.2 |
| No of conditions | | R ² 17.3% | | |
| <9 | 32.1 (27.3-37.2) | Ref | Ref | Ref |
| 9-19 | 43.2 (38.9-47.7) | 1.6 (1.2-2.2)** | 1.6 (1.3-2.0)*** | 0.2 |
| 20-43 | 51.4 (42.9-59.7) | 2.2 (1.5-3.3)*** | 2.1 (1.5-2.8)*** | 0.2 |
| ≥44 | 87.6 (81.3-92.0) | 15.0 (5.9-38.3)*** | 7.3 (3.5-15.0)*** | 0.05 |
| Setting | | R ² 4.8% | | |
| Community | 39.5 (36.1-43.1) | Ref | Ref | Ref |
| Primary care | 49.7 (39.1-60.4) | 1.5 (1.0-2.2)* | 1.2 (0.9-1.7) | 0.1 |
| Hospital | 59.6 (45.6-72.2) | 2.2 (1.3-3.9)** | 1.2 (0.7-1.8) | 0.2 |
| Source | | R ² 3.9% | | |
| Self-report | 40.0 (36.2-43.8) | Ref | Ref | Ref |
| Database | 52.7 (45.2-60.1) | 1.7 (1.2-2.4)** | 0.9 (0.7-1.3) | 0.2 |
| Continent | | R ² 4.1% | | |
| North America | 50.4 (43.6-57.3) | Ref | Ref | Ref |
| Europe | 44.8 (38.2-51.5) | 0.8 (0.5-1.2) | 0.6 (0.4-0.7)*** | 0.2 |
| Australasia | 35.8 (29.5-42.5) | 0.5 (0.3-1.0)** | 0.6 (0.4-0.9)** | 0.07 |
| Asia | 35.3 (29.3-50.0) | 0.5 (0.4-0.8)* | 0.6 (0.4-0.7)*** | 0.1 |
| South America | 47.5 (31.2-64.4) | 0.9 (0.5-1.7) | 0.8 (0.5-1.3) | 0.1 |
| Africa | 23.6 (16.3-32.8) | 0.3 (0.1-0.8) | 0.6 (0.3-1.1) | 0.2 |
| Multiple continents | 38.4 (29.1-48.6) | 0.6 (0.3-1.2) | 0.7 (0.4-1.1) | 0.1 |
| Country income | | R ² 0.8% | | |
| Low or middle-income | 37.8 (31.4-44.7) | Ref | | |
| High-income | 44.3 (40.3-48.4) | 1.3 (0.9-1.8) | | |
| Study risk of bias | | R ² 0.0% | | |
| Low risk | 33.3 (20.2-49.6) | Ref | | |
| Moderate risk | 42.7 (39.1-46.4) | 1.5 (0.8-2.9) | | |
| High risk | 46.4 (34.1-59.1) | 1.7 (0.8-3.8) | | |
| Publication year | | 1.0 (1.0-1.0) | | |

*<0.05 **<0.01 ***<0.001

Ref: Reference category. FMI: Fraction of missing information


References

1. World Health Organization. *Ageing and health*. 2018 [cited 2021 April 1]; Available from: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>.
2. The Academy of Medical Sciences, *Multimorbidity: a priority for global health research*. 2018.
3. Rivera-Almaraz, A., et al., *Disability, quality of life and all-cause mortality in older Mexican adults: association with multimorbidity and frailty*. BMC Geriatr, 2018. **18**(236): p. 1-9.
4. Arokiasamy, P., et al., *The impact of multimorbidity on adult physical and mental health in low- and middle-income countries: what does the study on global ageing and adult health (SAGE) reveal?* BMC Med, 2015. **13**(178): p. 1-16.
5. Barnett, K., et al., *Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study*. The Lancet, 2012. **380**(9836): p. 37-43.
6. Low, L.L., et al., *Epidemiologic Characteristics of Multimorbidity and Sociodemographic Factors Associated With Multimorbidity in a Rapidly Aging Asian Country*. JAMA Network Open, 2019. **2**(11): p. 1-14.
7. Fortin, M., et al., *Prevalence estimates of multimorbidity: a comparative study of two sources*. BMC health services research, 2010. **10**(111): p. 1-6.
8. Pearson-Stuttard, J., M. Ezzati, and E.W.J.T.L.P.H. Gregg, *Multimorbidity—a defining challenge for health systems*. The Lancet, 2019. **4**(12): p. e599-e600.
9. Moffat, K. and S.W.J.B.f.p. Mercer, *Challenges of managing people with multimorbidity in today's healthcare systems*. BMC Family Practice, 2015. **16**(1): p. 1-3.
10. Johnston, M.C., et al., *Defining and measuring multimorbidity: a systematic review of systematic reviews*. Eur J Public Health, 2019. **29**(1): p. 182-189.
11. Diederichs, C., K. Berger, and D.B. Bartels, *The measurement of multiple chronic diseases--a systematic review on existing multimorbidity indices*. J Gerontol A Biol Sci Med Sci, 2011. **66**(3): p. 301-11.
12. Huntley, A.L., et al., *Measures of multimorbidity and morbidity burden for use in primary care and community settings: a systematic review and guide*. Ann Fam Med, 2012. **10**(2): p. 134-41.
13. Xu, X., G.D. Mishra, and M. Jones, *Evidence on multimorbidity from definition to intervention: An overview of systematic reviews*. Ageing Res Rev, 2017. **37**: p. 53-68.
14. Harrison, C., et al., *Examining different measures of multimorbidity, using a large prospective cross-sectional study in Australian general practice*. BMJ Open, 2014. **4**(7): p. 1-9.
15. Ho, I.S.-S., et al., *Examining variation in the measurement of multimorbidity in research: a systematic review of 566 studies*. The Lancet Public Health, June 21, 2021 (published online: [https://www.thelancet.com/journals/lanpub/article/PIIS2468-2667\(21\)00107-9/fulltext](https://www.thelancet.com/journals/lanpub/article/PIIS2468-2667(21)00107-9/fulltext)).
16. Munn, Z., et al., *Methodological guidance for systematic reviews of observational epidemiological studies reporting prevalence and cumulative incidence data*. Int J Evid-based Hea, 2015. **13**(3): p. 147-153.
17. Covidence. *Covidence 2020* [cited 2020 January 14]; Available from: <https://www.covidence.org/>.

18. World Bank Group. *The world by income*. 2018 [cited 2021 01 June]; Available from: <https://datatopics.worldbank.org/world-development-indicators/the-world-by-income-and-region.html>.
19. Armijo-Olivo, S., et al., *Assessment of study quality for systematic reviews: a comparison of the Cochrane Collaboration Risk of Bias Tool and the Effective Public Health Practice Project Quality Assessment tool: methodological research*. J Eval Clin Pract, 2012. **18**(1): p. 12-18.
20. Viechtbauer, W. *Package 'metafor'*. 2020 [cited 2020 August 20]; Available from: <https://cran.r-project.org/web/packages/metafor/metafor.pdf>.
21. Schwarzer, G. *Package 'meta'*. 2021 [cited 2021 Feb 15]; Available from: <https://cran.r-project.org/web/packages/meta/meta.pdf>.
22. Viechtbauer, W. and M.W.L. Cheung, *Outlier and influence diagnostics for meta-analysis*. Research Synthesis Methods, 2010. **1**(2): p. 112-125.
23. Leys, C., et al., *Detecting multivariate outliers: Use a robust variant of the Mahalanobis distance*. Journal of Experimental Social Psychology, 2018. **74**: p. 150-156.
24. Tang, F. and H. Ishwaran, *Random forest missing data algorithms*. Statistical Analysis and Data Mining: The ASA Data Science Journal, 2017. **10**(6): p. 363-377.
25. van Buuren, S. *mice: Multivariate Imputation by Chained Equations. R package version: 4.0.1*. 2020 [cited 2020 1 Aug]; Available from: <https://cran.r-project.org/web/packages/mice/mice.pdf>.
26. White, I.R., P. Royston, and A.M. Wood, *Multiple imputation using chained equations: Issues and guidance for practice*. Stat Med, 2011. **30**(4): p. 377-99.
27. Li, X., et al. *Package 'metacart'*. 2020 [cited 2020 May 20]; Available from: <https://cran.r-project.org/web/packages/metacart/metacart.pdf>.
28. Langan, D., et al., *A comparison of heterogeneity variance estimators in simulated random-effects meta-analyses*. Res Synth Methods, 2019. **10**(1): p. 83-98.
29. Verbyla, A.P., *A note on model selection using information criteria for general linear models estimated using REML*. Australian & New Zealand Journal of Statistics, 2019. **61**(1): p. 39-50.
30. Gagnier, J.J., et al., *An empirical study using permutation-based resampling in meta-regression*. Systematic reviews, 2012. **1**(1): p. 1-9.
31. Nakagawa, S., et al., *The orchard plot: Cultivating a forest plot for use in ecology, evolution, and beyond*. Research Synthesis Methods, 2021. **12**(1): p. 4-12.
32. Shi, X., et al., *Effect comparison between Egger's test and Begg's test in publication bias diagnosis in meta-analyses: evidence from a pilot survey*. International Journal of Research Studies in Biosciences, 2017. **5**(5): p. 14-20.
33. Violán, C., et al., *Burden of multimorbidity, socioeconomic status and use of health services across stages of life in urban areas: a cross-sectional study*. BMC public health, 2014. **14**(530): p. 1-13.
34. Prazeres, F. and L. Santiago, *Prevalence of multimorbidity in the adult population attending primary care in Portugal: a cross-sectional study*. BMJ Open, 2015. **5**(9): p. 1-10.
35. Sullivan, P.W., V.H. Ghushchyan, and E.A. Bayliss, *The impact of co-morbidity burden on preference-based health-related quality of life in the United States*. Pharmacoeconomics, 2012. **30**(5): p. 431-42.

- 1
- 2
- 3
- 4 36. Fortin, M., et al., *Prevalence of multimorbidity among adults seen in family practice*. Ann Fam Med, 2005. **3**(3): p. 223-8.
- 5
- 6 37. Fortin, M., et al., *A systematic review of prevalence studies on multimorbidity: toward a more uniform methodology*. Ann Fam Med, 2012. **10**(2): p. 142-51.
- 7
- 8 38. Nguyen, H., et al., *Prevalence of multimorbidity in community settings: A systematic review and meta-analysis of observational studies*. J Comorb, 2019. **9**: p. 1-15.
- 9
- 10 39. Hu, R.-H., et al., *Increasing age- and gender-specific burden and complexity of multimorbidity in Taiwan, 2003–2013: a cross-sectional study based on nationwide claims data*. BMJ Open, 2019. **9**(6): p. 1-10.
- 11
- 12
- 13
- 14 40. Brodersen, J., et al., *Overdiagnosis: what it is and what it isn't*. BMJ Evidence-Based Medicine, 2018. **23**(1): p. 1-3.
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41






Figure 2: Country of origin of the included studies estimating the prevalence of multimorbidity (except studies from multiple countries)

1182x571mm (118 x 118 DPI)




Figure 3: Relationship between the prevalence of multimorbidity and mean age or number of conditions (the area of points is proportional to inverse variances)

1182x571mm (118 x 118 DPI)




Figure 4: The distribution of prevalence estimates within the subgroups of mean age and number of conditions (forest-like plot for a large review)

1182x571mm (118 x 118 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary appendix

Supplement to: Ho ISS, Azcoaga-Lorenzo A, Akbari A, et al. Variation in the estimated prevalence of multimorbidity: systematic review and meta-analysis of 194 studies.

For peer review only

Table of Contents

| | |
|---|----|
| Table S1: Search strategy..... | 3 |
| Table S2: Summary of outlying studies | 4 |
| Table S3: Characteristics of 24 outlying studies..... | 5 |
| Table S4: Characteristics of 194 included studies | 9 |
| Table S5: Associations between predictors | 25 |
| Table S6: Risk of bias assessment of included studies | 26 |
| Table S7: Output of adjusted meta-analytic model based on 218 studies | 39 |
| Table S8: Definition of variables..... | 40 |
| Figure S1: Graphical display of study effect sizes and heterogeneity | 41 |
| Figure S2: Process of examining and identifying outlying studies in meta-analysis | 42 |
| Figure S3: Summary of risk of bias assessment | 43 |
| Figure S4: Meta-regression trees for predicting the pooled estimated prevalence of multimorbidity (based on 'mean age' and 'number of conditions' predictors. unit: log(odds))..... | 44 |
| References..... | 45 |

Table S1: Search strategy

| Database | Search strategy |
|--|---|
| <div>Ovid Interface</div> <div>PsycINFO</div> <div>Embase</div> <div>Global Health</div> <div>Ovid MEDLINE</div> | <div>1. (multimorbidit\$ or multi-morbidit\$ or comorbidit\$ or co-morbidit\$ or polymorbidit\$ or poly-morbidit\$ or multicondition\$ or multicondition\$ or “multiple chronic condition\$” or “morbidity burden” or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) adj2 (disease\$ or illness\$ or condition\$ or diagnos\$ or morbid\$))).m_titl.</div> <div>2. (measure\$ or index or indices or instrument\$ or scale\$ or “disease count\$”).mp.</div> <div>3. 1 and 2</div> <div>4. Limit 3 to human</div> |
| <div>EBSCO Interface</div> <div>CINAHL Plus</div> | <div>1. MM (multimorbidit* or multi-morbidit* or comorbidit* or co-morbidit* or polymorbidit* or poly-morbidit* or multicondition* or multicondition* or “multiple chronic condition*” or “morbidity burden” or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) N2 (disease* or illness* or condition* or diagnos* or morbid*)))</div> <div>2. AB (measure* or index or indices or instrument* or scale*)</div> <div>3. 1 AND 2</div> <div>Limiters – Full Text; Human; Language: English</div> |
| Scopus | TITLE (multimorbidit* or multi-morbidit* or comorbidit* or co-morbidit* or polymorbidit* or poly-morbidit* or multicondition* or multicondition* or “multiple chronic condition*” or “morbidity burden” or ((multiple or coexisting or co-existing or concurrent or con-current or morbid or co-morbid) W/2 (disease* or illness* or condition* or diagnos?s or morbid*))) AND TITLE (measure* or index or indices or instrument* or scale* or “disease counts”) |
| Web of Science | (TI=(measure* or index or indices or instrument* or scale*))AND (TI=(multimorbidit* or multi-morbidit* or comorbidit* or co-morbidit* or polymorbidit* or poly-morbidit* or multicondition* or multicondition* or 'multiple chronic condition*' or 'morbidity burden' or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) NEAR/2 (disease* or illness* or condition* or diagnos* or morbid*)))) AND LANGUAGE: (English) |
| Cochrane library | (multimorbidity or multi-morbidity or comorbidity or co-morbidity or polymorbidity or poly-morbidity or multicondition or multicondition or 'multiple chronic conditions' or 'morbidity burden' or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) NEAR/2 (disease or illness or condition or diagnosis or morbid))) AND (measure or index or indices or instrument or scale or “disease count*”):ti |
| <div>ProQuest Dissertations & Theses</div> <div>Global</div> | <div>ti((multimorbidit* OR multi-morbidit* OR comorbidit* OR co-morbidit* OR polymorbidit* OR poly-morbidit* OR multicondition* OR multicondition* OR 'multiple chronic condition*' OR 'morbidity burden' OR ((multiple OR coexisting OR co-existing OR concurrent OR con-current OR morbid OR co-morbid) NEAR/2 (disease* OR illness* OR condition* OR diagnos?s OR morbid*)))) AND noft((measure* OR index OR indices OR instrument* OR scale*))</div> <div>Limited by: Manuscript type: Doctoral dissertations, Master's theses</div> <div>Language: English</div> |

Table S2: Summary of outlying studies

| Name of variable | Outlying studies (n=24) |
|--|--|
| Prevalence of multimorbidity (%) | Range: 7.3 to 89.1 Median: 28.1 (IQR 14.6 to 48.7) Mean: 34.3 (SD 23.5) |
| Mean age of study population (year) | Range: 39.6 to 82.2 Median: 56.6 (IQR 52.3 to 66.4) Mean: 59.3 (SD 11.5) |
| No of conditions (count) | Range: 7 to 259 Median: 34 (IQR 19 to 54) Mean: 52 (SD 58) |
| Country income (count, %) | |
| High income | 21 (87.5%) |
| Low- or Middle-income | 3 (11.5%) |
| Continent (count, %) | |
| Europe | 6 (25.0%) |
| North America | 7 (29.2%) |
| Asia | 7 (29.2%) |
| Australasia | 3 (12.5%) |
| Multiple continents | 1 (4.2%) |
| Study population (count, %) | |
| Only older people | 2 (8.3%) |
| Middle-aged and older | 1 (4.2%) |
| All adults | 15 (62.5%) |
| Only children | 1 (4.2%) |
| All age population | 5 (20.8%) |
| Setting (count, %) | |
| Community | 12 (50.0%) |
| Primary care | 7 (29.2%) |
| Hospital | 4 (16.7%) |
| Care home | 1 (4.2%) |
| Source (count, %) | |
| Self-report | 8 (33.3%) |
| Database | 16 (66.6%) |
| Risk of bias assessment (count, %) | |
| Low | 4 (16.7%) |
| Moderate | 19 (79.2%) |
| High | 1 (4.2%) |

IQR: Interquartile range. SD: Standard deviation. The percentages were rounded so they do not add up to 100%.

Table S3: Characteristics of 24 outlying studies

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias | Rationale for exclusion |
|-----------------------------------|-------------|---------------|----------------|-----------|---------------------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|---|
| ¹ Stanley et al (2018) | New Zealand | Australasia | High | Hospitals | All adults | Not reported | 3489747 | Medical records and administrative database | 61 | 275706 | 0.08 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) and the studentized residual of this study is more than 2 standard deviations away from its expected value. |
| ² Lenzi et al (2016) | Italy | Europe | High | Hospitals | All adults | 66.4 | 3759836 | Medical records and administrative database | 26 | 574208 | 0.15 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ³ Hu et al (2019) | Taiwan | Asia | High | Community | All adults | Not reported | 1429527 | Medical records and administrative database | 20 | 939485 | 0.66 | Low | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ⁴ Gawron et al (2020) | USA | North America | High | Hospitals | All adults but not older people | Not reported | 741612 | Medical records and administrative database | Not reported | 53824 | 0.07 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) and the studentized residual of this study is more than 2 standard deviations away from its expected value. |
| ⁵ Low et al (2019) | Singapore | Asia | High | Community | All adults | 39.6 | 1181024 | Self-report | 48 | 309428 | 0.26 | Low | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ⁶ Wang et al (2014) | China | Asia | Low or middle | Community | Whole population | Not reported | 162464 | Self-report | 40 | 17987 | 0.11 | Low | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ⁷ Gaulin et al (2019) | Canada | North America | High | Hospitals | All adults | 51.2 | 1316832 | Medical records and administrative database | 34 | 416282 | 0.32 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias | Rationale for exclusion |
|-------------------------------------|----------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|--|
| ⁸ Violan et al (2014) | Spain | Europe | High | Primary care | All adults | 47.4 | 1356761 | Medical records and administrative database | 146 | 645818 | 0.48 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ⁹ Nicholson et al (2019) | Canada | North America | High | Primary care | All adults | 52.3 | 367743 | Medical records and administrative database | 20 | 195838 | 0.53 | High | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ¹⁰ Bao et al (2019) | China | Asia | Low or middle | Community | Middle aged and older | 61.36 | 18137 | Self-report | 19 | 3773 | 0.21 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ¹¹ Fortin et al (2005) | Canada | North America | High | Primary care | All adults | 56.55 | 980 | Medical records and administrative database | 14 | 873 | 0.89 | Moderate | The studentized residual of this study is more than 2 standard deviations away from its expected value. |
| ¹² Prazeres et al (2015) | Portugal | Europe | High | Primary care | All adults | 56.3 | 1993 | Medical records and administrative database | 147 | 1449 | 0.73 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |
| ¹³ Lawson et al (2013) | UK | Europe | High | Community | All adults | 72.7 | 7054 | Medical records and administrative database | 40 | 1243 | 0.18 | Moderate | Irregular patterns found in compositional data (in scatter plot and Mahalanobis distance test)- low prevalence in studies with high mean participant age and a larger number of conditions |
| ¹⁴ Sullivan et al (2012) | USA | North America | High | Community | All adults | Not reported | 47178 | Medical records and administrative database | 259 | 19666 | 0.42 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias | Rationale for exclusion |
|--------------------------------------|-------------|---------------|----------------|--------------|-------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|---|
| ¹⁵ Peng et al (2020) | China | Asia | Low or middle | Community | Only older people | 71.6 | 1321 | Self-report | 15 | 589 | 0.45 | Moderate | Contributing to high levels of heterogeneity of effect sizes (in leave-one-out analysis) |
| ¹⁶ Excoffier et al (2018) | Switzerland | Europe | High | Primary care | All adults | 56.5 | 2904 | Medical records and administrative database | 75 | 1513 | 0.52 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |
| ¹⁷ Chung et al (2015) | Hong Kong | Asia | High | Community | All adults | Not reported | 25780 | Self-report | 46 | 3227 | 0.13 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |
| ¹⁸ Ki et al (2017) | South Korea | Asia | High | Community | All adults | 57.05 | 19942 | Medical records and administrative database | 66 | 5979 | 0.30 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |
| ¹⁹ Bobo et al (2016) | USA | North America | High | Community | Whole population | Not reported | 138858 | Self-report | 19 | 33682 | 0.24 | Moderate | Infrequent values in compositional categorical data (few studies focused on whole population) |
| ²⁰ Randall et al (2018) | Australia | Australasia | High | Community | Whole population | Not reported | 5437018 | Self-report | 30 | 660449 | 0.12 | Moderate | Infrequent values in compositional categorical data (few studies focused on whole population) |
| ²¹ Russell et al (2020) | New Zealand | Australasia | High | Community | Only children | Not reported | 3838 | Self-report | 7 | 374 | 0.10 | Moderate | Infrequent values in compositional categorical data (only one study focused on children population) |
| ²² Barnett et al (2012) | UK | Europe | High | Primary care | Whole population | Not reported | 1751841 | Medical records and administrative database | 40 | 406427 | 0.23 | Low | Infrequent values in compositional categorical data (few studies focused on whole population) |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias | Rationale for exclusion |
|--------------------------------------|---|---------------------|----------------|--------------|-------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|---|
| ²³ St Sauver et al (2015) | USA | North America | High | Primary care | Whole population | Not reported | 106061 | Medical records and administrative database | 20 | 34592 | 0.33 | Moderate | Infrequent values in compositional categorical data (few studies focused on whole population) |
| ²⁴ Vetrano et al (2016) | Denmark, Finland, Iceland, Italy, the Netherlands, Norway, United Kingdom, Czech Republic, France, Sweden and Germany, Canada | Multiple continents | High | Care homes | Only older people | 82.2 | 6903 | Medical records and administrative database | 13 | 5098 | 0.74 | Moderate | Infrequent values in compositional categorical data (only one study focused on care home) |

MM: Multimorbidity. No of participants: The total number of participants in the denominator for estimating prevalence in a study (which could be a subset in some included studies)

Table S4: Characteristics of 194 included studies

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-----------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ²⁵ Aarts et al (2012) | The Netherlands | Europe | High | Primary care | All adults | 55.4 | 1184 | Medical records and administrative database | 23 | 420 | 0.35 | Moderate |
| ²⁶ Aarts et al (2011a) | The Netherlands | Europe | High | Community | Middle aged and older | 70 | 15188 | Self-report | Not reported | 7729 | 0.51 | Moderate |
| ²⁷ Aarts et al (2011b) | The Netherlands | Europe | High | Primary care | All adults | 55.4 | 1763 | Medical records and administrative database | 23 | 985 | 0.56 | Moderate |
| ²⁸ Abizanda et al (2014) | Spain | Europe | High | Primary care | Only older people | 78.6 | 842 | Medical records and administrative database | 14 | 580 | 0.69 | Moderate |
| ²⁹ Agborsangaya et al (2012) | Canada | North America | High | Community | All adults | 46.6 | 4003 | Self-report | 16 | 919 | 0.23 | Moderate |
| ³⁰ Agborsangaya et al (2013) | Canada | North America | High | Community | All adults | 47.8 | 4803 | Self-report | 16 | 1729 | 0.36 | Moderate |
| ³¹ Agborsangaya et al (2014) | Canada | North America | High | Community | All adults | 47.7 | 4752 | Self-report | 16 | 1597 | 0.34 | Moderate |
| ³² Ahrenfeldt et al (2019) | Europe | Europe | High | Community | Middle aged and older | 66.25 | 244258 | Self-report | 10 | 90652 | 0.37 | Moderate |
| ³³ Alimohammadian et al (2017) | Iran | Asia | Low or middle | Community | Middle aged and older | Not reported | 49946 | Self-report | 8 | 10035 | 0.20 | Moderate |
| ³⁴ Angst et al (2002) | Switzerland | Europe | High | Primary care | All adults | Not reported | 591 | Medical records and administrative database | 10 | 201 | 0.34 | High |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|---|---------------------|----------------|--------------|-----------------------|---------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ³⁵ Appa et al (2014) | USA | North America | High | Community | All adults | 60.2 | 1997 | Self-report | 16 | 1417 | 0.71 | Moderate |
| ³⁶ Adams et al (2017) | USA | North America | High | Community | All adults | Not reported | 400000 | Self-report | 12 | 191600 | 0.48 | Moderate |
| ³⁷ Ahmadi et al (2016) | Iran | Asia | Low or middle | Community | Middle aged and older | 52.1 | 49946 | Self-report | 8 | 10035 | 0.20 | Moderate |
| ³⁸ Amaral et al (2018) | Brazil | South America | Low or middle | Community | Only older people | Not reported | 264 | Self-report | 8 | 175 | 0.66 | Moderate |
| ³⁹ An et al (2016) | South Korea | Asia | High | Community | Middle aged and older | 54.8 | 10118 | Self-report | 8 | 3228 | 0.32 | Moderate |
| ⁴⁰ Araujo et al (2018) | Brazil | South America | Low or middle | Community | All adults | Not reported | 4001 | Self-report | 12 | 1160 | 0.29 | Moderate |
| ⁴¹ Arnold-Reed et al (2018) | Australia | Australasia | High | Primary care | All adults | 38.2 | 4285 | Medical records and administrative database | 43 | 2269 | 0.53 | Moderate |
| ⁴² Arokiasamy et al (2015) | 6 low middle income countries (China, Ghana, India, Mexico, Russia, South Africa) | Multiple continents | Low or middle | Community | All adults | Not reported | 42236 | Self-report | 8 | 9250 | 0.22 | Moderate |
| ⁴³ Sinnige et al (2015) | The Netherlands | Europe | High | Primary care | Middle aged and older | 66.9 | 120480 | Medical records and administrative database | 29 | 74733 | 0.62 | Moderate |
| ⁴⁴ Zemedikun et al (2018) | UK | Europe | High | Community | Middle aged and older | Median age 58 | 502643 | Medical records and administrative database | 36 | 95710 | 0.19 | Moderate |
| ⁴⁵ Wensing et al (2001) | The Netherlands | Europe | High | Primary care | All adults | Not reported | 3867 | Self-report | 25 | 626 | 0.16 | Moderate |
| ⁴⁶ Mounce et al (2018) | UK | Europe | High | Community | Middle aged and older | Not reported | 4564 | Self-report | 15 | 1553 | 0.34 | Moderate |
| ⁴⁷ Taylor et al (2010) | Australia | Australasia | High | Community | All adults | Not reported | 3206 | Self-report | 7 | 547 | 0.17 | Low |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|---|---------------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁴⁸ Vancampfort et al (2019) | Six low and middle income countries (China, Ghana, India, Mexico, Russia, and South Africa) | Multiple continents | Low or middle | Community | Middle aged and older | 62.4 | 34129 | Self-report | 11 | 15529 | 0.46 | Moderate |
| ⁴⁹ Vancampfort et al (2018) | Six low and middle income countries (China, Ghana, India, Mexico, Russia, and South Africa) | Multiple continents | Low or middle | Community | Only older people | 72.6 | 14585 | Self-report | 11 | 8780 | 0.60 | Moderate |
| ⁵⁰ Aubert et al (2016) | Switzerland | Europe | High | Primary care | Middle aged and older | 63.5 | 1002 | Medical records and administrative database | 17 | 676 | 0.67 | Moderate |
| ⁵¹ Autenrieth et al (2013) | Germany | Europe | High | Community | Only older people | 75.7 | 1007 | Self-report | 13 | 658 | 0.65 | Moderate |
| ⁵² Bahler et al (2015) | Switzerland | Europe | High | Community | Only older people | 74.9 | 229493 | Medical records and administrative database | 22 | 175752 | 0.77 | Moderate |
| ⁵³ Vancampfort et al (2017) | 44 low and middle income countries | Multiple continents | Low or middle | Community | All adults | 38.3 | 194431 | Self-report | 11 | 27518 | 0.14 | Moderate |
| ⁵⁴ Banjare et al (2014) | India | Asia | Low or middle | Community | Only older people | Not reported | 310 | Self-report | 20 | 176 | 0.57 | Moderate |
| ⁵⁵ Barra et al (2015) | USA | North America | High | Community | All adults | 45.36 | 43079 | Self-report | Not reported | 22412 | 0.52 | Moderate |
| ⁵⁶ Bernard et al (2016) | Australia | Australasia | High | Hospitals | Only older people | 81.8 | 306 | Medical records and administrative database | 19 | 125 | 0.41 | High |
| ⁵⁷ Biswas et al (2019) | Bangladesh | Asia | Low or middle | Community | All adults | Not reported | 8763 | Self-report | 3 | 1078 | 0.12 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|---|---------------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁵⁸ Blakemore et al (2016) | UK | Europe | High | Primary care | Only older people | 75 | 4377 | Self-report | 24 | 2631 | 0.60 | Moderate |
| ⁵⁹ Blyth et al (2008) | Australia | Australasia | High | Community | Only older people | 76.9 | 1685 | Self-report | 18 | 920 | 0.55 | Moderate |
| ⁶⁰ Bowling et al (2019) | USA | North America | High | Community | Middle aged and older | 56.7 | 4217 | Self-report | 12 | 3053 | 0.72 | Moderate |
| ⁶¹ Britt et al (2008) | Australia | Australasia | High | Primary care | All adults | Not reported | 9156 | Medical records and administrative database | 18 | 3398 | 0.37 | Moderate |
| ⁶² Broeiro-Goncalves et al (2019) | Portugal | Europe | High | Hospitals | All adults | 59.8 | 800376 | Medical records and administrative database | 22 | 335357 | 0.42 | Moderate |
| ⁶³ Bruce et al (2010) | Canada | North America | High | Community | All adults | 37.8 | 453 | Self-report | 4 | 163 | 0.36 | High |
| ⁶⁴ Burgers et al (2010) | France, Germany, Canada, Australia, Netherlands, New Zealand, UK, USA | Multiple continents | High | Community | All adults | Not reported | 8973 | Self-report | 7 | 4037 | 0.45 | Moderate |
| ⁶⁵ Burke et al (2017) | US, Europe, Asia | Multiple continents | High | Community | Only older people | Not reported | 4668 | Self-report | 9 | 2165 | 0.46 | Moderate |
| ⁶⁶ Buurman et al (2016) | The Netherlands | Europe | High | Hospitals | Only older people | 78.2 | 639 | Medical records and administrative database | 35 | 440 | 0.69 | Moderate |
| ⁶⁷ Calderon-Larranaga et al (2017) | Sweden | Europe | High | Primary care | Only older people | 74.6 | 3363 | Self-report | 60 | 2980 | 0.89 | Moderate |
| ⁶⁸ Camargo-Casas et al (2018) | Colombia | South America | Low or middle | Community | Only older people | 71.1 | 2000 | Self-report | 12 | 808 | 0.40 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|---|---------------------|----------------|--------------|-----------------------|----------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁶⁹ Canevelli et al (2019) | Italy | Europe | High | Primary care | Only older people | 75.1 | 185 | Medical records and administrative database | 18 | 162 | 0.88 | High |
| ⁷⁰ Chamberlain et al (2020) | USA | North America | High | Community | All adults | Not reported | 198941 | Self-report | 21 | 78527 | 0.39 | Low |
| ⁷¹ Chen et al (2018) | China | Asia | Low or middle | Community | Only older people | Not reported | 30774 | Medical records and administrative database | 33 | 25101 | 0.82 | Low |
| ⁷² Chen et al (2018) | China | Asia | Low or middle | Community | Middle aged and older | Not reported | 3737 | Self-report | 16 | 1722 | 0.46 | Moderate |
| ⁷³ Cheung et al (2013) | Hong Kong (SAR of China) | Asia | High | Community | Middle aged and older | 71.3 | 1145 | Self-report | 18 | 654 | 0.57 | Moderate |
| ⁷⁴ Chu et al (2018) | Hong Kong (SAR of China) | Asia | High | Primary care | Middle aged and older | Not reported | 382 | Medical records and administrative database | 40 | 206 | 0.54 | Moderate |
| ⁷⁵ Chudasama et al (2019) | UK | Europe | High | Community | Middle aged and older | Median age:58 | 491939 | Medical records and administrative database | 36 | 96622 | 0.20 | Moderate |
| ⁷⁶ Cimarras-Otal et al (2014) | Spain | Europe | High | Community | All adults | Not reported | 22190 | Self-report | 20 | 7830 | 0.35 | Moderate |
| ⁷⁷ Chin et al (2016) | Hong Kong (SAR of China) | Asia | High | Primary care | All adults | Median age: 48 | 9259 | Self-report | 8 | 2350 | 0.25 | Moderate |
| ⁷⁸ Agrawal et al (2016) | India, China, Russia, Mexico, South Africa, Ghana | Multiple continents | Low or middle | Community | All adults | 57.8 | 40166 | Self-report | 9 | 9238 | 0.23 | Moderate |
| ⁷⁹ Gu et al (2018) | China | Asia | Low or middle | Community | Only older people | Not reported | 411 | Self-report | 13 | 232 | 0.56 | Moderate |
| ⁸⁰ Gunn et al (2012) | Australia | Australasia | High | Primary care | All adults | 50.89 | 6864 | Self-report | 12 | 2154 | 0.31 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-----------|---------------|----------------|--------------|-----------------------|----------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁸¹ Han et al (2013) | USA | North America | High | Primary care | Only older people | 76 | 159 | Medical records and administrative database | 18 | 117 | 0.74 | High |
| ⁸² Hanlon et al (2018) | UK | Europe | High | Community | All adults | Not reported | 493737 | Medical records and administrative database | 42 | 161576 | 0.33 | Low |
| ⁸³ Jantsch et al (2018) | Brazil | South America | Low or middle | Community | All adults | 42 | 3092 | Self-report | 11 | 912 | 0.29 | Moderate |
| ⁸⁴ John et al (2003) | USA | North America | High | Community | Only older people | 71.3 | 992 | Self-report | 11 | 732 | 0.74 | High |
| ⁸⁵ Johnson-Lawrence et al (2017) | USA | North America | High | Community | All adults | 49.9 | 115097 | Self-report | 9 | 27278 | 0.24 | Moderate |
| ⁸⁶ Johnston et al (2019) | UK | Europe | High | Community | All adults | 48 | 7184 | Self-report | Not reported | 388 | 0.05 | Moderate |
| ⁸⁷ Jones et al (2016) | USA | North America | High | Community | Only older people | Not reported | 6964 | Self-report | 10 | 4951 | 0.71 | Moderate |
| ⁸⁸ Jovic et al (2016) | Serbia | Europe | Low or middle | Community | All adults | 49.4 | 13103 | Self-report | 13 | 3522 | 0.27 | Moderate |
| ⁸⁹ Juul-Larsen et al (2020) | Denmark | Europe | High | Hospitals | Only older people | Median age: 78 | 369 | Self-report | 34 | 311 | 0.84 | Moderate |
| ⁹⁰ Hudon et al (2008) | Canada | North America | High | Community | All adults | Not reported | 16782 | Self-report | 25 | 5343 | 0.32 | Low |
| ⁹¹ Hussain et al (2015) | Indonesia | Asia | Low or middle | Community | Middle aged and older | Not reported | 9438 | Self-report | 12 | 3369 | 0.36 | Moderate |
| ⁹² Ie et al (2017) | USA | North America | High | Hospitals | Only older people | Not reported | 1084 | Medical records and administrative database | 24 | 1036 | 0.96 | High |
| ⁹³ Ishizaki et al (2019) | Japan | Asia | High | Community | Only older people | 76.9 | 2525 | Self-report | 9 | 1121 | 0.44 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁹⁴ Danon-Hersch et al (2012) | Switzerland | Europe | High | Community | Only older people | Not reported | 1283 | Self-report | 12 | 448 | 0.35 | Moderate |
| ⁹⁵ de Heer et al (2013) | USA | North America | High | Community | All adults | 47.72 | 1002 | Self-report | 19 | 378 | 0.38 | Moderate |
| ⁹⁶ Demirchyan et al (2013) | Armenia | Asia | Low or middle | Community | All adults | 58.8 | 721 | Self-report | Not reported | 564 | 0.78 | High |
| ⁹⁷ Fabbri et al (2015) | Italy | Europe | High | Community | Only older people | 73.6 | 1018 | Self-report | 15 | 458 | 0.45 | Moderate |
| ⁹⁸ Fillenbaum et al (2000) | USA | North America | High | Community | Only older people | 73.44 | 4034 | Self-report | 5 | 1181 | 0.29 | Moderate |
| ⁹⁹ Kaneko et al (2019) | Japan | Asia | High | Community | Only older people | Not reported | 253 | Self-report | Not reported | 135 | 0.53 | Moderate |
| ¹⁰⁰ Kang et al (2017) | South Korea | Asia | High | Primary care | All adults | 32.2 | 590 | Medical records and administrative database | 14 | 153 | 0.26 | Moderate |
| ¹⁰¹ Gandhi et al (2020) | USA | North America | High | Community | All adults | Not reported | 9499 | Self-report | 8 | 3379 | 0.36 | Moderate |
| ¹⁰² Costa et al (2018) | Brazil | South America | Low or middle | Community | Only older people | Not reported | 1451 | Self-report | 29 | 1343 | 0.93 | Moderate |
| ¹⁰³ Rizzuto et al (2017) | Sweden | Europe | High | Community | Only older people | Not reported | 1099 | Self-report | 36 | 774 | 0.70 | Moderate |
| ¹⁰⁴ Dhalwani et al (2017) | UK | Europe | High | Community | Middle aged and older | Not reported | 5476 | Self-report | 18 | 1156 | 0.21 | Moderate |
| ¹⁰⁵ Elixhauser et al (1998) | USA | North America | High | Hospitals | All adults | 57.1 | 1779167 | Medical records and administrative database | 30 | 619150 | 0.35 | Low |
| ¹⁰⁶ Fabbri et al (2015) | USA | North America | High | Hospitals | Only older people | 72.3 | 695 | Self-report | 15 | 440 | 0.63 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---------------------------------------|---|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁰⁷ Fortin et al (2014) | Canada | North America | High | Community | Middle aged and older | 57.8 | 1196 | Self-report | 14 | 599 | 0.50 | Moderate |
| ¹⁰⁸ Fuchs et al (1998) | Israel | Asia | High | Community | Only older people | Not reported | 1820 | Self-report | 14 | 1174 | 0.65 | Moderate |
| ¹⁰⁹ Galenkamp et al (2011) | The Netherlands | Europe | High | Community | Middle aged and older | 69.2 | 2046 | Self-report | 7 | 876 | 0.43 | High |
| ¹¹⁰ Galenkamp et al (2016) | Germany, UK, Italy, The Netherlands, Spain and Sweden | Europe | High | Community | Only older people | 74.2 | 2792 | Self-report | 8 | 1358 | 0.49 | Moderate |
| ¹¹¹ Gamma et al (2001) | Switzerland | Europe | High | Community | All adults | Not reported | 407 | Self-report | 14 | 53 | 0.13 | High |
| ¹¹² Ge et al (2018) | Singapore | Asia | High | Community | All adults | 51.4 | 1940 | Self-report | 17 | 715 | 0.37 | Moderate |
| ¹¹³ Ge et al (2019) | Singapore | Asia | High | Community | All adults | 51.3 | 1932 | Self-report | 17 | 564 | 0.29 | Moderate |
| ¹¹⁴ Gould et al (2016) | USA | North America | High | Community | Only older people | 74.82 | 4184 | Self-report | 7 | 2932 | 0.70 | Moderate |
| ¹¹⁵ Habib et al (2014) | Lebanon | Asia | Low or middle | Community | All adults | 46.6 | 2501 | Self-report | Not reported | 665 | 0.27 | Moderate |
| ¹¹⁶ Harrison et al (2017) | Australia | Australasia | High | Primary care | All adults | Not reported | 8707 | Medical records and administrative database | 28 | 2838 | 0.33 | Moderate |
| ¹¹⁷ Hayek et al (2017) | Israel | Asia | High | Community | All adults | 47.2 | 4325 | Self-report | 10 | 1579 | 0.37 | Moderate |
| ¹¹⁸ Henninger et al (2012) | USA | North America | High | Community | Only older people | 76 | 3212 | Self-report | 9 | 1753 | 0.55 | Moderate |
| ¹¹⁹ Hernandez et al (2019) | Ireland | Europe | High | Community | Middle aged and older | Not reported | 6101 | Self-report | 31 | 4468 | 0.73 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|---|---------------------|----------------|-----------|-----------------------|--------------|--------------------|-------------|---------------------------|----------------|--------------------|--------------|
| ¹²⁰ Ho et al (2014) | Singapore | Asia | High | Community | Middle aged and older | 66.15 | 1844 | Self-report | 12 | 830 | 0.45 | Moderate |
| ¹²¹ Khan et al (2019) | Bangladesh | Asia | Low or middle | Community | All adults | 58.6 | 12338 | Self-report | 6 | 1031 | 0.08 | Low |
| ¹²² Kiliari et al (2013) | Cyprus | Europe | High | Community | All adults | 53 | 465 | Self-report | Not reported | 132 | 0.28 | Moderate |
| ¹²³ King et al (2018) | USA | North America | High | Community | All adults | Not reported | 5541 | Self-report | 11 | 3342 | 0.60 | Moderate |
| ¹²⁴ Kingston et al (2018) | UK | Europe | High | Community | All adults | Not reported | 9723900 | Self-report | 12 | 5250906 | 0.54 | High |
| ¹²⁵ Koyanagi et al (2018) | China, Ghana, India, Mexico, Russia, and South Africa | Multiple continents | Low or middle | Community | Middle aged and older | 62.1 | 32715 | Self-report | 10 | 16324 | 0.50 | Moderate |
| ¹²⁶ Kriegsman et al (2004) | The Netherlands | Europe | High | Community | Middle aged and older | 69.2 | 2489 | Self-report | 7 | 519 | 0.21 | Moderate |
| ¹²⁷ Kristensen et al (2019) | Germany | Europe | High | Community | Middle aged and older | 63.47 | 19605 | Self-report | 13 | 12600 | 0.64 | Moderate |
| ¹²⁸ Kristensen et al (2019) | Germany | Europe | High | Community | Middle aged and older | 64.37 | 7604 | Self-report | 13 | 5140 | 0.68 | Moderate |
| ¹²⁹ Kunna et al (2017) | China, Ghana | Multiple continents | Low or middle | Community | Middle aged and older | Not reported | 15864 | Self-report | 7 | 4731 | 0.30 | Low |
| ¹³⁰ Kuwornu et al (2014) | Canada | North America | High | Community | All adults | 51.05 | 3284 | Self-report | 15 | 1143 | 0.35 | Moderate |
| ¹³¹ Lai et al (2019) | Hong Kong (SAR of China) | Asia | High | Community | All adults | Not reported | 69636 | Self-report | 14 | 3898 | 0.06 | Moderate |
| ¹³² Lai et al (2018) | Hong Kong (SAR of China) | Asia | High | Community | All adults | Not reported | 300 | Self-report | 11 | 48 | 0.16 | Moderate |
| ¹³³ Laires et al (2019) | Portugal | Europe | High | Community | All adults | Not reported | 15196 | Self-report | 13 | 6671 | 0.44 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-----------------------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹³⁴ Lang et al (2015) | USA | North America | High | Community | Middle aged and older | 53.4 | 3058 | Self-report | 6 | 948 | 0.31 | Moderate |
| ¹³⁵ Le Cossec et al (2016) | France | Europe | High | Community | Middle aged and older | 70 | 15325 | Self-report | 4 | 3528 | 0.23 | Moderate |
| ¹³⁶ Lee et al (2007) | USA | North America | High | Hospitals | Middle aged and older | Not reported | 741847 | Medical records and administrative database | 11 | 302792 | 0.41 | Low |
| ¹³⁷ Lee et al (2018) | Taiwan | Asia | High | Community | Only older people | Not reported | 20898 | Medical records and administrative database | Not reported | 4234 | 0.20 | High |
| ¹³⁸ Li et al (2016) | UK | Europe | High | Primary care | All adults | Not reported | 27806 | Self-report | 12 | 10332 | 0.37 | Moderate |
| ¹³⁹ Li et al (2019) | USA | North America | High | Community | Middle aged and older | 67.4 | 14996 | Self-report | 8 | 9805 | 0.65 | Moderate |
| ¹⁴⁰ Lujic et al (2017) | Australia | Australasia | High | Community | Middle aged and older | 70.2 | 90352 | Self-report | 8 | 33792 | 0.37 | Moderate |
| ¹⁴¹ Lupianez-Villanueva et al (2018) | 14 European countries | Europe | High | Community | All adults | Not reported | 14000 | Self-report | 13 | 3416 | 0.24 | Moderate |
| ¹⁴² Zhou et al (2018) | Bangladesh, India and China | Asia | Low or middle | Community | All adults | Not reported | 18696 | Self-report | 9 | 3512 | 0.19 | Moderate |
| ¹⁴³ Zhang et al (2019) | China | Asia | Low or middle | Community | Only older people | 70.5 | 11707 | Self-report | 11 | 5104 | 0.44 | Moderate |
| ¹⁴⁴ Wong et al (2010) | Canada | North America | High | Community | Only older people | Not reported | 740 | Self-report | 7 | 489 | 0.66 | Moderate |
| ¹⁴⁵ Weimann et al (2016) | South Africa | Africa | Low or middle | Community | All adults | 34 | 18526 | Self-report | 4 | 5057 | 0.27 | Moderate |
| ¹⁴⁶ Wang et al (2017) | Australia | Australasia | High | Community | All adults | 44 | 8820 | Self-report | 8 | 2539 | 0.29 | Moderate |
| ¹⁴⁷ Wang et al (2019) | South Africa | Africa | Low or middle | Community | Only older people | Not reported | 2627 | Self-report | 5 | 439 | 0.17 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-----------------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁴⁸ Wade et al (2019) | New Zealand | Australasia | High | Community | All adults | 59.05 | 7654 | Self-report | 12 | 2786 | 0.36 | Moderate |
| ¹⁴⁹ Maciejewski et al (2019) | USA | North America | High | Community | Only older people | 77.1 | 20124230 | Medical records and administrative database | 19 | 14425446 | 0.72 | Moderate |
| ¹⁵⁰ Marengoni et al (2016) | Sweden | Europe | High | Community | Only older people | 74.4 | 3155 | Medical records and administrative database | 14 | 1654 | 0.52 | Moderate |
| ¹⁵¹ Marengoni et al (2009) | Sweden | Europe | High | Community | Only older people | Not reported | 1099 | Self-report | 22 | 575 | 0.52 | Moderate |
| ¹⁵² Marques et al (2018) | 13 European countries | Europe | High | Community | All adults | 50.2 | 32931 | Self-report | 6 | 7113 | 0.22 | Moderate |
| ¹⁵³ Mavaddat et al (2014) | UK | Europe | High | Primary care | Middle aged and older | 58.7 | 11439 | Self-report | 6 | 1006 | 0.09 | Moderate |
| ¹⁵⁴ McDaid et al (2013) | Ireland | Europe | High | Community | Middle aged and older | Not reported | 6018 | Self-report | 8 | 733 | 0.12 | High |
| ¹⁵⁵ Melis et al (2014) | Sweden | Europe | High | Hospitals | Only older people | 83.75 | 390 | Medical records and administrative database | 39 | 213 | 0.55 | Moderate |
| ¹⁵⁶ Min et al (2007) | USA | North America | High | Community | Only older people | 81 | 372 | Self-report | 9 | 230 | 0.62 | High |
| ¹⁵⁷ Momtaz et al (2010) | Malaysia | Asia | High | Community | Only older people | 69.26 | 385 | Self-report | 16 | 165 | 0.43 | Moderate |
| ¹⁵⁸ Mondor et al (2018) | Canada | North America | High | Community | All adults | Not reported | 27195 | Medical records and administrative database | 17 | 11390 | 0.42 | Moderate |
| ¹⁵⁹ Muggah et al (2012) | Canada | North America | High | Community | All adults | Not reported | 28450000 | Medical records and administrative database | 9 | 4523550 | 0.16 | Moderate |
| ¹⁶⁰ Nagel et al (2008) | Germany | Europe | High | Community | Middle aged and older | 56.5 | 13781 | Self-report | 15 | 9275 | 0.67 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|--------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁶¹ Niedzwiedz et al (2019) | USA | North America | High | Community | Middle aged and older | 67.2 | 2272 | Self-report | 8 | 1491 | 0.66 | Moderate |
| ¹⁶² Nunes et al (2016) | Brazil | South America | Low or middle | Community | All adults | 45.75 | 2927 | Self-report | 11 | 852 | 0.29 | Moderate |
| ¹⁶³ Nunes et al (2017) | Brazil | South America | Low or middle | Community | All adults | 43.7 | 60202 | Self-report | 22 | 13365 | 0.22 | Moderate |
| ¹⁶⁴ Nunes et al (2015) | Brazil | South America | Low or middle | Community | Only older people | Not reported | 1593 | Self-report | 17 | 1295 | 0.81 | Moderate |
| ¹⁶⁵ Olaya et al (2017) | Spain | Europe | High | Community | Only older people | 71.75 | 2113 | Self-report | 7 | 1088 | 0.51 | Moderate |
| ¹⁶⁶ Olivares et al (2017) | Argentina | South America | High | Community | All adults | 43 | 1044 | Self-report | Not reported | 346 | 0.33 | Moderate |
| ¹⁶⁷ Park et al (2018) | South Korea | Asia | High | Community | Middle aged and older | 62.7 | 5996 | Self-report | 25 | 1607 | 0.27 | Moderate |
| ¹⁶⁸ Patel et al (2006) | Mexico | South America | Low or middle | Community | Middle aged and older | 73 | 7852 | Self-report | 5 | 1833 | 0.23 | Moderate |
| ¹⁶⁹ Pati et al (2016) | India | Asia | Low or middle | Community | All adults | 44.96 | 103 | Self-report | 18 | 24 | 0.23 | Moderate |
| ¹⁷⁰ Pati et al (2019) | India | Asia | Low or middle | Primary care | All adults | 44 | 1649 | Self-report | 21 | 567 | 0.34 | Moderate |
| ¹⁷¹ Pati et al (2017) | India | Asia | Low or middle | Primary care | All adults | 44 | 1649 | Self-report | 21 | 467 | 0.28 | Moderate |
| ¹⁷² Payne et al (2013) | UK | Europe | High | Primary care | All adults | 49 | 180815 | Medical records and administrative database | 40 | 54945 | 0.30 | Moderate |
| ¹⁷³ Perez et al (2020) | Sweden | Europe | High | Community | Only older people | 72.8 | 2596 | Self-report | 60 | 2213 | 0.85 | Moderate |
| ¹⁷⁴ Petersen et al (2019) | South Africa | Africa | Low or middle | Primary care | All adults | Not reported | 2549 | Self-report | Not reported | 893 | 0.35 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|--|---------------|----------------|--------------|-----------------------|----------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁷⁵ Pfortmueller et al (2013) | Switzerland | Europe | High | Hospitals | All adults | Median age: 28 | 3170 | Medical records and administrative database | 18 | 1183 | 0.37 | High |
| ¹⁷⁶ Pressley et al (1999) | USA | North America | High | Hospitals | Only older people | Not reported | 5934 | Medical records and administrative database | Not reported | 3534 | 0.60 | Moderate |
| ¹⁷⁷ Prior et al (2016) | Denmark | Europe | High | Community | All adults | Not reported | 118410 | Self-report | 39 | 33937 | 0.29 | Moderate |
| ¹⁷⁸ Ribeiro et al (2018) | Brazil | South America | High | Community | Only older people | 70 | 820 | Self-report | 8 | 270 | 0.33 | Moderate |
| ¹⁷⁹ Ruel et al (2014) | Australia | Australasia | High | Community | All adults | 50 | 1854 | Self-report | 8 | 585 | 0.32 | Moderate |
| ¹⁸⁰ Ruel et al (2014) | China | Asia | Low or middle | Community | All adults | 49 | 1020 | Self-report | 11 | 346 | 0.34 | Moderate |
| ¹⁸¹ Ryan et al (2018) | Ireland | Europe | High | Community | Middle aged and older | Not reported | 4823 | Self-report | 16 | 2588 | 0.54 | Moderate |
| ¹⁸² Schmidt et al (2016) | Austria, Belgium, Denmark, France, Germany, Italy, the Netherlands, Spain, Sweden, and Switzerland | Europe | High | Community | Only older people | Not reported | 56609 | Self-report | 11 | 13794 | 0.24 | Moderate |
| ¹⁸³ Schottker et al (2016) | Germany | Europe | High | Primary care | Middle aged and older | Median age:70 | 2547 | Medical records and administrative database | 14 | 251 | 0.10 | Moderate |
| ¹⁸⁴ Seo et al (2017) | South Korea | Asia | High | Community | Middle aged and older | Not reported | 156747 | Self-report | 15 | 42006 | 0.27 | Moderate |
| ¹⁸⁵ She et al (2019) | China | Asia | Low or middle | Hospitals | Only older people | 68.9 | 1497 | Self-report | 22 | 1255 | 0.84 | Moderate |
| ¹⁸⁶ Singh et al (2019) | India | Asia | Low or middle | Community | All adults | 41 | 16287 | Self-report | 5 | 1531 | 0.09 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|---|---------------------|----------------|--------------|-----------------------|----------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁸⁷ Stepanova et al (2015) | USA | North America | High | Community | All adults | 34.7 | 26225 | Self-report | 13 | 9992 | 0.38 | High |
| ¹⁸⁸ Stickley et al (2020) | USA | North America | High | Community | All adults | 44.9 | 15311 | Self-report | 9 | 3996 | 0.26 | High |
| ¹⁸⁹ Streit et al (2014) | Switzerland | Europe | High | Primary care | Middle aged and older | 63.5 | 1002 | Medical records and administrative database | 17 | 676 | 0.67 | Moderate |
| ¹⁹⁰ Stubbs et al (2018) | China, Ghana, India, Mexico, Russia, South Africa | Multiple continents | Low or middle | Community | Middle aged and older | 62.4 | 34129 | Self-report | 13 | 19317 | 0.57 | Moderate |
| ¹⁹¹ Su et al (2016) | China | Asia | Low or middle | Community | Only older people | Not reported | 2058 | Self-report | 10 | 1012 | 0.49 | Moderate |
| ¹⁹² Sundstrup et al (2017) | USA | North America | High | Community | All adults | 43.5 | 10427 | Self-report | 8 | 2489 | 0.24 | High |
| ¹⁹³ Takahashi et al (2016) | USA | North America | High | Hospitals | All adults | 57 | 6402 | Medical records and administrative database | Not reported | 3140 | 0.49 | High |
| ¹⁹⁴ Tinetti et al (2011) | USA | North America | High | Community | Only older people | 72.6 | 5298 | Self-report | 5 | 1200 | 0.23 | High |
| ¹⁹⁵ Troelstra et al (2020) | The Netherlands | Europe | High | Community | All adults | Not reported | 604 | Self-report | 26 | 321 | 0.53 | High |
| ¹⁹⁶ van Zon et al (2020) | USA | North America | High | Community | Middle aged and older | 53.8 | 10719 | Self-report | 8 | 2390 | 0.22 | Moderate |
| ¹⁹⁷ Vancampfort et al (2017) | China, Ghana, India, Mexico, Russia, and South Africa | Multiple continents | Low or middle | Community | All adults | Median age: 62 | 32585 | Self-report | 11 | 14524 | 0.45 | Moderate |
| ¹⁹⁸ Vassilaki et al (2015) | USA | North America | High | Primary care | Only older people | 78.5 | 2176 | Medical records and administrative database | 17 | 1884 | 0.87 | Moderate |
| ¹⁹⁹ Vassilaki et al (2016) | USA | North America | High | Primary care | Only older people | 79 | 1449 | Medical records and administrative database | 17 | 1237 | 0.85 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|-----------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ²⁰⁰ Villarreal et al (2015) | Panama | South America | High | Primary care | Only older people | 78.2 | 304 | Self-report | 7 | 227 | 0.75 | Moderate |
| ²⁰¹ Violan et al (2019) | Spain | Europe | High | Primary care | Only older people | 75.4 | 916619 | Medical records and administrative database | 60 | 853085 | 0.93 | Moderate |
| ²⁰² von Strauss et al (2000) | Sweden | Europe | High | Community | Only older people | Not reported | 502 | Self-report | 15 | 155 | 0.31 | Moderate |
| ²⁰³ Vos et al (2013) | The Netherlands | Europe | High | Community | Only older people | 71.9 | 315 | Self-report | 21 | 202 | 0.64 | Moderate |
| ²⁰⁴ Vu et al (2019) | Vietnam | Asia | Low or middle | Hospitals | Only older people | 71.9 | 405 | Medical records and administrative database | Not reported | 146 | 0.36 | High |
| ²⁰⁵ Wang et al (2018) | USA | North America | High | Community | All adults | 47 | 3086 | Self-report | 20 | 1109 | 0.36 | Moderate |
| ²⁰⁶ Wang et al (2017) | China | Asia | Low or middle | Community | Only older people | 69.24 | 2705 | Self-report | 17 | 1230 | 0.45 | Moderate |
| ²⁰⁷ Wijers et al (2019) | Spain | Europe | High | Community | Middle aged and older | 74.2 | 707 | Self-report | 21 | 491 | 0.69 | Moderate |
| ²⁰⁸ Williams et al (2016) | USA | North America | High | Community | All adults | Not reported | 23789 | Self-report | 9 | 9213 | 0.39 | Moderate |
| ²⁰⁹ Woldeamayrat et al (2018) | Ethiopia | Africa | Low or middle | Primary care | All adults | Not reported | 411 | Self-report | 18 | 73 | 0.18 | Moderate |
| ²¹⁰ Yao et al (2020) | China | Asia | Low or middle | Community | Middle aged and older | 57.7 | 10084 | Self-report | 15 | 3243 | 0.32 | Moderate |
| ²¹¹ Yorke et al (2017) | USA | North America | High | Community | Middle aged and older | 66.6 | 5877 | Self-report | 7 | 3391 | 0.58 | Moderate |
| ²¹² You et al (2019) | China | Asia | Low or middle | Community | Only older people | 72 | 5296 | Self-report | 27 | 2201 | 0.42 | Moderate |
| ²¹³ Zhang et al (2020) | China | Asia | Low or middle | Community | Only older people | 74.14 | 4348 | Self-report | 15 | 2338 | 0.54 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|-------------------------------------|--------------------------|---------------|----------------|--------------|-------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ²¹⁴ Khanam et al (2011) | Bangladesh | Asia | Low or middle | Community | Only older people | 69.5 | 452 | Medical records and administrative database | 9 | 243 | 0.54 | Moderate |
| ²¹⁵ Cornell et al (2009) | USA | North America | High | Primary care | All adults | 62.4 | 1645314 | Medical records and administrative database | 45 | 1327382 | 0.81 | Moderate |
| ²¹⁶ Cassell et al (2018) | UK | Europe | High | Primary care | All adults | Not reported | 403985 | Medical records and administrative database | 36 | 109884 | 0.27 | Moderate |
| ²¹⁷ Wong et al (2019) | Hong Kong (SAR of China) | Asia | High | Community | All adults | 45.67 | 1014 | Self-report | 5 | 124 | 0.12 | Moderate |
| ²¹⁸ Puth et al (2017) | Germany | Europe | High | Community | All adults | Not reported | 19294 | Self-report | 17 | 7640 | 0.40 | Moderate |

MM: Multimorbidity. No of participants is the total number of participants in the denominator for estimating prevalence in a study (which could be a subset in some included studies)

Table S5: Associations between predictors

| | Mean age (lm) Unadjusted coefficient estimates | No of conditions (nb) Unadjusted incident rate ratio |
|-----------------------|--|--|
| Mean age | | 1.0 (1.0-1.0) |
| Source | | |
| Self-report | 59.5 (intercept) | Ref |
| Database | 7.2 (1.7-12.7)* | 1.8 (1.5-2.2)*** |
| Continent | | |
| Europe | 66.8 (62.7-70.9) (intercept) | Ref |
| North America | -7.0 (-12.8 to -1.1)* | 0.6 (0.5-0.8)*** |
| Australasia | -8.0 (-17.5-1.6) | 0.8 (0.6-1.1) |
| Asia | -8.9 (-15.1 to -2.7)** | 0.7 (0.6-0.8)*** |
| South America | -8.5 (-18.1-1.1) | 0.6 (0.4-0.9)** |
| Africa | -32.8 (-57.7 to -8.0)** | 0.4 (0.2-0.8)* |
| Multiple continents | -7.6 (-18.4-3.2) | 0.5 (0.3-0.7)*** |
| Setting | | |
| Community | 59.8 (intercept) | Ref |
| Primary care | 2.6 (-3.3-8.6) | 1.7 (1.4-2.1)*** |
| Hospitals | 10.2 (1.5-19.0)* | 1.8 (1.3-2.4)*** |
| Study population | | |
| All adults | 48.2 (intercept) | Ref |
| Middle-aged and older | 15.5 (12.8-18.1)*** | 0.9 (0.7-1.1) |
| Only older people | 26.3 (23.8-28.8)*** | 1.14 (1.0-1.4) |

*<0.05 **<0.01 ***<0.001

Ref: Reference category. lm: Linear regression. nb: Negative binomial regression

Table S6: Risk of bias assessment of included studies

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{1.} Aarts et al (2012) | Moderate | Moderate | Moderate | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{2.} Aarts et al (2011) | Low | High | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | No |
| ^{3.} Aarts et al (2011) | Moderate | Moderate | Moderate | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{4.} Abizanda et al (2014) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{5.} Agborsangaya et al (2012) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{6.} Agborsangaya et al (2013) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{7.} Agborsangaya et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{8.} Ahrenfeldt et al (2019) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | No |
| ^{9.} Alimohammadian et al (2017) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | Yes |
| ^{10.} Angst et al (2002) | Moderate | Moderate | Moderate | High | Low | High | High | Unclear | High | No |
| ^{11.} Appa et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{12.} Adams et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{13.} Ahmadi et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{14.} Amaral et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{15.} An et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{16.} Araujo et al (2018) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{17.} Arnold-Reed et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{18.} Arokiasamy et al (2015) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{19.} Sinnige et al (2015) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{20.} Zemedikun et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{21.} Wensing et al (2001) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Unclear | Moderate | Yes |
| ^{22.} Mounce et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{23.} Taylor et al (2010) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Low | Yes |
| ^{24.} Vancampfort et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{25.} Vancampfort et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{26.} Aubert et al (2016) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{27.} Autenrieth et al (2013) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{28.} Bahler et al (2015) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{29.} Vancampfort et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{30.} Banjare et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{31.} Barra et al (2015) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{32.} Bernard et al (2016) | High | Moderate | High | High | Moderate | Low | Moderate | Low | High | No |
| ^{33.} Biswas et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{34.} Blakemore et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{35.} Blyth et al (2008) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{36.} Bowling et al (2019) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{37.} Britt et al (2008) | Low | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{38.} Broeiro-Goncalves (2019) | Low | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{39.} Bruce et al (2010) | High | Moderate | Moderate | High | Low | High | Moderate | Unclear | High | No |
| ^{40.} Burgers et al (2010) | Low | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{41.} Burke et al (2017) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{42.} Buurman et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{43.} Calderon-Larranaga et al (2017) | Moderate | Moderate | Moderate | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{44.} Camargo-Casas et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{45.} Canevelli et al (2019) | High | High | High | High | Moderate | High | Moderate | Low | High | Yes |
| ^{46.} Chamberlain et al (2020) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Low | Yes |
| ^{47.} Chen et al (2018) | Low | Moderate | High | High | Low | Low | Moderate | Low | Low | Yes |
| ^{48.} Chen et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{49.} Cheung et al (2013) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{50.} Chu et al (2018) | Low | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{51.} Chudasama et al (2019) | Moderate | Moderate | Low | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{52.} Cimarras-Otal et al (2014) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{53.} Chin et al (2016) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{54.} Agrawal et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{55.} Gu et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{56.} Gunn et al (2012) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{57.} Han et al (2013) | High | High | Moderate | High | Moderate | High | Moderate | Unclear | High | No |
| ^{58.} Hanlon et al (2018) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Low | Yes |
| ^{59.} Jantsch et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{60.} John et al (2003) | Moderate | High | Moderate | High | Low | High | Moderate | Low | High | No |
| ^{61.} Johnson-Lawrence et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{62.} Johnston et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{63.} Jones et al (2016) | Low | Moderate | Moderate | High | Low | Low | Moderate | Unclear | Moderate | Yes |
| ^{64.} Jovic et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{65.} Juul-Larsen et al (2020) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{66.} Hudon et al (2008) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Low | Yes |
| ^{67.} Hussain et al (2015) | Moderate | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{68.} Ie et al (2017) | High | High | Moderate | High | Moderate | Low | Moderate | Low | High | Yes |
| ^{69.} Ishizaki et al (2019) | Moderate | Moderate | Low | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{70.} Danon-Hersch et al (2012) | Moderate | High | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{71.} de Heer et al (2013) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{72.} Demirchyan et al (2013) | High | Moderate | Low | High | Moderate | High | Moderate | Low | High | No |
| ^{73.} Fabbri et al (2015) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{74.} Fillenbaum et al (2000) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{75.} Kaneko et al (2019) | Moderate | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | No |
| ^{76.} Kang et al (2017) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{77.} Gandhi et al (2020) | Moderate | Moderate | Moderate | High | High | High | Moderate | Low | Moderate | Yes |
| ^{78.} Costa et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{79.} Rizzuto et al (2017) | High | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{80.} Dhalwani et al (2017) | Moderate | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{81.} Elixhauser et al (1998) | Low | Moderate | High | High | Low | Low | Moderate | Unclear | Low | Yes |
| ^{82.} Fabbri et al (2015) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{83.} Fortin et al (2014) | Low | Moderate | Low | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{84.} Fuchs et al (1998) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | No |
| ^{85.} Galenkamp et al (2011) | Low | Moderate | High | High | Moderate | High | Moderate | Unclear | High | No |
| ^{86.} Galenkamp et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{87.} Gamma et al (2001) | High | Moderate | High | High | Moderate | High | Moderate | Unclear | High | No |
| ^{88.} Ge et al (2018) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{89.} Ge et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{90.} Gould et al (2016) | Moderate | Moderate | Moderate | High | High | High | Moderate | Unclear | Moderate | Yes |
| ^{91.} Habib et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{92.} Harrison et al (2017) | Low | Moderate | High | High | Moderate | Low | Moderate | Unclear | Moderate | No |
| ^{93.} Hayek et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{94.} Henninger et al (2012) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{95.} Hernandez et al (2019) | Moderate | Moderate | Moderate | High | High | High | Moderate | Unclear | Moderate | Yes |
| ^{96.} Ho et al (2014) | Moderate | Moderate | High | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{97.} Khan et al (2019) | Low | Moderate | Low | High | Low | High | Moderate | Low | Low | Yes |
| ^{98.} Kiliari et al (2013) | High | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | No |
| ^{99.} King et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{100.} Kingston et al (2018) | Low | Moderate | High | High | Moderate | High | Moderate | Unclear | High | Yes |
| ^{101.} Koyanagi et al (2018) | Low | Moderate | Moderate | High | Moderate | Low | High | Low | Moderate | Yes |
| ^{102.} Kriegsman et al (2004) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{103.} Kristensen et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{104.} Kristensen et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{105.} Kunna et al (2017) | Low | Moderate | Low | High | Moderate | Low | High | Low | Low | Yes |
| ^{106.} Kuwornu et al (2014) | Moderate | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{107.} Lai et al (2019) | Low | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{108.} Lai et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{109.} Laires et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{110.} Lang et al (2015) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{111.} Le Cossec et al (2016) | Low | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{112.} Lee et al (2007) | Low | Moderate | High | High | Low | Low | Moderate | Low | Low | Yes |
| ^{113.} Lee et al (2018) | Low | Moderate | High | High | High | Low | Moderate | Unclear | High | No |
| ^{114.} Li et al (2016) | Low | Low | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{115.} Li et al (2019) | Low | Moderate | Low | High | Moderate | Moderate | Moderate | Low | Moderate | No |
| ^{116.} Lujic et al (2017) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | Yes |
| ^{117.} LupianezUnclearVillanueva et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{118.} Zhou et al (2018) | Moderate | Moderate | Moderate | High | Moderate | Low | High | Low | Moderate | Yes |
| ^{119.} Zhang et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{120.} Wong et al (2010) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{121.} Weimann et al (2016) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{122.} Wang et al (2017) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{123.} Wang et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{124.} Wade et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{125.} Maciejewski et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{126.} Marengoni et al (2016) | Moderate | Moderate | High | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{127.} Marengoni et al (2009) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{128.} Marques et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{129.} Mavaddat et al (2014) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{130.} McDaid et al (2013) | Low | Moderate | High | High | Moderate | High | Moderate | Low | High | Yes |
| ^{131.} Melis et al (2014) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{132.} Min et al (2007) | High | Moderate | High | High | Moderate | High | Moderate | Unclear | High | Yes |
| ^{133.} Momtaz et al (2010) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{134.} Mondor et al (2018) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{135.} Muggah et al (2012) | Low | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | No |
| ^{136.} Nagel et al (2008) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{137.} Niedzwiedz et al (2019) | Moderate | Moderate | High | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{138.} Nunes et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{139.} Nunes et al (2017) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | Yes |
| ^{140.} Nunes et al (2015) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{141.} Olaya et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{142.} Olivares et al (2017) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{143.} Park et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{144.} Patel et al (2006) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{145.} Pati et al (2016) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{146.} Pati et al (2019) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{147.} Pati et al (2017) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Unclear | Moderate | Yes |
| ^{148.} Payne et al (2013) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{149.} Perez et al (2020) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{150.} Petersen et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{151.} Pfortmueller et al (2013) | Moderate | Moderate | High | High | High | High | Moderate | Unclear | High | No |
| ^{152.} Pressley et al (1999) | Low | Moderate | High | High | Moderate | Low | Moderate | Unclear | Moderate | No |
| ^{153.} Prior et al (2016) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{154.} Ribeiro et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{155.} Ruel et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{156.} Ruel et al (2014) | Moderate | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | Yes |
| ^{157.} Ryan et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{158.} Schmidt et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{159.} Schottker et al (2016) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{160.} Seo et al (2017) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | No |
| ^{161.} She et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{162.} Singh et al (2019) | Low | Moderate | Moderate | High | Low | Low | Moderate | Unclear | Moderate | Yes |
| ^{163.} Stepanova et al (2015) | Low | High | High | High | High | High | High | Unclear | High | Yes |
| ^{164.} Stickley et al (2020) | Low | Moderate | High | High | Moderate | High | Moderate | Low | High | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{165.} Streit et al (2014) | Moderate | Moderate | Moderate | High | High | High | Moderate | Unclear | Moderate | Yes |
| ^{166.} Stubbs et al (2018) | Low | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{167.} Su et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{168.} Sundstrup et al (2017) | Low | Moderate | High | High | Moderate | High | Moderate | Unclear | High | Yes |
| ^{169.} Takahashi et al (2016) | Moderate | Moderate | High | High | High | Low | Moderate | Low | High | No |
| ^{170.} Tinetti et al (2011) | Low | Moderate | High | High | High | High | Moderate | Unclear | High | No |
| ^{171.} Troelstra et al (2020) | High | Moderate | High | High | Moderate | Low | Moderate | Unclear | High | Yes |
| ^{172.} van Zon et al (2020) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{173.} Vancampfort et al (2017) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{174.} Vassilaki et al (2015) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{175.} Vassilaki et al (2016) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{176.} Villarreal et al (2015) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{177.} Violan et al (2019) | Low | Moderate | Moderate | High | High | Low | Moderate | Low | Moderate | Yes |
| ^{178.} von Strauss et al (2000) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | No |
| ^{179.} Vos et al (2013) | Moderate | Moderate | High | High | Moderate | High | Moderate | Low | Moderate | No |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{180.} Vu et al (2019) | High | Moderate | High | High | Moderate | High | Moderate | Low | High | No |
| ^{181.} Wang et al (2018) | Moderate | Moderate | Low | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{182.} Wang et al (2017) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{183.} Wijers et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{184.} Williams et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{185.} Woldeamayrat et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{186.} Yao et al (2020) | Moderate | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{187.} Yorke et al (2017) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{188.} You et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{189.} Zhang et al (2020) | Moderate | Moderate | Low | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{190.} Khanam et al (2011) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{191.} Cornell et al (2009) | Low | Moderate | High | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{192.} Cassell et al (2018) | Low | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | No |
| ^{193.} Wong et al (2019) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{194.} Puth et al (2017) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |

Table S7: Output of adjusted meta-analytic model based on 218 studies


| | Pooled prevalence of multimorbidity of each subgroup (% , 95% CI) | Meta-regression Unadjusted Odds Ratio (95% CI) | Meta-regression Adjusted Odds Ratio (95% CI) R ² 45.1% | FMI |
|---------------------|---|--|---|------|
| Group of mean age | | R ² 30.9% | | |
| <59 | 29.3 (26.2-32.7) | Ref | Ref | Ref |
| 59-73 | 47.0 (41.3-52.7) | 2.1 (1.6-2.8)*** | 2.5 (2.0-3.2)*** | 0.3 |
| ≥74 | 66.1 (60.5-71.2) | 4.7 (3.4-6.4)*** | 4.4 (3.3-5.9)*** | 0.2 |
| No of conditions | | R ² 6.2% | | |
| <9 | 30.8 (26.0-35.9) | Ref | Ref | Ref |
| 9-19 | 45.0 (40.8-49.3) | 1.8 (1.3-2.6)*** | 1.8 (1.4-2.3)*** | 0.1 |
| 20-43 | 44.3 (35.7-53.3) | 1.8 (1.2-2.7)** | 1.8 (1.2-2.5)** | 0.2 |
| ≥44 | 51.6 (32.3-70.4) | 2.4 (1.3-4.3)** | 2.5 (1.5-4.0)*** | 0.2 |
| Setting | | R ² 3.3% | | |
| Community | 38.2 (34.9-41.7) | Ref | Ref | Ref |
| Primary care | 50.6 (41.2-59.9) | 1.7 (1.2-2.4)** | 1.8 (1.3-2.6)*** | 0.2 |
| Hospital | 47.1 (31.9-63.0) | 1.4 (0.9-2.4) | 1.0 (0.7-1.6) | 0.09 |
| Care home | 73.9 (72.8-74.9) | 4.6 (0.6-34.5) | 1.8 (0.3-9.2) | 0.03 |
| Source | | R ² 2.7% | | |
| Self-report | 38.6 (34.8-42.4) | Ref | Ref | Ref |
| Database | 48.9 (42.3-55.5) | 1.5 (1.1-2.1)** | 0.8 (0.6-1.1) | 0.1 |
| Continent | | R ² 5.3% | | |
| North America | 48.9 (42.0-55.7) | Ref | Ref | Ref |
| Europe | 44.0 (37.7-50.4) | 0.8 (0.6-1.2) | 0.6 (0.4-0.8)*** | 0.1 |
| Australasia | 28.2 (20.3-37.6) | 0.4 (0.2-0.7)** | 0.4 (0.3-0.7)*** | 0.07 |
| Asia | 34.2 (28.6-40.3) | 0.5 (0.4-0.8)** | 0.5 (0.4-0.7)*** | 0.1 |
| South America | 47.5 (31.2-64.4) | 0.9 (0.5-1.8) | 0.8 (0.5-1.4) | 0.1 |
| Africa | 23.6 (12.3-32.8) | 0.3 (0.1-0.9)* | 0.3 (0.2-0.8)* | 0.1 |
| Multiple continents | 41.4 (31.0-52.6) | 0.7 (0.4-1.4) | 0.6 (0.4-1.1) | 0.1 |

*<0.05 **<0.01 ***<0.001
Ref: Reference category. FMI: Fraction of missing information.

Table S8: Definition of variables

| Variable name | Definition |
|--|---|
| Study setting | |
| Community | Studies that used population surveys, insurance claims databases, or research databases |
| Primary care | Studies that were carried out in primary care settings |
| Hospital | Studies that were carried out in hospital settings |
| Data source | |
| Self-report | Studies that collected data using self-report or interviews |
| Medical records and administrative databases | Studies that collected data using electronic medical records, medical chart reviews, insurance claims databases, pharmacy databases, or research databases |
| Study population | |
| All adults | Studies with a sample of population aged 18 and older (n=45), aged 20 and older (n=8), aged 21 and older (n=3), aged 25 and older (n=2), or others (n=27) (e.g. aged 16 and older, or aged 17 and older) |
| Middle-aged and older | Studies with a sample of population aged 50 and older (n=25), aged 40 and older (n=5), aged 40 and older (n=10), or others (n=6) (e.g. aged 57 and older, or aged 45 and older) |
| Only older people | Studies with a sample of population aged 65 and older (n=22), aged 60 and older (n=25), aged 70 and older (n=5) or others (n=11) (e.g. aged 68 and older, aged 77 and older, aged 78 and older, or aged 80 and older) |

Figure S1: Graphical display of study effect sizes and heterogeneity



No obvious subgroup effects were identified

Figure S2: Process of examining and identifying outlying studies in meta-analysis




Figure S3: Summary of risk of bias assessment





Figure S4: Meta-regression trees for predicting the pooled estimated prevalence of multimorbidity (based on ‘mean age’ and ‘number of conditions’ predictors. unit: log(odds))



References

1 Stanley J, Semper K, Millar E, Sarfati D. Epidemiology of multimorbidity in New Zealand: A cross-sectional study using national-level hospital and pharmaceutical data. *BMJ Open* 2018; **8**:1-12.

2 Lenzi J, Avaldi VM, Rucci P, Pieri G, Fantini MP. Burden of multimorbidity in relation to age, gender and immigrant status: A cross-sectional study based on administrative data. *BMJ Open* 2016; **6**:1-12.

3 Hu RH, Hsiao FY, Chen LJ, Huang PT, Hsu WW. Increasing age- and gender-specific burden and complexity of multimorbidity in taiwan, 2003-2013: A cross-sectional study based on nationwide claims data. *BMJ Open* 2019; **9**:1-10.

4 Gawron LM, Sanders JN, Sward K, Poursaid AE, Simmons R, Turok DK. Multimorbidity and highly effective contraception in reproductive-age women in the us intermountain west: A retrospective cohort study. *J Gen Intern Med* 2020; **35**:637-42.

5 Low LL, Kwan YH, Ko MSM, et al. Epidemiologic characteristics of multimorbidity and sociodemographic factors associated with multimorbidity in a rapidly aging asian country. *JAMA Network Open* 2019; **2**:1-14.

6 Wang HHX, Wang JJ, Wong SYS, et al. Epidemiology of multimorbidity in china and implications for the healthcare system: Cross-sectional survey among 162,464 community household residents in southern china. *BMC Medicine* 2014; **12**:1-12.

7 Gaulin M, Simard M, Candas B, Lesage A, Sirois C. Combined impacts of multimorbidity and mental disorders on frequent emergency department visits: A retrospective cohort study in quebec, canada. *Canadian Medical Association Journal* 2019; **191**:E724-E32.

8 Violan C, Foguet-Boreu Q, Flores-Mateo G, et al. Prevalence, determinants and patterns of multimorbidity in primary care: A systematic review of observational studies. *PLOS One* 2014; **9**:1-9.

9 Nicholson K, Terry AL, Fortin M, Williamson T, Bauer M, Thind A. Prevalence, characteristics, and patterns of patients with multimorbidity in primary care: A retrospective cohort analysis in canada. *British Journal of General Practice* 2019; **69**:1-10.

10 Bao X-Y, Xie Y-X, Zhang X-X, et al. The association between multimorbidity and health-related quality of life: A cross-sectional survey among community middle-aged and elderly residents in southern china. *Health and quality of life outcomes* 2019; **17**:107-09.

11 Fortin M, Bravo G, Hudon C, Vanasse A, Lapointe L. Prevalence of multimorbidity among adults seen in family practice. *Ann Fam Med* 2005; **3**:223-8.

12 Prazeres F, Santiago L. Prevalence of multimorbidity in the adult population attending primary care in portugal: A cross-sectional study. *BMJ Open* 2015; **5**:1-10.

13 Lawson KD, Mercer SW, Wyke S, et al. Double trouble: The impact of multimorbidity and deprivation on preference-weighted health related quality of life a cross sectional analysis of the scottish health survey. *Int J Equity Health* 2013; **12**:1-9.

14 Sullivan PW, Ghushchyan VH, Bayliss EA. The impact of co-morbidity burden on preference-based health-related quality of life in the United States. *Pharmacoeconomics* 2012; **30**:431-42.

15 Peng X, Bao X, Xie Y, et al. The mediating effect of pain on the association between multimorbidity and disability and impaired physical performance among community-dwelling older adults in southern china. *Aging Clin Exp Res* 2020; **32**:1327-34.

- 16 Excoffier S, Herzig L, N’Goran AA, Déruaz-Luyet A, Haller DM. Prevalence of multimorbidity in general practice: A cross-sectional study within the Swiss sentinel surveillance system (sentinella). *BMJ Open* 2018; **8**:1-7.
- 17 Chung RY, Mercer S, Lai FTT, Yip BHK, Wong MCS, Wong SYS. Socioeconomic determinants of multimorbidity: A population-based household survey of Hong Kong Chinese. *PLOS ONE* 2015; **10**:1-15.
- 18 Ki M, Lee YH, Kim Y-S, Shin J-Y, Lim J, Nazroo J. Socioeconomic inequalities in health in the context of multimorbidity: A Korean panel study. *PLOS ONE* 2017; **12**:1-14.
- 19 Bobo WV, Yawn BP, St Sauver JL, Grossardt BR, Boyd CM, Rocca WA. Prevalence of combined somatic and mental health multimorbidity: Patterns by age, sex, and race/ethnicity. *J Gerontol - A Biol Sci Med Sci* 2016; **71**:1483-91.
- 20 Randall DA, Lujic S, Havard A, Eades SJ, Jorm L. Multimorbidity among aboriginal people in New South Wales contributes significantly to their higher mortality. *Med J Aust* 2018; **209**:19-23.
- 21 Russell J, Grant CC, Morton SMB. Multimorbidity in early childhood and socioeconomic disadvantage: Findings from a large New Zealand child cohort. *Acad Pediatr* 2020; **20**:619-27.
- 22 Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: A cross-sectional study. *The Lancet* 2012; **380**:37-43.
- 23 St Sauver JL, Boyd CM, Grossardt BR, et al. Risk of developing multimorbidity across all ages in an historical cohort study: Differences by sex and ethnicity. *BMJ Open* 2015; **5**:1-13.
- 24 Vetrano DL, Foebel AD, Marengoni A, et al. Chronic diseases and geriatric syndromes: The different weight of comorbidity. *Eur J Intern Med* 2016; **27**:62-7.
- 25 Aarts S, den Akker Mv, Bosma H, et al. The effect of multimorbidity on health related functioning: Temporary or persistent? Results from a longitudinal cohort study. *J Psychosom Res* 2012; **73**:211-17.
- 26 Aarts S, van den Akker M, Hajema KJ, et al. Multimorbidity and its relation to subjective memory complaints in a large general population of older adults. *Int Psychogeriatr* 2011; **23**:616-24.
- 27 Aarts S, van den Akker M, Tan FES, Verhey FRJ, Metsemakers JFM, van Boxtel MPJ. Influence of multimorbidity on cognition in a normal aging population: A 12-year follow-up in the maastricht aging study. *Int J Geriatr Psychiatry* 2011; **26**:1046-53.
- 28 Abizanda P, Romero L, Sanchez-Jurado PM, Martinez-Reig M, Alfonso-Silguero SA, Rodriguez-Manas L. Age, frailty, disability, institutionalization, multimorbidity or comorbidity. Which are the main targets in older adults? *J Nutr; Health Aging* 2014; **18**:622-27.
- 29 Agborsangaya CB, Lau D, Lahtinen M, Cooke T, Johnson JA. Health-related quality of life and healthcare utilization in multimorbidity: Results of a cross-sectional survey. *Qual Life Res* 2012; **22**:791-99.
- 30 Agborsangaya CB, Ngwakongnwi E, Lahtinen M, Cooke T, Johnson JA. Multimorbidity prevalence in the general population: The role of obesity in chronic disease clustering. *BMC Public Health* 2013; **13**:1161-61.
- 31 Agborsangaya CB, Lahtinen M, Cooke T, Johnson JA. Comparing the eq-5d 3l and 5l: Measurement properties and association with chronic conditions and multimorbidity in the general population. *Health Qual Life Outcomes* 2014; **12**:1-7.

- 32 Ahrenfeldt LJ, Möller S, Thinggaard M, Christensen K, Lindahl-Jacobsen R. Sex differences in comorbidity and frailty in Europe. *Int J Public Health* 2019; **64**:1025-36.
- 33 Alimohammadian M, Majidi A, Yaseri M, et al. Multimorbidity as an important issue among women: Results of a gender difference investigation in a large population-based cross-sectional study in west asia. *BMJ Open* 2017; **7**:1-8.
- 34 Angst J, Sellaro R, Ries Merikangas K. Multimorbidity of psychiatric disorders as an indicator of clinical severity. *Eur Arch Psychiatry Clin Neurosci* 2002; **252**:147-54.
- 35 Appa AA, Creasman J, Brown JS, et al. The impact of multimorbidity on sexual function in middle-aged and older women: Beyond the single disease perspective. *J Sex Med* 2014; **11**:2744-55.
- 36 Adams ML, Grandpre J, Katz DL, Shenson D. Linear association between number of modifiable risk factors and multiple chronic conditions: Results from the behavioral risk factor surveillance system. *Prev Med* 2017; **105**:169-75.
- 37 Ahmadi B, Alimohammadian M, Yaseri M, et al. Multimorbidity: Epidemiology and risk factors in the golesan cohort study, iran a cross-sectional analysis. *Medicine* 2016; **95**:1-7.
- 38 Amaral TLM, Amaral CdA, de Lima NS, Herculano PV, Do Prado PR, Monteiro GTR. Multimorbidity, depression and quality of life among elderly people assisted in the family health strategy in senador guiomard, acre, Brazil. *Ciencia e Saude Coletiva* 2018; **23**:3077-84.
- 39 An KO Kim J. Association of sarcopenia and obesity with multimorbidity in Korean adults: A nationwide cross-sectional study. *J Am Med Dir Assoc* 2016; **17**:e1-7.
- 40 Araujo MEA, Silva MT, Galvao TF, Nunes BP, Pereira MG. Prevalence and patterns of multimorbidity in amazon region of Brazil and associated determinants: A cross-sectional study. *BMJ Open* 2018; **8**:1-11.
- 41 Arnold-Reed D, Troeung L, Brett T, et al. Increasing multimorbidity in an Australian street health service: A 10-year retrospective cohort study. *Aust J Gen Pract* 2018; **47**:181-89.
- 42 Arokiasamy P, Uttamacharya U, Jain K, et al. The impact of multimorbidity on adult physical and mental health in low- and middle-income countries: What does the study on global ageing and adult health (sage) reveal? *BMC Med* 2015; **13**:1-16.
- 43 Sinnige J, Korevaar JC, Westert GP, Spreeuwenberg P, Schellevis FG, Braspenning JC. Multimorbidity patterns in a primary care population aged 55 years and over. *Fam Pract* 2015; **32**:505-13.
- 44 Zemedikun DT, Gray LJ, Khunti K, Davies MJ, Dhalwani NN. Patterns of multimorbidity in middle-aged and older adults: An analysis of the uk biobank data. *Mayo Clin Proc* 2018; **93**:857-66.
- 45 Wensing M, Vingerhoets E, Grol R. Functional status, health problems, age and comorbidity in primary care patients. *Qual Life Res* 2001; **10**:141-48.
- 46 Mounce LTAP, Campbell JLMD, Henley WEP, Arreal MCTPMPHLCP, Porter IP, Valderas JMBPMPH. Predicting incident multimorbidity. *Ann Fam Med* 2018; **16**:322-29.
- 47 Taylor AW, Price K, Gill TK, et al. Multimorbidity - not just an older person's issue. Results from an Australian biomedical study. *BMC Public Health* 2010; **10**:1-10.
- 48 Vancampfort D, Stubbs B, Firth J, Koyanagi A. Handgrip strength, chronic physical conditions and physical multimorbidity in middle-aged and older adults in six low- and middle income countries. *Eur J Intern Med* 2019; **61**:96-102.

- 1
- 2
- 3
- 4 49 Vancampfort D, Smith L, Stubbs B, et al. Associations between active travel and
- 5 physical multi-morbidity in six low- and middle-income countries among community-
- 6 dwelling older adults: A cross-sectional study. *PloS one* 2018; **13**:1-15.
- 7 50 Aubert CE, Streit S, Da Costa BR, et al. Polypharmacy and specific comorbidities in
- 8 university primary care settings. *Eur J Intern Med* 2016; **35**:35-42.
- 9 51 Autenrieth CS, Kirchberger I, Heier M, et al. Physical activity is inversely associated
- 10 with multimorbidity in elderly men: Results from the kora-age augsburg study. *Prev*
- 11 *Med* 2013; **57**:17-19.
- 12 52 Bähler C, Huber CA, Brüngger B, Reich O. Multimorbidity, health care utilization
- 13 and costs in an elderly community-dwelling population: A claims data based
- 14 observational study. *BMC Health Services Res* 2015; **15**:1-12.
- 15 53 Vancampfort DK, A.: Ward, P. B.: Veronese, N.: Carvalho, A. F.: Solmi, M.: Mugisha,
- 16 J.: Rosenbaum, S.: De Hert, M.: Stubbs, B. Perceived stress and its relationship with
- 17 chronic medical conditions and multimorbidity among 229,293 community-dwelling
- 18 adults in 44 low- and middle-income countries. *Am J Epidemiol* 2017; **186**:979-89.
- 19 54 Banjare P Pradhan J. Socio-economic inequalities in the prevalence of multi-
- 20 morbidity among the rural elderly in Bargarh district of Odisha (India). *PLoS ONE*
- 21 2014; **9**:1-10.
- 22 55 Barra M, Augestad LA, Whitehurst DGT, Rand-Hendriksen K. Examining the
- 23 relationship between health-related quality of life and increasing numbers of
- 24 diagnoses. *Qual life Res* 2015; **24**:2823-32.
- 25 56 Bernard S, Inderjeeth C, Raymond W. Higher charlson comorbidity index scores do
- 26 not influence functional independence measure score gains in older rehabilitation
- 27 patients. *Aust J Ageing* 2016; **35**:236-41.
- 28 57 Biswas T, Townsend N, Islam MS, et al. Association between socioeconomic status
- 29 and prevalence of non-communicable diseases risk factors and comorbidities in
- 30 bangladesh: Findings from a nationwide cross-sectional survey. *BMJ Open* 2019; **9**:1-
- 31 9.
- 32 58 Blakemore A, Hann M, Howells K, et al. Patient activation in older people with long-
- 33 term conditions and multimorbidity: Correlates and change in a cohort study in the
- 34 united kingdom. *BMC Health Services Res* 2016; **16**:582-82.
- 35 59 Blyth FM, Rochat S, Cumming RG, et al. Pain, frailty and comorbidity on older men:
- 36 The champ study. *Pain* 2008; **140**:224-30.
- 37 60 Bowling CB, Deng L, Sakhuja S, Morey MC, Jaeger BC, Muntner P. Prevalence of
- 38 activity limitations and association with multimorbidity among us adults 50 to
- 39 64 years old. *J Gen Intern Med* 2019; **34**:2390-96.
- 40 61 Britt HC, Harrison CM, Miller GC, Knox SA. Prevalence and patterns of
- 41 multimorbidity in australia. *Med J Aust* 2008; **189**:72-77.
- 42 62 Broeiro-Gonçalves P, Nogueira P, Aguiar P. Multimorbidity and disease severity by
- 43 age groups, in inpatients: Cross-sectional study. *Portuguese J Public Health* 2019;
- 44 **37**:1-9.
- 45 63 Bruce SG, Riediger ND, Zacharias JM, Young TK. Obesity and obesity-related
- 46 comorbidities in a canadian first nation population. *Chronic Dis Can* 2010; **31**:27-32.
- 47 64 Burgers JS, Voerman GE, Grol R, Faber MJ, Schneider EC. Quality and coordination
- 48 of care for patients with multiple conditions: Results from an international survey of
- 49 patient experience. *Eval Health Prof* 2010; **33**:343-64.
- 50 65 Burke BT. *Connections that count: A network analysis of multimorbidity and gait*
- 51 *speed in older adults*, in *Department of Epidemiology (dissertation)*. 2017, Johns
- 52 Hopkins University: Baltimore, Maryland.1-224.
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 66 Buurman BM, Frenkel WJ, Abu-Hanna A, Parlevliet JL, de Rooij SE. Acute and chronic diseases as part of multimorbidity in acutely hospitalized older patients. *Eur J Intern Med* 2016; **27**:68-75.
- 67 Calderón-Larrañaga A, Vetrano DL, Onder G, et al. Assessing and measuring chronic multimorbidity in the older population: A proposal for its operationalization. *J Gerontol A Biol Sci Med Sci* 2017; **72**:1417-23.
- 68 Camargo-Casas S, Suarez-Monsalve S, Zepeda MUP, García-Peña C, Cano-Gutierrez CA. Multimorbidity, depressive symptoms, and self-reported health in older adults: A secondary analysis of the sabe bogota study. *Rev Invest Clin* 2018; **70**:192-97.
- 69 Canevelli M, Raganato R, Remiddi F, et al. Counting deficits or diseases? The agreement between frailty and multimorbidity in subjects with cognitive disturbances. *Aging Clin Exp Res* 2019; **32**:179-82.
- 70 Chamberlain AM, Rutten LJF, Wilson PM, et al. Neighborhood socioeconomic disadvantage is associated with multimorbidity in a geographically-defined community. *BMC Public Health* 2020; **20**:1-10.
- 71 Chen H, Chen Y, Cui B. The association of multimorbidity with healthcare expenditure among the elderly patients in Beijing, China. *Arch Gerontol Geriatr* 2018; **79**:32-38.
- 72 Chen H, Cheng M, Zhuang Y, Broad JBJG, International G. Multimorbidity among middle-aged and older persons in urban china: Prevalence, characteristics and health service utilization. *Geriatr Gerontol Int* 2018; **18**:1447-52.
- 73 Cheung CL, Nguyen US, Au E, Tan KC, Kung AW. Association of handgrip strength with chronic diseases and multimorbidity: A cross-sectional study. *Age (Dordr)* 2013; **35**:929-41.
- 74 Chu T, Lau P, Cheng R, Chan M, Liang J. A study on the prevalence of multimorbidities of diseases and utilisation of public healthcare services in the new territories west area of Hong Kong. *Hong Kong Practitioner* 2018; **40**:43-50.
- 75 Chudasama YV, Khunti KK, Zaccardi F, et al. Physical activity, multimorbidity, and life expectancy: A UK biobank longitudinal study. *BMC Med* 2019; **17**:1-13.
- 76 Cimarras-Otal C, Calderón-Larrañaga A, Poblador-Plou B, et al. Association between physical activity, multimorbidity, self-rated health and functional limitation in the Spanish population. *BMC Public Health* 2014; **14**:1-10.
- 77 Chin WY, Choi EP, Wan EY, Lam CL. Health-related quality of life mediates associations between multi-morbidity and depressive symptoms in Chinese primary care patients. *Fam Pract* 2016; **33**:61-8.
- 78 Agrawal S, Agrawal PK. Association between body mass index and prevalence of multimorbidity in low-and middle-income countries: A cross-sectional study. *Int J Med Public Health* 2016; **6**:73-83.
- 79 Gu J, Chao J, Chen W, et al. Multimorbidity and health-related quality of life among the community-dwelling elderly: A longitudinal study. *Arch Gerontol Geriatr* 2018; **74**:133-40.
- 80 Gunn JM, Ayton DR, Densley K, et al. The association between chronic illness, multimorbidity and depressive symptoms in an australian primary care cohort. *Soc Psychiatry Psychiatr Epidemiol* 2010; **47**:175-84.
- 81 Han MA, Tinetti ME, Agostini JV, Han L, Lee H, Walke LM. The impact of medication use on symptom burden in older patients with multiple medical morbidities. *J Clin Gerontol and Geriatr* 2013; **4**:84-88.
- 82 Hanlon P, Nicholl BI, Jani BD, Lee D, McQueenie R, Mair FS. Frailty and pre-frailty in middle-aged and older adults and its association with multimorbidity and mortality:

- A prospective analysis of 493 737 UK biobank participants. *Lancet Public Health* 2018; **3**:e323-e32.
- 83 Jantsch AG, Alves RFS, Faerstein E. Educational inequality in Rio de Janeiro and its impact on multimorbidity: Evidence from the pró-saúde study. A cross-sectional analysis. *Sao Paulo Med J* 2018; **136**:51-58.
- 84 John R, Kerby DS, Hagan Hennessy C. Patterns and impact of comorbidity and multimorbidity among community-resident American Indian elders. *Gerontol* 2003; **43**:649-60.
- 85 Johnson-Lawrence V, Zajacova A, Sneed R. Education, race/ethnicity, and multimorbidity among adults aged 30-64 in the national health interview survey. *SSM Popul Health* 2017; **3**:366-72.
- 86 Johnston MC, Black C, Mercer SW, Prescott GJ, Crilly MA. Impact of educational attainment on the association between social class at birth and multimorbidity in middle age in the Aberdeen children of the 1950s cohort study. *BMJ Open* 2019; **9**:1-9.
- 87 Jones SM, Amtmann D, Gell NM. A psychometric examination of multimorbidity and mental health in older adults. *Aging Ment Health* 2016; **20**:309-17.
- 88 Jovic D, Marinkovic J, Vukovic D. Association between body mass index and prevalence of multimorbidity: A cross-sectional study. *Public Health* 2016; **139**:103-11.
- 89 Juul-Larsen HG, Andersen O, Bandholm T, et al. Differences in function and recovery profiles between patterns of multimorbidity among older medical patients the first year after an acute admission—an exploratory latent class analysis. *Arch Gerontol Geriatr* 2020; **86**:1-11.
- 90 Hudon C, Soubhi H, Fortin M. Relationship between multimorbidity and physical activity: Secondary analysis from the Quebec health survey. *BMC Public Health* 2008; **8**:1-8.
- 91 Hussain MA, Huxley RR, Al Mamun A. Multimorbidity prevalence and pattern in Indonesian adults: An exploratory study using national survey data. *BMJ Open* 2015; **5**:1-10.
- 92 Ie K, Felton M, Springer S, Wilson SA, Albert SM. Multimorbidity and polypharmacy in family medicine residency practices. *J Pharm Technol* 2017; **33**:219-24.
- 93 Ishizaki T, Kobayashi E, Fukaya T, Takahashi Y, Shinkai S, Liang J. Association of physical performance and self-rated health with multimorbidity among older adults: Results from a nationwide survey in japan. *Arch Gerontol Geriatr* 2019; **84**:1-9.
- 94 Danon-Hersch N, Rodondi N, Spagnoli J, Santos-Eggimann B. Prefrailty and chronic morbidity in the youngest old: An insight from the Lausanne cohort Lc65+. *J Am Geriatr Soc* 2012; **60**:1687-94.
- 95 de Heer HD, Balcázar HG, Morera OF, et al. Barriers to care and comorbidities along the u.S.-mexico border. *Public Health Rep* 2013; **128**:480-8.
- 96 Demirchyan A, Khachadourian V, Armenian HK, Petrosyan V. Short and long term determinants of incident multimorbidity in a cohort of 1988 earthquake survivors in Armenia. *Int. J. Equity Health* 2013; **12**:68-68.
- 97 Fabbri E, An Y, Zoli M, et al. Aging and the burden of multimorbidity: Associations with inflammatory and anabolic hormonal biomarkers. *J Gerontol A Biol Sci Med Sci* 2015; **70**:63-70.
- 98 Fillenbaum GGP, C. F.: Cohen, H. J.: Cornoni-Huntley, J. C.: Guralnik, J. M. Comorbidity of five chronic health conditions in elderly community residents: Determinants and impact on mortality. *J Gerontol: Med Sci* 2000; **55**:M84-89.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
- 99 Kaneko M, Boven Kv, Takayanagi H, Kusaba T, Yamada T, Matsushima MJFp. Multicentre descriptive cross-sectional study of japanese home visit patients: Reasons for encounter, health problems and multimorbidity. *Fam Pract* 2019; **37**:227-33.
- 100 Kang J Song Y-M. The association between submarine service and multimorbidity: A cross-sectional study of korean naval personnel. *BMJ Open* 2017; **7**:1-6.
- 101 Gandhi K, Lim E, Davis J, Chen JJ. Racial-ethnic disparities in self-reported health status among us adults adjusted for sociodemographics and multimorbidities, national health and nutrition examination survey 2011-2014. *Ethn Health* 2020; **25**:65-78.
- 102 Costa CDS, Flores TR, Wendt A, et al. Inequalities in multimorbidity among elderly: A population-based study in a city in southern brazil. *Cad Saude Publica* 2018; **34**:1-12.
- 103 Rizzuto DM, R. J. F.: Angleman, S.: Qiu, C.: Marengoni, A. Effect of chronic diseases and multimorbidity on survival and functioning in elderly adults. *J Am Geriatr Soc* 2017; **65**:1056-60.
- 104 Dhalwani NN, Zaccardi F, O'Donovan G, et al. Association between lifestyle factors and the incidence of multimorbidity in an older english population. *J Gerontol A Biol Sci Med Sci* 2017; **72**:528-34.
- 105 Elixhauser A, Steiner C, Harris DR, Coffey RM. Comorbidity measures for use with administrative data. *Med Care* 1998; **36**:8-27.
- 106 Fabbri E, An Y, Schrack JA, et al. Energy metabolism and the burden of multimorbidity in older adults: Results from the baltimore longitudinal study of aging. *J Gerontol A Biol Sci Med Sci* 2015; **70**:1297-303.
- 107 Fortin M, Haggerty J, Almirall J, Bouhali T, Sasseville M, Lemieux M. Lifestyle factors and multimorbidity: A cross sectional study. *BMC Public Health* 2014; **14**:1-6.
- 108 Fuchs ZB, T.: Novikov, I.: Walter-Ginzburg, A.: Lyanders, M.: Gindin, J.: Habet, B.: Modan, B. Morbidity, comorbidity, and their association with disability among community-dwelling oldest-old in israel. *J Gerontol: Med Sci* 1998; **53**:M447-55.
- 109 Galenkamp H, Braam AW, Huisman M, Deeg DJ. Somatic multimorbidity and self-rated health in the older population. *J Gerontol B Psychol Sci Soc Sci* 2011; **66**:380-6.
- 110 Galenkamp H, Gagliardi C, Principi A, et al. Predictors of social leisure activities in older europeans with and without multimorbidity. *Eur J Ageing* 2016; **13**:129-43.
- 111 Gamma A Angst J. Concurrent psychiatric comorbidity and multimorbidity in a community study: Gender differences and quality of life. *Eur Arch Psychiatry Clin Neurosci* 2001; **251**:43-46.
- 112 Ge L, Yap CW, Heng BH. Sex differences in associations between multimorbidity and physical function domains among community-dwelling adults in Singapore. *PLoS One* 2018; **13**:1-14.
- 113 Ge L, Ong R, Yap CW, Heng BH. Effects of chronic diseases on health-related quality of life and self-rated health among three adult age groups. *Nurs Health Sci* 2019; **21**:214-22.
- 114 Gould CE, O'Hara R, Goldstein MK, Beaudreau SA. Multimorbidity is associated with anxiety in older adults in the health and retirement study. *Int J Geriatr Psychiatry* 2016; **31**:1105-15.
- 115 Habib RR, Hojeij S, Elzein K, Chaaban J, Seyfert K. Associations between life conditions and multi-morbidity in marginalized populations: The case of palestinian refugees. *Eur J Public Health* 2014; **24**:727-33.
- 116 Harrison C, Henderson J, Miller G, Britt H. The prevalence of diagnosed chronic conditions and multimorbidity in australia: A method for estimating population prevalence from general practice patient encounter data. *PLOS ONE* 2017; **12**:1-13.

- 117 Hayek S, Ifrah A, Enav T, Shohat T. Prevalence, correlates, and time trends of multiple chronic conditions among israeli adults: Estimates from the Israeli national health interview survey, 2014-2015. *Prev Chronic Dis* 2017; **14**:1-14.
- 118 Henninger DE, Whitson HE, Cohen HJ, Ariely D. Higher medical morbidity burden is associated with external locus of control. *J Am Geriatr Soc* 2012; **60**:751-55.
- 119 Hernández B, Reilly RB, Kenny RA. Investigation of multimorbidity and prevalent disease combinations in older irish adults using network analysis and association rules. *Sci Rep* 2019; **9**:1-12.
- 120 Ho C, Feng L, Fam J, Mahendran R, Kua EH, Ng TP. Coexisting medical comorbidity and depression: Multiplicative effects on health outcomes in older adults. *Int Psychogeriatr* 2014; **26**:1221-9.
- 121 Khan N, Rahman M, Mitra D, Afsana K. Prevalence of multimorbidity among bangladeshi adult population: A nationwide cross-sectional study. *BMJ Open* 2019; **9**:1-8.
- 122 Kiliari N, Theodosopoulou E, Papanastasiou E. Multimorbidity and unmet citizens' needs and expectations urge for reforms in the health system of cyprus: A questionnaire survey. *JRSM Open* 2014; **5**:1-11.
- 123 King DE, Xiang J, Pilkerton CS. Multimorbidity trends in united states adults, 1988-2014. *J Am Board Fam Med* 2018; **31**:503-13.
- 124 Kingston AR, L.: Booth, H.: Knapp, M.: Jagger, C.: Modem project. Projections of multi-morbidity in the older population in england to 2035: Estimates from the population ageing and care simulation (pacsim) model. *Age Ageing* 2018; **47**:374-80.
- 125 Koyanagi A, Lara E, Stubbs B, et al. Chronic physical conditions, multimorbidity, and mild cognitive impairment in low- and middle-income countries. *J Am Geriatr Soc* 2018; **66**:721-27.
- 126 Kriegsman DMD, D. J.: Stalman, W. A. Comorbidity of somatic chronic diseases and decline in physical functioning;; the longitudinal aging study amsterdam. *J Clin Epidemiol* 2004; **57**:55-65.
- 127 Kristensen K, König HH, Hajek A. The longitudinal association of multimorbidity on loneliness and network size: Findings from a population-based study. *Int J Geriatr Psychiatry* 2019; **34**:1490-97.
- 128 Kristensen K, König HH, Hajek A. The association of multimorbidity, loneliness, social exclusion and network size: Findings from the population-based german ageing survey. *BMC Public Health* 2019; **19**:1-10.
- 129 Kunna R, San Sebastian M, Stewart Williams J. Measurement and decomposition of socioeconomic inequality in single and multimorbidity in older adults in china and ghana: Results from the who study on global ageing and adult health (sage). *Int J Equity Health* 2017; **16**:2-17.
- 130 Kusumastuti S, Gerds TA, Lund R, Mortensen EL, Westendorp RGJ. Discrimination ability of comorbidity, frailty, and subjective health to predict mortality in community-dwelling older people: Population based prospective cohort study. *Eur J Intern Med* 2017; **42**:29-38.
- 131 Lai FTT, Guthrie B, Wong SYS, et al. Sex-specific intergenerational trends in morbidity burden and multimorbidity status in hong kong community: An age-period-cohort analysis of repeated population surveys. *BMJ Open* 2019; **9**:1-10.
- 132 Lai FTT, Ma TW, Hou WK. Multimorbidity is associated with more subsequent depressive symptoms in three months: A prospective study of community-dwelling adults in hong kong. *Int Psychogeriatr* 2018; **31**:1367-71.
- 133 Laires PA Perelman J. The current and projected burden of multimorbidity: A cross-sectional study in a southern europe population. *Eur J Ageing* 2019; **16**:181-92.

- 1
2
3 134 Lang K, Alexander IM, Simon J, et al. The impact of multimorbidity on quality of life
4 among midlife women: Findings from a u.S. Nationally representative survey. *J*
5 *Women's Health* 2015; **24**:374-83.
6
7 135 Le Cossec C, Perrine AL, Beltzer N, Fuhrman C, Carcaillon-Bentata L. Pre-frailty,
8 frailty, and multimorbidity: Prevalences and associated characteristics from two
9 french national surveys. *J Nutr Health Aging* 2016; **20**:860-69.
10 136 Lee TA, Shields AE, Vogeli C, et al. Mortality rate in veterans with multiple chronic
11 conditions. *J Gen Intern Med* 2007; **22** 403-7.
12 137 Lee WJ, Peng LN, Lin CH, Lin HP, Loh CH, Chen LK. The synergic effects of frailty
13 on disability associated with urbanization, multimorbidity, and mental health:
14 Implications for public health and medical care. *Sci Rep* 2018; **8**:1-7.
15 138 Li J, Green M, Kearns B, et al. Patterns of multimorbidity and their association with
16 health outcomes within yorkshire, england: Baseline results from the yorkshire health
17 study. *BMC Public Health* 2016; **16**:1-9.
18 139 Li Y, White K, O'Shields KR, McLain AC, Merchant AT. Light-intensity physical
19 activity and cardiometabolic risk among older adults with multiple chronic conditions.
20 *Am J Health Promot* 2019; **33**:507-15.
21 140 Lujic S, Simpson JM, Zwar N, Hosseinzadeh H, Jorm L. Multimorbidity in australia:
22 Comparing estimates derived using administrative data sources and survey data. *PLoS*
23 *One* 2017; **12**:1-18.
24 141 Lupiáñez-Villanueva F, Anastasiadou D, Codagnone C, Nuño-Solinís R, Garcia-
25 Zapirain Soto MB. Electronic health use in the european union and the effect of
26 multimorbidity: Cross-sectional survey. *J Med Internet Res* 2018; **20**:1-14.
27 142 Zhou C-h, Tang S-f, Wang X-h, et al. Satisfaction about patient-centeredness and
28 healthcare system among patients with chronic multimorbidity. *Current Med Sci*
29 2018; **38**:184-90.
30 143 Zhang R, Lu Y, Shi L, Zhang S, Chang F. Prevalence and patterns of multimorbidity
31 among the elderly in china: A cross-sectional study using national survey data. *BMJ*
32 *Open* 2019; **9**:1-9.
33 144 Wong CH, Weiss D, Sourial N, et al. Frailty and its association with disability and
34 comorbidity in a community-dwelling sample of seniors in montreal: A cross-
35 sectional study. *Aging Clin Exp Res* 2010; **22**:54-62.
36 145 Weimann A, Dai D, Oni T. A cross-sectional and spatial analysis of the prevalence of
37 multimorbidity and its association with socioeconomic disadvantage in south africa: A
38 comparison between 2008 and 2012. *Soc Sci Med* 2016; **163**:144-56.
39 146 Wang L, Palmer AJ, Cocker F, Sanderson K. Multimorbidity and health-related
40 quality of life (hrqol) in a nationally representative population sample: Implications of
41 count versus cluster method for defining multimorbidity on hrqol. *Health Qual life*
42 *Outcomes* 2017; **15**:1-12.
43 147 Wang C, Pu R, Li Z, et al. Subjective health and quality of life among elderly people
44 living with chronic multimorbidity and difficulty in activities of daily living in rural
45 South Africa. *Clin Interv Aging* 2019; **14**:1285-96.
46 148 Wade A, Hobbs M, Green MA. Investigating the relationship between multimorbidity
47 and dental attendance: A cross-sectional study of uk adults. *British Dental Journal*
48 2019; **226**:138-43.
49 149 Maciejewski ML Hammill BG. Measuring the burden of multimorbidity among
50 medicare beneficiaries via condition counts and cumulative duration. *Health Serv Res*
51 2019; **54**:484-91.
52 150 Marengoni A, Angleman S, Meinow B, et al. Coexisting chronic conditions in the
53 older population: Variation by health indicators. *Eur J Intern Med* 2016; **31**:29-34.
54
55
56
57
58
59
60

- 151 Marengoni A, von Strauss E, Rizzuto D, Winblad B, Fratiglioni L. The impact of chronic multimorbidity and disability on functional decline and survival in elderly persons. A community-based, longitudinal study. *J Intern Med* 2009; **265**:288-95.
- 152 Marques A, Santos DA, Peralta M, Sardinha LB, González Valeiro M. Regular physical activity eliminates the harmful association of television watching with multimorbidity. A cross-sectional study from the european social survey. *Prev Med* 2018; **109**:28-33.
- 153 Mavaddat N, Valderas JM, van der Linde R, Khaw KT, Kinmonth AL. Association of self-rated health with multimorbidity, chronic disease and psychosocial factors in a large middle-aged and older cohort from general practice: A cross-sectional study. *BMC Fam Pract* 2014; **15**:1-11.
- 154 McDaid O, Hanly MJ, Richardson K, Kee F, Kenny RA, Savva GM. The effect of multiple chronic conditions on self-rated health, disability and quality of life among the older populations of northern ireland and the republic of ireland: A comparison of two nationally representative cross-sectional surveys. *BMJ Open* 2013; **3**:1-9.
- 155 Melis R, Marengoni A, Angleman S, Fratiglioni L. Incidence and predictors of multimorbidity in the elderly: A population-based longitudinal study. *PLoS One* 2014; **9**:1-8.
- 156 Min LC, Wenger NS, Fung C, et al. Multimorbidity is associated with better quality of care among vulnerable elders. *Med Care* 2007; **45**:480-8.
- 157 Momtaz YA, Hamid TA, Yahaya N, Ibrahim R. Effects of chronic comorbidity on psychological well-being among older persons in northern peninsular Malaysia. *Appl Res Qual Life* 2010; **5**:133-46.
- 158 Mondor L, Cohen D, Khan AI, Wodchis WP. Income inequalities in multimorbidity prevalence in ontario, canada: A decomposition analysis of linked survey and health administrative data. *Int J Equity Health* 2018; **17**:1-13.
- 159 Muggah E, Graves E, Bennett C, Manuel DG. The impact of multiple chronic diseases on ambulatory care use; a population based study in ontario, canada. *BMC Health Serv Res* 2012; **12**:1-6.
- 160 Nagel A, Jungert A, Spinneker A, Neuhäuser-Berthold M. The impact of multimorbidity on resting metabolic rate in community-dwelling women over a ten-year period: A cross-sectional and longitudinal study. *J Nutr Health Aging* 2017; **21**:781-86.
- 161 Niedzwiedz CL, Katikireddi SV, Pell JP, Smith DJ. Sex differences in the association between salivary telomere length and multimorbidity within the us health & retirement study. *Age Ageing* 2019; **48**:703-10.
- 162 Nunes BP, Camargo-Figuera FA, Guttier M, et al. Multimorbidity in adults from a southern brazilian city: Occurrence and patterns. *Int J Public Health* 2016; **61**:1013-20.
- 163 Nunes BP, Chiavegatto Filho ADP, Pati S, et al. Contextual and individual inequalities of multimorbidity in brazilian adults: A cross-sectional national-based study. *BMJ Open* 2017; **7**:1-12.
- 164 Nunes BP, Thumé E, Facchini LA. Multimorbidity in older adults: Magnitude and challenges for the brazilian health system. *BMC Public Health* 2015; **15**:1-11.
- 165 Olaya B, Domènech-Abella J, Moneta MV, et al. All-cause mortality and multimorbidity in older adults: The role of social support and loneliness. *Exp Gerontol* 2017; **99**:120-26.
- 166 Olivares DEV, Chambi FRV, Chañi EMM, Craig WJ, Pacheco SOS, Pacheco FJ. Risk factors for chronic diseases and multimorbidity in a primary care context of

- central argentina: A web-based interactive and cross-sectional study. *Int J Environ Res Public Health* 2017; **14**:1-22.
- 167 Park B, Ock M, Lee HA, et al. Multimorbidity and health-related quality of life in koreans aged 50 or older using knhanes 2013-2014. *Health Qual Life Outcomes* 2018; **16**:1-10.
- 168 Patel KV, Peek MK, Wong R, Markides KS. Comorbidity and disability in elderly mexican and mexican american adults: Findings from mexico and the southwestern united states. *J Aging Health* 2006; **18**:315-29.
- 169 Pati S, Hussain MA, Swain S, et al. Development and validation of a questionnaire to assess multimorbidity in primary care: An indian experience. *BioMed Res Int* 2016; **2016**:1-9.
- 170 Pati SS, Subhashisa: Knottnerus, J.: Metsemakers, Job F.: van den Akker, Marjan. Health related quality of life in multimorbidity: A primary-care based study from odisha, India. *Health Qual Life Outcomes* 2019; **17**:1-11.
- 171 Pati S, Swain S, Metsemakers J, Knottnerus JA, van den Akker M. Pattern and severity of multimorbidity among patients attending primary care settings in Odisha, india. *PLoS One* 2017; **12**:1-19.
- 172 Payne RA, Abel GA, Guthrie B, Mercer SW. The effect of physical multimorbidity, mental health conditions and socioeconomic deprivation on unplanned admissions to hospital: A retrospective cohort study. *Can Med Assoc J* 2013; **185**:E221-8.
- 173 Pérez LM, Hooshmand B, Mangialasche F, et al. Glutathione serum levels and rate of multimorbidity development in older adults. *J Gerontol A Biol Sci Med Sci* 2020; **75**:1089-94.
- 174 Petersen I, Rathod S, Kathree T, Selohilwe O, Bhana A. Risk correlates for physical-mental multimorbidities in south africa: A cross-sectional study. *Epidemiol Psychiatr Sci* 2019; **28**:418-26.
- 175 Pfortmueller CA, Stotz M, Lindner G, Müller T, Rodondi N, Exadaktylos AK. Multimorbidity in adult asylum seekers: A first overview. *PloS one* 2013; **8**:1-7.
- 176 Pressley JC Patrick CH. Frailty bias in comorbidity risk adjustments of community-dwelling elderly populations. *J Clin Epidemiol* 1999; **52**:753-60.
- 177 Prior A, Fenger-Grøn M, Larsen KK, et al. The association between perceived stress and mortality among people with multimorbidity: A prospective population-based cohort study. *Am J Epidemiol* 2016; **184**:199-210.
- 178 Ribeiro AS, Pereira LC, Silva DRP, et al. Physical activity and sitting time are specifically associated with multiple chronic diseases and medicine intake in brazilian older adults. *J Aging Phys Act* 2018; **26**:608-13.
- 179 Ruel G, Lévesque J-F, Stocks N, et al. Understanding the evolution of multimorbidity: Evidences from the north west adelaide health longitudinal study (nwahs). *PLOS ONE* 2014; **9**:1-11.
- 180 Ruel G, Shi Z, Zhen S, et al. Association between nutrition and the evolution of multimorbidity: The importance of fruits and vegetables and whole grain products. *Clin Nutr* 2014; **33**:513-20.
- 181 Ryan A, Murphy C, Boland F, Galvin R, Smith SM. What is the impact of physical activity and physical function on the development of multimorbidity in older adults over time? A population-based cohort study. *J Gerontol A Biol Sci Med Sci* 2018; **73**:1538-44.
- 182 Schmidt AE, Ilinca S, Schulmann K, et al. Fit for caring: Factors associated with informal care provision by older caregivers with and without multimorbidity. *Eur J Ageing* 2016; **13**:103-13.

- 183 Schöttker B, Saum KU, Jansen EH, Holleczeck B, Brenner H. Associations of metabolic, inflammatory and oxidative stress markers with total morbidity and multi-morbidity in a large cohort of older german adults. *Age Ageing* 2016; **45**:127-35.
- 184 Seo JC, B.: Kim, S.: Lee, H.: Oh, D. The relationship between multiple chronic diseases and depressive symptoms among middle-aged and elderly populations: Results of a 2009 korean community health survey of 156,747 participants. *BMC Public Health* 2017; **17**:1-10.
- 185 She R, Yan Z, Jiang H, Vetrano DL, Lau JTF, Qiu C. Multimorbidity and health-related quality of life in old age: Role of functional dependence and depressive symptoms. *J Am Med Dir Assoc* 2019; **20**:1143-49.
- 186 Singh K, Patel SA, Biswas S, et al. Multimorbidity in south asian adults: Prevalence, risk factors and mortality. *J Public Health (Oxf)* 2019; **41**:80-89.
- 187 Spangenberg L, Forkmann T, Brähler E, Glaesmer H. The association of depression and multimorbidity in the elderly: Implications for the assessment of depression. *Psychogeriatr* 2011; **11**:227-34.
- 188 Stickley A, Koyanagi A, Ueda M, Inoue Y, Waldman K, Oh H. Physical multimorbidity and suicidal behavior in the general population in the united states. *J Affect Disord* 2020; **260**:604-09.
- 189 Streit S, da Costa BR, Bauer DC, et al. Multimorbidity and quality of preventive care in swiss university primary care cohorts. *PLOS ONE* 2014; **9**:1-7.
- 190 Stubbs B, Vancampfort D, Veronese N, et al. Multimorbidity and perceived stress: A population-based cross-sectional study among older adults across six low- and middle-income countries. *Maturitas* 2018; **107**:84-91.
- 191 Su P, Ding H, Zhang W, et al. The association of multimorbidity and disability in a community-based sample of elderly aged 80 or older in shanghai, china. *BMC Geriatr* 2016; **16**:1-7.
- 192 Sundstrup E, Jakobsen MD, Mortensen OS, Andersen LL. Joint association of multimorbidity and work ability with risk of long-term sickness absence: A prospective cohort study with register follow-up. *Scand J Work Environ Health* 2017; **43**:146-54.
- 193 Takahashi PY, Ryu E, Hathcock MA, et al. A novel housing-based socioeconomic measure predicts hospitalisation and multiple chronic conditions in a community population. *J Epidemiol Community Health* 2016; **70**:286-91.
- 194 Tinetti MEM, G. J.: Chang, S. S.: Newman, A. B.: Fitzpatrick, A. L.: Fried, T. R.: Peduzzi, P. N. Contribution of multiple chronic conditions to universal health outcomes. *J Am Geriatr Soc* 2011; **59**:1686-91.
- 195 Troelstra SA, Straker L, Harris M, Brown S, van der Beek AJ, Coenen P. Multimorbidity is common among young workers and related to increased work absenteeism and presenteeism: Results from the population-based raine study cohort. *Scand J Work Environ Health* 2020; **46**:218-27.
- 196 van Zon SKR, Reijneveld SA, Galaurchi A, Mendes de Leon CF, Almansa J, Bültmann U. Multimorbidity and the transition out of full-time paid employment: A longitudinal analysis of the health and retirement study. *J Gerontol B Psychol Sci Soc Sci* 2020; **75**:705-15.
- 197 Vancampfort D, Stubbs B, Koyanagi A. Physical chronic conditions, multimorbidity and sedentary behavior amongst middle-aged and older adults in six low- and middle-income countries. *Int J Behav Nutr Phys Act* 2017; **14**:1-13.
- 198 Vassilaki M, Aakre JA, Cha RH, et al. Multimorbidity and risk of mild cognitive impairment. *J Am Geriatr Soc* 2015; **63**:1783-90.

- 199 Vassilaki M, Aakre JA, Mielke MM, et al. Multimorbidity and neuroimaging biomarkers among cognitively normal persons. *Neurology* 2016; **86**:2077-84.
- 200 Villarreal AEG, S.: Lopez, L.: Britton, G. B.: Initiative, P. A. R. Cognitive impairment, depression, and cooccurrence of both among the elderly in panama: Differential associations with multimorbidity and functional limitations. *BioMed Res Int* 2015; **2015**:1-7.
- 201 Violán C, Foguet-Boreu Q, Fernández-Bertolín S, et al. Soft clustering using real-world data for the identification of multimorbidity patterns in an elderly population: Cross-sectional study in a mediterranean population. *BMJ Open* 2019; **9**:1-14.
- 202 von Strauss E, Fratiglioni L, Viitanen M, Forsell Y, Winblad B. Morbidity and comorbidity in relation to functional status: A community-based study of the oldest old (90+ years). *J Am Geriatr Soc* 2000; **48**:1462-9.
- 203 Vos HM, Bor HH, Rangelrooij-Minkels MJ, Schellevis FG, Lagro-Janssen AL. Multimorbidity in older women: The negative impact of specific combinations of chronic conditions on self-rated health. *Eur J Gen Pract* 2013; **19**:117-22.
- 204 Vu HMN, L. H.: Tran, T. H.: Pham, K. T. H.: Phan, H. T.: Nguyen, H. N.: Tran, B. X.: Latkin, C. A.: Ho, C. S. H.: Ho, R. C. M. Effects of chronic comorbidities on the health-related quality of life among older patients after falls in vietnamese hospitals. *Int J Environ Res Public Health* 2019; **16**:1-9.
- 205 Wang L, Cocker F, Kilpatrick M, et al. The associations of multimorbidity with health-related productivity loss in a large and diverse public sector setting: A cross-sectional survey. *J Occup Environ Med* 2018; **60**:528-35.
- 206 Wang X-X, Lin W-Q, Chen X-J, et al. Multimorbidity associated with functional independence among community-dwelling older people: A cross-sectional study in southern china. *Health Qual Life Outcomes* 2017; **15**:73-9.
- 207 Wijers IGM, Ayala A, Rodriguez-Blazquez C, et al. The disease burden morbidity assessment in older adults and its association with mortality and other health outcomes. *Eur J Ageing* 2019; **16**:193-203.
- 208 Williams JSE, L. E. The association between multimorbidity and quality of life, health status and functional disability. *Am J Med Sci* 2016; **352**:45-52.
- 209 Woldeamayrat EM, Kassa A, Gari T, Dangisso MH. Chronic diseases multimorbidity among adult patients at hawassa university comprehensive specialized hospital. *BMC Public Health* 2018; **18**:1-8.
- 210 Yao SS, Cao GY, Han L, et al. Associations between somatic multimorbidity patterns and depression in a longitudinal cohort of middle-aged and older chinese. *J Am Med Dir Assoc* 2020; **21**:1282-87.
- 211 Yorke AM, Curtis AB, Shoemaker M, Vangsnes E. The impact of multimorbidity on grip strength in adults age 50 and older: Data from the health and retirement survey (hrs). *Arch Gerontol Geriatr* 2017; **72**:164-68.
- 212 You L, Yu Z, Zhang X, et al. Association between multimorbidity and depressive symptom among community-dwelling elders in eastern china. *Clin Interv Aging* 2019; **14**:2273-80.
- 213 Zhang L, Ma L, Sun F, Tang Z, Chan P. A multicenter study of multimorbidity in older adult inpatients in china. *J Nutr Health Aging* 2020; **24**:269-76.
- 214 Khanam MA, Streatfield PK, Kabir ZN, Qiu C, Cornelius C, Wahlin Å. Prevalence and patterns of multimorbidity among elderly people in rural bangladesh: A cross-sectional study. *J Health, Popul Nutr* 2011; **29**:406-14.
- 215 Cornell J, Pugh J, Williams, Jr., et al. Multimorbidity clusters: Clustering binary data from multimorbidity clusters: Clustering binary data from a large administrative medical database. *Appl Multivar Res* 2009; **12**:1-6.

- 1
2
3 216 Cassell A, Edwards D, Harshfield A, et al. The epidemiology of multimorbidity in
4 primary care: A retrospective cohort study. *Br J GenPract* 2018; **68**:e245-e51.
5 217 Wong ELY, Xu RH, Cheung AWL. Measuring the impact of chronic conditions and
6 associated multimorbidity on health-related quality of life in the general population in
7 hong kong sar, china: A cross-sectional study. *PLOS ONE* 2019; **14**:1-16.
8 218 Puth M-T, Weckbecker K, Schmid M, Münster E. Prevalence of multimorbidity in
9 germany: Impact of age and educational level in a cross-sectional study on 19,294
10 adults. *BMC Public Health* 2017; **17**:1-7.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



PRISMA Checklist

| Section/topic | # | Checklist item | Reported on page # |
|------------------------------------|----|---|------------------------|
| TITLE | | | |
| Title | 1 | Identify the report as a systematic review, meta-analysis, or both. | Page 1 |
| ABSTRACT | | | |
| Structured summary | 2 | Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number. | Page 2 |
| INTRODUCTION | | | |
| Rationale | 3 | Describe the rationale for the review in the context of what is already known. | Page 4 |
| Objectives | 4 | Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS). | Page 5 |
| METHODS | | | |
| Protocol and registration | 5 | Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number. | Page 2 |
| Eligibility criteria | 6 | Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. | Page 5 |
| Information sources | 7 | Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched. | Page 5 |
| Search | 8 | Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated. | Supplementary Table S1 |
| Study selection | 9 | State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis). | Page 6, Figure 1 |
| Data collection process | 10 | Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators. | Page 6 |
| Data items | 11 | List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made. | Supplementary Table S8 |
| Risk of bias in individual studies | 12 | Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis. | Page 6 Appendix p26 |
| Summary measures | 13 | State the principal summary measures (e.g., risk ratio, difference in means). | Page 6-8 |
| Synthesis of results | 14 | Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I ²) for each meta-analysis. | Page 6-8 |

| Section/topic | # | Checklist item | Reported on page # |
|---------------|---|----------------|--------------------|
|---------------|---|----------------|--------------------|



PRISMA Checklist

| | | | |
|-------------------------------|----|--|---|
| Risk of bias across studies | 15 | Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies). | Page 10 and Table 2 |
| Additional analyses | 16 | Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified. | Page 6-8 |
| RESULTS | | | |
| Study selection | 17 | Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram. | Figure 1 |
| Study characteristics | 18 | For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations. | Page 8-9, Table 1; Supplementary Table S4 |
| Risk of bias within studies | 19 | Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12). | Supplementary Table S6 |
| Results of individual studies | 20 | For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot. | Page 9-10 Figure 2-4 |
| Synthesis of results | 21 | Present results of each meta-analysis done, including confidence intervals and measures of consistency. | Page 9-10 Table 2 |
| Risk of bias across studies | 22 | Present results of any assessment of risk of bias across studies (see Item 15). | Page 9-10, Table 1 and Table 2 |
| Additional analysis | 23 | Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]). | Page 10-11 |
| DISCUSSION | | | |
| Summary of evidence | 24 | Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers). | Page 11,12 |
| Limitations | 25 | Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias). | Page 12 |
| Conclusions | 26 | Provide a general interpretation of the results in the context of other evidence, and implications for future research. | Page 13 |
| FUNDING | | | |
| Funding | 27 | Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. | Page 14 |

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

BMJ Open

Variation in the estimated prevalence of multimorbidity: systematic review and meta-analysis of 193 international studies

| | |
|---------------------------------|---|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2021-057017.R1 |
| Article Type: | Original research |
| Date Submitted by the Author: | 11-Mar-2022 |
| Complete List of Authors: | Ho, Iris Szu-Szu; The University of Edinburgh Usher Institute of Population Health Sciences and Informatics Azcoaga-Lorenzo, Amaya; University of St Andrews Bute Medical School Akbari, Ashley; Swansea University Medical School Davies, Jim; University of Oxford, Department of Computer Science Hodgins, Peter; The University of Edinburgh Usher Institute of Population Health Sciences and Informatics Khunti, Kamlesh; University of Leicester College of Life Sciences Kadam, Umesh; University of Leicester College of Life Sciences Lyons, Ronan; Swansea University, Health Data Research UK McCowan, Colin; University of St Andrews, School of Medicine Mercer, Stewart; The University of Edinburgh, Usher Institute Nirantharakumar, Krishnarajah; University of Birmingham Guthrie, Bruce; University of Edinburgh |
| Primary Subject Heading: | Global health |
| Secondary Subject Heading: | Public health, Geriatric medicine |
| Keywords: | EPIDEMIOLOGY, GENERAL MEDICINE (see Internal Medicine), GERIATRIC MEDICINE |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Variation in the estimated prevalence of multimorbidity: systematic review and meta-analysis of 193 international studies

Iris Szu-Szu Ho,¹ Amaya Azcoaga-Lorenzo,² Ashley Akbari,³ Professor Jim Davies,⁴ Peter Hodgins,¹ Kamlesh Khunti,⁵ Umesh Kadam,⁵ Ronan A Lyons,³ Colin McCowan,² Stewart W Mercer,¹ Krishnarajah Nirantharakumar,⁶ Bruce Guthrie.¹

1. University of Edinburgh, Usher Institute, Medical School, Teviot Place, Edinburgh, EH8 9AG, United Kingdom
2. University of St Andrews, School of Medicine, Medical and Biological Sciences, North Haugh, St Andrews, KY16 9TF, United Kingdom
3. Swansea University, Medical School, Population Health Science, Singleton Park, Swansea, SA2 8PP, United Kingdom
4. University of Oxford, Department of Computer Science, Wolfson Building, Parks Road, Oxford, OX1 3QD, United Kingdom
5. University of Leicester, Leicester General Hospital, Gwendolen Road, Leicester, LE5 4PW, United Kingdom
6. University of Birmingham, Institute of Applied Health Research, Birmingham, B15 2TT, United Kingdom

Corresponding author: Iris Szu-Szu Ho. Email: iris.s.ho@outlook.com

Abstract

Objective. (1) To estimate the pooled prevalence of multimorbidity in all age groups, globally. (2) To examine how measurement of multimorbidity impacted the estimated prevalence.

Methods. In this systematic review and meta-analysis, we conducted searches in nine bibliographic databases (PsycINFO, Embase, Global Health, Medline, Scopus, Web of Science, Cochrane Library, CINAHL, and ProQuest Dissertations & Theses Global) for prevalence studies published between database inception and 21 January 2020. Studies reporting the prevalence of multimorbidity (in all age groups and in community, primary care, care home and hospital settings) were included. Studies with an index condition or those that did not include people with no long-term conditions in the denominator were excluded. Retrieved studies were independently reviewed by two reviewers, and relevant data were extracted using pre-designed pro-forma. We used meta-analysis to pool the estimated prevalence of multimorbidity across studies, and used random-effects meta-regression and subgroup analysis to examine the association of heterogeneous prevalence estimates with study and measure characteristics.

Results. 13,807 titles were screened, of which 193 met inclusion criteria for meta-analysis. The pooled prevalence of multimorbidity was 42.4% (95%CI=38.9%-46.0%) with high heterogeneity ($I^2>99\%$). In adjusted meta-regression models, participant mean age and the number of conditions included in a measure accounted for 47.8% of heterogeneity in effect sizes. The estimated prevalence of multimorbidity was significantly higher in studies with older adults and those that included larger numbers of conditions. There was no significant difference in estimated prevalence between low- or middle-income countries (36.8%) and high-income countries (44.3%), or between self-report (40.0%) and administrative/clinical databases (52.7%).

Conclusions. The pooled prevalence of multimorbidity was significantly higher in older populations and when studies included a larger number of baseline conditions. The findings suggest that, to improve study comparability and quality of reporting, future studies should use a common core conditions set for multimorbidity measurement and report multimorbidity prevalence stratified by socio-demographics.

Funding: HDRUK (CFC0110)

Review registration: PROSPERO (CRD42020172409)

Strengths and limitations of this study

- This study used meta-regression to examine the variation of estimated prevalence of multimorbidity and how measure and study characteristics influenced prevalence estimates.
- The use of multiple imputation in this study minimised biased estimates caused by missing values and unbalanced classes and enhanced statistical accuracy.
- The inclusion of studies with various measure and study characteristics enabled a better understanding of the contributing factors of the heterogeneity of multimorbidity prevalence.
- Due to inconsistent reporting of multimorbidity prevalence and data unavailability, the estimated multimorbidity prevalence stratified by sex, ethnicity and socio-economic status could not be explored in this study.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Population ageing is a worldwide phenomenon, with the World Health Organization (WHO, 2018) estimating that the proportion of the global population aged 60 and older will double from 12% to 22% between 2015 and 2050 [1]. A key implication of population ageing is that increasing numbers of people will be living with multimorbidity. Multimorbidity, commonly defined as the co-occurrence of two or more long-term conditions [2], adversely affects people’s risk of death, health-related quality of life, functional ability, and mental well-being [3, 4]. Multimorbidity affects all groups of society, but is known to be more common in older people, in women, and in those from low socio-economic backgrounds particularly in high-income countries [5-7]. In low- and middle-income countries, people living in urban areas, on the other hand, were found to have a higher rate of multimorbidity prevalence [8]. Multimorbidity poses major challenges to the delivery of care in health systems internationally which are often focused on the management of single diseases and lack appropriate coordination and continuity of care across different sectors [9, 10]. Disparities in health and health and social care could be found at any stage along the continuum of chronic diseases, from prevention to the management of diseases. To understand these disparities among multimorbid populations requires consistently monitoring the populations (e.g. incidence, prevalence, health impact, risk factors and delivery of care) defined by race and ethnicity, gender, age, socio-economic status, physical environment and geographic factors.

Previous systematic reviews have identified issues in the measurement of multimorbidity, related to the choice of chronic conditions counted in measures, the categorisation of conditions and diseases, and the counting or weighting method used [11-13]. Although weighted measures are often used when the purpose of measurement is to predict future outcomes, a simple count of conditions remains the most commonly-used method for the measurement of multimorbidity, and is optimal for estimating multimorbidity prevalence [13, 14]. However, the estimated prevalence of multimorbidity varies widely in the literature ranging from 3.5% to 100% [15], likely reflecting a combination of varying measures and varying populations studied [16]. Much of the research up to now has not quantitatively investigated the variation in multimorbidity prevalence and its influencing factors in much detail. Understanding the links between prevalence estimates and measurement approaches can better inform and support future development of multimorbidity measurement guidelines. Therefore, this review aimed to examine the pooled prevalence of

multimorbidity in all age groups, globally and how measurement of multimorbidity impacted the estimated prevalence.

Research questions

- What is the pooled prevalence of multimorbidity and does it differ between different age groups?
- What are the factors that influenced the variation in prevalence estimates across studies?

Methods

The systematic review and meta-analysis reported here is part of a larger review which aimed to examine 1) how multimorbidity has been constructed and 2) measured by international studies and 3) variation in the estimated prevalence of multimorbidity across studies. Analysis in relation to the first two registered objectives has been reported [13], and this paper reports the third registered objective. The PROSPERO registration number for this paper is therefore the same as for the first published paper from this work [13].

Inclusion and exclusion criteria

The eligibility criteria for this review were defined based on the CoCoPop framework—Condition, Context, and Population [17]. The condition included in this review is prevalence of multimorbidity. The majority of studies defined multimorbidity as the co-existence of two or more chronic conditions, and used the cut-off to estimate its prevalence in a population of interest. We therefore included studies that used this definition for examining multimorbidity prevalence across international studies. For this analysis, we included studies carried out in the community, primary care, care home and hospitals, and those estimating the prevalence of multimorbidity in the population studied. Studies that did not include a relevant denominator population – for example, only examining patients with an index condition or excluding patients who did not have multimorbidity – were excluded. Qualitative research, studies not published in English, and conference abstracts were also excluded.

Search strategy

The search strategy for this review was developed in collaboration with a specialist medical librarian (Supplementary Table S1). Key terms relevant to multimorbidity and measurement were

combined using Boolean logic to identify studies that met the inclusion criteria. We included medical subject headings to provide a sensitive search for relevant literature. Databases included in the search were Ovid interface (PsycINFO, Embase, Global Health, Medline), Scopus, Web of Science, Cochrane Library, EBSCO interface (CINAHL Plus), and ProQuest Dissertations & Theses Global, from inception to 21 January 2020 (we are not aware of any large recently published studies since that date). In addition to the database searches, our secondary search strategy included hand-searching reference lists of retrieved articles and tracked citations to maximise the yield.

Study screening and selection

Articles retrieved from databases were organised using EndNote X9 bibliographic software and Excel, and then were imported to Covidence for screening [18]. Titles, abstracts, and full-texts of retrieved articles were screened against the eligibility criteria by two reviewers. Throughout the review process, any disagreement that arose was resolved through discussion between the two reviewers (IS-SH and PH), and through the involvement of a third reviewer (BG) if necessary. The study selection process is summarised in Figure 1.

Data extraction

We extracted data on the characteristics of the included studies using pre-designed data extraction pro-forma. The extracted data include 1) authors, 2) publication year, 3) study purpose, 4) method, 5) country, 6), continent, 7) country income (classified as ‘high’ and ‘low or medium’ [combined because of small numbers] allocated based on the World Bank Group at the time of review [19]), 8) study participants, 9) mean age, 10) sample size, 11) number of conditions, 12) setting, 13) data collection method/data source, 14) number of multimorbidity cases, and 15) proportion of multimorbidity (calculated based on item 10 and 14). Data on the estimated prevalence stratified by sex, ethnicity and socio-economic status were fragmented and unavailable in many studies, and thus these could not be retrieved for analyses.

Risk of bias assessment

We used the Effective Public Health Practice Project (EPHPP) quality assessment tool for quantitative studies to assess the risk of bias and the quality of each of the included studies, in terms of 1) selection bias, 2) study design, 3) confounders, 4) blinding, 5) data collection method, 6) withdrawals and dropouts [20]. We assessed also publication bias (rated high if there was

selective reporting within studies) and conflict of interest (rated unclear if conflict of interest declaration was not reported). Each study was rated and assigned an overall risk of bias as 'high', 'moderate', or 'low' (please see the details in appendix p26).

Data analysis

Descriptive statistics were used to summarise study characteristics. Since distributions were skewed, median and interquartile range were used to measure the central tendency and examine variability of variables such as mean age and number of conditions. Categorical (e.g. continent, study population, and data source) and ordinal data (e.g. country income and risk of bias) were examined using frequency tables. To investigate the association between continuous/count predictor (mean age/number of conditions) and categorical predictors, univariate generalized linear models were used. We summarised the prevalence of multimorbidity using metaprop [21, 22]. The presence of effect size heterogeneity was examined using the Q statistic and I-squared. Significant heterogeneity was identified, so we used subgroup analysis and meta-regression with random-effects models to identify potential moderating factors.

Outlying studies were identified using studentised residuals, leave-one-out analysis and Mahalanobis distance. Studies with studentised residuals that were larger than 2 or 3 and those that contributed to heterogeneity in leave-one-out analyses were scrutinized [23]. Mahalanobis distance was used for pattern recognition and multivariate outlier detection [24]. Study effect sizes were graphically displayed to identify outlying studies and explore subgroup effects (Supplementary Figure S1). In initial analysis of heterogeneity and outliers, 24 studies were found to make a significant contribution to the high level of observed heterogeneity in multimorbidity prevalence and significant changes in the summary effect size. The 24 studies were excluded for one or more of the following reasons: 1) their contribution to high levels of heterogeneity in the leave-one-out test, 2) being identified as an outlying value in the studentised residuals test ($z\text{-score} \geq 2$), 3) their Mahalanobis distance exceeding the chi-squared critical value at a 0.01 significance level, 4) infrequent values in compositional categorical data (e.g. only one study examined prevalence in children). The process of identifying outliers, the rationale for exclusion of each study, and the characteristics of outlying studies are documented in Supplementary Figure S2 and Table S2 and Table S3. Sensitivity analysis was performed to explore the impact of excluding the 24 studies in meta-analysis.

There was missingness in two predictors, with 37% missingness in the ‘mean age’ of the study population variable (some of which reported it categorically, and thus were treated as missing data) and 6% missingness in the ‘number of conditions’ included in the multimorbidity measure variable. Previous research has shown that complete case removal (removing missing data in a data set) in meta-regression could lead to biased coefficient estimates of predictors (varied widely from complete-data estimates), whereas multiple imputation was found to perform well at generating estimates that were close to complete-data estimates [25]. Therefore, in this review, multiple imputation with 60 imputed datasets and 10 iterations was conducted where random forest was used to impute missing data [26, 27]. Following multiple imputation, fraction of missing information (FMI) was computed to quantify the impact of missing data, which ranged from 0.05 to 0.3 indicating that the uncertainty in the values imputed for missing data is small/moderate [28].

A random-effects regression tree approach with ten-fold cross-validation was used to identify subgroups (cut-offs) of the ‘mean age’ and ‘number of conditions’ variables with differential effect sizes [29]. Given considerable variation in the effect sizes, we conducted meta-regression with the restricted maximum likelihood (REML) estimator to examine the possible sources of heterogeneity in effect sizes [21, 22, 30]. As the variable ‘multimorbidity prevalence’ did not follow the normal distribution (positively skewed), we applied logit transformation to the variable for analyses and converted the logits back to odds ratios (e^{logit}) and proportions ($p = e^{\text{logit}} / (e^{\text{logit}} + 1)$) for reporting. For model selection, we refitted the models using maximum likelihood and then conducted a log-likelihood test to compare the fit of models [31]. A permutation test with 1000 permuted datasets was conducted to validate the robustness of the final model by rearranging and shuffling the order of the data and re-calculating p-values to check whether there is type 1 error [32]. Subgroup analysis with the REML method was used to estimate the pooled multimorbidity prevalence of subgroups of each variable (age, the number of conditions included in a measure, setting, data source, continent, country income, study risk of bias). Forest-like plots were used to display the effect sizes of included studies [33]. The presence of publication bias was assessed using Egger’s test, which did not find evidence of publication bias [34]. All statistical tests were performed using R version 4.0.4.

Patients and public involvement

No patients were involved in the development of the research question, outcome measures, study design and implementation. Nonetheless, we have previously discussed preliminary review findings and issues relevant to multimorbidity measurement with our patient and public involvement group. We plan to disseminate the review findings to researchers, clinicians, policy makers and public audiences through news media, social media and seminars.

Results

After screening 13,807 titles and abstracts, 217 studies were identified which estimated the prevalence of multimorbidity using a cut-off of 'two or more' conditions. Following the removal of 24 outlying studies, 193 studies were included in the meta-analysis (Table 1, Supplementary Table S4). Of the 193 studies, 64 studies were from Europe, 47 from North America, 44 from Asia, 11 from Australasia, 12 from South America, and four from Africa (Table 1 and Figure 2).

Seventy-five percent of studies were from high-income countries (n=145) and 24.9% from low- and middle-income countries (LMICs) (one from low-income, eight from lower middle-income, 29 from upper middle-income, and 10 from multiple low- and middle-income countries). The majority of studies (n=147) estimated the prevalence of multimorbidity in community settings, followed by primary care (n=32) and hospital setting (n=14). Prevalence data were collected through either self-report (n=150) or medical records and administrative databases (n=43). In a univariate linear regression (Supplementary Table S5), we found that studies from Europe, database studies and studies conducted in hospital settings were more likely to measure multimorbidity in an older population and included a larger number of conditions in a multimorbidity measure, compared to those from other continents, self-report studies, and studies conducted in primary care and community settings. In respect to risk of bias in included studies (Supplementary Table S6 and Figure S3), 11.4% were rated as high risk of bias, 83.9% as moderate risk of bias, and 4.7% as low risk of bias.

The pooled estimate of multimorbidity prevalence across the 193 studies was 42.4% (95%CI 38.9%-46.0%), τ^2 is 1.0 (95%CI 0.9-1.3) with high heterogeneity ($I^2>99\%$), and meta-regression was therefore used to examine study characteristics associated with heterogeneity. Mean age ($F=89.8$, $p<0.0001$, $R^2=31.7\%$) and number of conditions ($F=39.2$, $p<0.0001$, $R^2=16.7\%$) were the strongest univariate predictors and positively associated with the estimated prevalence of

multimorbidity (Figure 3). Meta-regression tree analysis (Supplementary Figure S4) partitioned the mean age variable into three homogeneous subgroups (aged <59, aged 59-73, aged ≥74) and the number of conditions variable into four homogeneous subgroups (<9, 9-19, 20-43, ≥44). The categorical ‘mean age’ and ‘number of conditions’ variables explained 35.9% and 19.5% of the heterogeneity in effect sizes respectively (larger than the original numerical variables). Therefore, the categorical variables identified from the regression trees for meta-analyses were used for meta-regression.

In univariate meta-regression, primary care studies (pooled multimorbidity prevalence 50.5%, OR 1.6, 95%CI 1.1-2.3) and hospital based studies (pooled multimorbidity prevalence 59.6%, OR 2.3, 95%CI 1.3-4.0) had significantly higher rates of multimorbidity than community-based studies (39.1%) (Table 2). Multimorbidity prevalence was significantly higher in database studies (pooled multimorbidity prevalence 52.7%, OR 1.7, 95%CI 1.2-2.4) than self-report studies (pooled multimorbidity prevalence 40.0%). In the mean age categorical variable, the pooled prevalence estimates of the three subgroups were statistically significantly different from one another, and considerably higher in studies with mean participant age ≥74 (pooled multimorbidity prevalence 67.0%, OR 5.2, 95%CI 3.8-7.2) and mean participant age 59-73 (pooled multimorbidity prevalence 47.6%, OR 2.3, 95%CI 1.8-3.0) than those with mean participant age <59 (pooled multimorbidity prevalence 28.0%) (Table 2 and Figure 4). Similar patterns were also found in the number of conditions variable where studies including ≥44 conditions in measurement (pooled multimorbidity prevalence 87.6%, OR 16.5, 95%CI 6.4-42.6), 20-43 conditions (pooled multimorbidity prevalence 52.1%, OR 2.5, 95%CI 1.7-3.7), and 9-19 conditions (pooled multimorbidity prevalence 43.7%, OR 1.8, 95%CI 1.3-2.5) yielded higher prevalence estimates than studies including <9 conditions in measurement (pooled multimorbidity prevalence 30.1%) with a dose-response relationship. The estimated prevalence of multimorbidity was 44.3% in high-income countries compared to 36.8% in low or middle income countries, but the difference was not statistically significantly different (OR 1.4, 95%CI 1.0-1.9). In study risk of bias, no statistically significant difference in pooled prevalence of multimorbidity was found between studies with low, moderate and high risk of bias.

In the adjusted meta-regression model, compared to studies where participant mean age was <59, multimorbidity prevalence remained significantly higher in studies with mean participant age 59-73 (OR 2.2, 95%CI 1.7-2.8) and in studies with mean participant age ≥74 (OR 4.4, 95%CI 3.3-

5.9). Compared to measures including <9 conditions, multimorbidity prevalence was higher in measures including ≥ 44 conditions (OR 8.2, 95%CI 3.8-17.5), 20-43 conditions (OR 2.3, 95%CI 1.6-3.2), and 9-19 conditions (OR 1.8, 95%CI 1.4-2.3). In respect to study settings, the pooled prevalence was significantly higher in primary care settings compared to community settings (OR 1.6, 95%CI 1.1-2.3). Compared to studies from North America, prevalence was lower in studies from Europe (OR 0.5, 95%CI 0.4-0.7), Australasia (OR 0.5, 95%CI 0.3-0.8), Asia (OR 0.6, 95%CI 0.4-0.8), or Africa (OR 0.3 95%CI 0.1-0.6). No significant difference in prevalence estimates between self-report and routine database studies was evident after controlling for study and measure characteristics. The model explained 54.3% of the heterogeneity in multimorbidity prevalence, with the mean age and number of conditions variables providing most explanatory power (47.8% of the heterogeneity).

Sensitivity analysis including the 24 outlying studies (Supplementary Table S7) was similar to primary analysis except for “number of conditions” variable. The mean participant age and number of conditions variables remained the strongest predictors of multimorbidity prevalence in sensitivity analysis. However, the estimated prevalence in sensitivity analysis (including outlying studies) was much lower in studies including ≥ 44 conditions in a multimorbidity measure (pooled multimorbidity prevalence 54.5, OR 2.8, 95%CI 1.5-5.4) compared to primary analysis excluding outlying studies (pooled multimorbidity prevalence 87.6, OR 16.5, 95%CI 6.4-42.6). The difference in estimates was mainly attributed to the three outlying studies that included 146, 147 and 259 conditions in a measure respectively but yielded relatively low mean multimorbidity prevalence (mean prevalence 54.3%)[35-37].

Discussion

The overall estimate of multimorbidity prevalence in adults across all the included studies was 42.4% (95%CI=38.9%-46.0%), but with very high heterogeneity. More than half of the observed heterogeneity was explained by study mean participant age and the number of conditions included in the multimorbidity measure, with older age and larger number of conditions strongly associated with a higher prevalence of multimorbidity. The difference in estimated prevalence was small between self-report and administrative/clinical databases, and between study settings. No significant difference was found between studies from low- or middle-income and high-income

countries, but North American studies had higher estimated prevalence and African studies had the lowest estimated prevalence than other continents.

Three prior systematic reviews examined the prevalence of multimorbidity across studies [38-40]. Fortin et al. (2012) and Violan et al (2014) conducted a narrative review and found various operationalisations of multimorbidity and a large variation in the prevalence of multimorbidity, particularly in studies with older adult populations or those with low socioeconomic status [38, 40]. Nguyen et al. (2019) meta-analysed the prevalence of multimorbidity across 70 studies from community settings and found that the pooled estimated prevalence was 33.1% with high levels of heterogeneity ($I^2 > 99\%$) [39]. The pooled prevalence of multimorbidity in Nguyen et al study is lower than in this study, likely because we have included studies from primary care and hospital settings (the pooled prevalence of multimorbidity in community-based studies in this analysis was 39.5%). Nguyen et al. (2019) did not carry out a meta-regression, but in narrative analysis comment that the prevalence of multimorbidity appeared higher in older adults and women [39]. Our review findings are consistent with previous literature finding that age is most important determinant of multimorbidity [5, 38, 39, 41]. While we did not find a significant difference between low and middle-income and high-income countries, Nguyen et al. in their review showed a statistically significantly higher pooled prevalence in high-income countries (the pooled prevalence from 18 studies was 37% compared to 36.8% in this review of 145 studies) than low or middle-income countries (the pooled prevalence from 31 studies was 29% compared to 44.3% in this review of 48 studies). This difference in findings may be due to the inclusion in our review of a larger number of studies from high-income or upper middle-income countries. The low number of included studies from low-income countries in this review could be explained by fewer attention paid to this relatively new research field (multimorbidity) in low-income countries and our literature search restricted to English language (proficient language of reviewers). The estimated prevalence of multimorbidity in North America was higher compared to other continents in this study despite older study populations and larger numbers of conditions found in studies from Europe. A possible explanation for the higher prevalence in North America is that private or insurance-based healthcare systems are more likely to code conditions since it affects remuneration, as well as cultural differences in relation to over-diagnosis and medicalisation [42]. On the other hand, the lower estimated multimorbidity prevalence in African studies could be attributed to the

predominance of infectious diseases and inadequate access to medical care including diagnostic services [43].

The strengths of this review are searches conducted in multiple databases, the large number of studies identified and the use of meta-analytic approaches to examine factors associated with heterogeneity of estimated multimorbidity prevalence. We examined and handled outlying studies and missing data (multiple imputation) with rigour and excluded studies that did not take into account 'healthy' populations (populations with no long-term conditions) to minimize biased estimates of multimorbidity prevalence. This review has limitations. Sensitivity analysis including all studies had similar findings with one exception, namely that sensitivity analysis found: a weaker (but still statistically significant) association with the number of conditions included in the multimorbidity measure than primary analysis. Although we examined associations with study characteristics including mean participant age, a limitation is the lack of information in the reviewed studies on prevalence estimates stratified by participant characteristics including sex, ethnicity, and socio-economic status. An additional uncontrolled factor is how studies measured multimorbidity in terms of the type (as opposed to the number) of the conditions included in measures, which varied substantially across studies with too much heterogeneity to model [13]. The exclusion of non-English studies in this review may also limit the generalisability of the research findings. Last but not least, measurement of multimorbidity is a relatively new research field and its labelling has been used variably. Thus, it is likely that not all relevant studies were identified and included in this review, but we were rigorous in our application of inclusion/exclusion criteria and did not favour adding known papers that did not appear in the search or were excluded through the process.

In spite of the methodological limitations, this review adds to our understanding of how study and measure characteristics can influence the estimated prevalence of multimorbidity. Mean age of the study population and the number of conditions included in the multimorbidity measure were the major factors associated with varying estimated prevalence of multimorbidity. A key implication is that comparing prevalence between studies requires more stratified estimates of multimorbidity prevalence. We therefore strongly recommend that as well as overall prevalence, future studies should clearly report multimorbidity prevalence stratified by age, in 5-year age bands to ensure granularity, and by sex at a minimum, and ideally by ethnicity and socio-economic status. This will allow readers to capture a more holistic picture of multimorbidity prevalence in the population

studied, and allow better comparison of prevalence in different populations, and accurate pooled estimates of prevalence in reviews.

Additionally, the number of conditions included in a measure is strongly associated with estimated multimorbidity prevalence. It would be ideal if studies additionally reported prevalence using a common core set of conditions agreed by consensus. Parallel reporting of the bespoke set chosen for the context and purpose, and a core set would improve comparability of prevalence estimates, and help identify the additional value of any bespoke multimorbidity measures. The lack of any significant difference in estimated prevalence between self-report and clinical/administrative databases in this review suggests that provided careful attention is paid to the number and type of conditions included in measures, exactly how data is collected may be less important.

To conclude, in recent years, there has been an increasing interest in the epidemiology of multimorbidity internationally. This review finds that population characteristics and measurement content are the major factors that influenced prevalence estimates of multimorbidity. Studies with older populations and larger numbers of conditions yielded a higher estimate of multimorbidity prevalence. However, heterogeneity between studies has made comparison of multimorbidity prevalence across studies difficult. To improve comparability and quality of reporting, this review suggests that future studies should use common core condition set for the measurement of multimorbidity and clearly report the prevalence of multimorbidity stratified by socio-demographics.

Contributorship statement

All authors have made substantial contributions: CMC, KN, UK, KK, RAL, JD, AA, AAL and SWM were involved in conception of the work, acquisition of funding, and critically commenting on the manuscript. IS-SH led and BG substantially contributed to the design, analysis, and interpretation of data for the review, and are responsible for the decision to submit the manuscript. IS-SH and PH screened and reviewed retrieved studies. All authors contributed to the edits of the manuscript and had access to the data. The final draft has been approved by all authors.

Competing interests

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: we had financial support from HDRUK for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Funding

This project was funded by Health Data Research UK (HDRUK: CFC0110).

Data sharing statement

Study data are available in supplementary appendix.

Figure legends

Figure 1: Preferred reporting items for systematic reviews and meta-analyses (PRISMA) flow diagram

Figure 2: Country of origin of the included studies estimating the prevalence of multimorbidity (except studies from multiple countries)

Figure 3: Relationship between the prevalence of multimorbidity and mean age or number of conditions (the area of points is proportional to inverse variances)

Figure 4: The distribution of prevalence estimates within the subgroups of mean age and number of conditions (forest-like plot for a large review)

Table 1: Summary of study characteristics (Supplementary Table S8 shows the definition of variables)

| Name of variable | Descriptive statistics (n=193) |
|--|---|
| Prevalence of multimorbidity (%) | Range: 2.7 to 95.6 Pooled prevalence with the REML estimator: 42.4 (38.9-46.0) |
| Mean age of study population (year) | Range of mean age: 32.2-83.8 Median of mean age: 62.6 (Q1, Q3: 50.1, 72.4) |
| No of conditions (count) | Range: 3-60 Median: 13 (Q1, Q3: 9, 19) |
| Country income (count, %) High income Low- or Middle-income | 145 (75.1%) 48 (24.9%) |
| Continent (count, %) Europe North America Asia Australasia South America Africa Multiple continents | 64 (33.2%) 47 (24.4%) 44 (22.8%) 11 (5.7%) 12 (6.2%) 4 (2.1%) 11 (5.7%) |
| Study population (count, %) Only older people Middle-aged and older All adults | 63 (32.6%) 46 (23.8%) 84 (43.5%) |
| Setting (count, %) Community Primary care Hospital | 147 (76.2%) 32 (16.6%) 14 (7.3%) |
| Source (count, %) Self-report Database | 150 (77.7%) 43 (22.3%) |
| Risk of bias assessment (count, %) Low Moderate High | 9 (4.7%) 162 (83.9%) 22 (11.4%) |

IQR: Interquartile range. SD: Standard deviation. The percentages were rounded so they do not add to 100%.

Table 2: Output of meta-analytic models (n=193)

| | Pooled prevalence of multimorbidity of each subgroup (%, 95% CI) | Meta-regression Unadjusted Odds Ratio (95% CI) | Meta-regression Adjusted Odds Ratio (95% CI) R ² 54.3% | FMI |
|---------------------------|--|--|--|------|
| Group of mean age | | R ² 35.9% | | |
| <59 | 28.0 (24.9-31.5) | Ref | Ref | Ref |
| 59-73 | 47.6 (42.5-52.8) | 2.3 (1.8-3.0)*** | 2.2 (1.7-2.8)*** | 0.3 |
| ≥74 | 67.0 (60.4-72.9) | 5.2 (3.8-7.2)*** | 4.4 (3.3-5.9)*** | 0.2 |
| No of conditions | | R ² 19.5% | | |
| <9 | 30.1 (24.9-35.7) | Ref | Ref | Ref |
| 9-19 | 43.7 (39.5-48.0) | 1.8 (1.3-2.5)*** | 1.8 (1.4-2.3)*** | 0.1 |
| 20-43 | 52.1 (43.8-60.3) | 2.5 (1.7-3.7)*** | 2.3 (1.6-3.2)*** | 0.2 |
| ≥44 | 87.6 (81.3-92.0) | 16.5 (6.4-42.6)*** | 8.2 (3.8-17.5)*** | 0.06 |
| Setting | | R ² 5.1% | | |
| Community | 39.1 (35.5-42.8) | Ref | Ref | Ref |
| Primary care | 50.5 (39.6-61.3) | 1.6 (1.1-2.3)* | 1.6 (1.1-2.3)** | 0.2 |
| Hospital | 59.6 (45.6-72.2) | 2.3 (1.3-4.0)** | 1.5 (1.0-2.4) | 0.2 |
| Source | | R ² 4.0% | | |
| Self-report | 40.0 (36.2-43.8) | Ref | Ref | Ref |
| Database | 52.7 (45.2-60.1) | 1.7 (1.2-2.4)** | 0.7 (0.5-1.0) | 0.2 |
| Continent | | R ² 6.8% | | |
| North America | 50.4 (43.6-57.3) | Ref | Ref | Ref |
| Europe | 44.8 (38.2-51.5) | 0.8 (0.5-1.2) | 0.5 (0.4-0.7)*** | 0.1 |
| Australasia | 35.8 (29.5-42.5) | 0.5 (0.3-1.1) | 0.5 (0.3-0.8)** | 0.08 |
| Asia | 35.3 (29.3-42.0) | 0.5 (0.4-0.8)** | 0.6 (0.4-0.8)*** | 0.1 |
| South America | 47.5 (31.2-64.4) | 0.9 (0.5-1.7) | 0.8 (0.5-1.3) | 0.1 |
| Africa | 13.8 (4.5-32.8) | 0.2 (0.06-0.4)*** | 0.3 (0.1-0.6)** | 0.1 |
| Multiple continents | 38.4 (29.1-48.6) | 0.6 (0.3-1.2) | 0.7 (0.4-1.1) | 0.1 |
| Country income | | R ² 1.2% | | |
| Low or middle-income | 36.8 (29.7-44.4) | Ref | | |
| High-income | 44.3 (40.3-48.4) | 1.4 (1.0-1.9) | | |
| Study risk of bias | | R ² 0.0% | | |
| Low risk | 33.3 (20.2-49.6) | Ref | | |
| Moderate risk | 42.4 (38.6-46.3) | 1.5 (0.7-3.0) | | |
| High risk | 46.4 (34.1-59.1) | 1.7 (0.8-3.9) | | |
| Publication year | | 1.0 (1.0-1.0) | | |

* <0.05 ** <0.01 *** <0.001

Ref: Reference category. FMI: Fraction of missing information

References

1. World Health Organization. *Ageing and health*. 2018 [cited 2021 April 1]; Available from: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>.

2. The Academy of Medical Sciences, *Multimorbidity: a priority for global health research*. 2018.

3. Rivera-Almaraz, A., et al., *Disability, quality of life and all-cause mortality in older Mexican adults: association with multimorbidity and frailty*. BMC Geriatr, 2018. **18**(236): p. 1-9.

4. Arokiasamy, P., et al., *The impact of multimorbidity on adult physical and mental health in low- and middle-income countries: what does the study on global ageing and adult health (SAGE) reveal?* BMC Med, 2015. **13**(178): p. 1-16.

5. Barnett, K., et al., *Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study*. The Lancet, 2012. **380**(9836): p. 37-43.

6. Low, L.L., et al., *Epidemiologic Characteristics of Multimorbidity and Sociodemographic Factors Associated With Multimorbidity in a Rapidly Aging Asian Country*. JAMA Network Open, 2019. **2**(11): p. 1-14.

7. Fortin, M., et al., *Prevalence estimates of multimorbidity: a comparative study of two sources*. BMC health services research, 2010. **10**(111): p. 1-6.

8. Asogwa, O.A., et al., *Multimorbidity of non-communicable diseases in low-income and middle-income countries: a systematic review and meta-analysis*. BMJ Open, 2022. **12**(1): p. 1-16.

9. Pearson-Stuttard, J., M. Ezzati, and E.W.J.T.L.P.H. Gregg, *Multimorbidity—a defining challenge for health systems*. The Lancet, 2019. **4**(12): p. e599-e600.

10. Moffat, K. and S.W.J.B.f.p. Mercer, *Challenges of managing people with multimorbidity in today’s healthcare systems*. BMC Family Practice, 2015. **16**(1): p. 1-3.

11. Johnston, M.C., et al., *Defining and measuring multimorbidity: a systematic review of systematic reviews*. Eur J Public Health, 2019. **29**(1): p. 182-189.

12. Diederichs, C., K. Berger, and D.B. Bartels, *The measurement of multiple chronic diseases--a systematic review on existing multimorbidity indices*. J Gerontol A Biol Sci Med Sci, 2011. **66**(3): p. 301-11.

13. Ho, I.S.-S., et al., *Examining variation in the measurement of multimorbidity in research: a systematic review of 566 studies*. The Lancet Public Health, June 21, 2021 (published online: [https://www.thelancet.com/journals/lanpub/article/PIIS2468-2667\(21\)00107-9/fulltext](https://www.thelancet.com/journals/lanpub/article/PIIS2468-2667(21)00107-9/fulltext)). **6**(1): p. e587-e597.

14. Huntley, A.L., et al., *Measures of multimorbidity and morbidity burden for use in primary care and community settings: a systematic review and guide*. Ann Fam Med, 2012. **10**(2): p. 134-41.

15. Xu, X., G.D. Mishra, and M. Jones, *Evidence on multimorbidity from definition to intervention: An overview of systematic reviews*. Ageing Res Rev, 2017. **37**: p. 53-68.

16. Harrison, C., et al., *Examining different measures of multimorbidity, using a large prospective cross-sectional study in Australian general practice*. BMJ Open, 2014. **4**(7): p. 1-9.

17. Munn, Z., et al., *Methodological guidance for systematic reviews of observational epidemiological studies reporting prevalence and cumulative incidence data*. Int J Evid-based Hea, 2015. **13**(3): p. 147-153.
18. Covidence. *Covidence 2020* [cited 2020 January 14]; Available from: <https://www.covidence.org/>.
19. World Bank Group. *The world by income*. 2018 [cited 2021 01 June]; Available from: <https://datatopics.worldbank.org/world-development-indicators/the-world-by-income-and-region.html>.
20. Effective Public Health Practice Project. *Quality Assessment Tool for Quantitative Studies Dictionary*. 2009 [cited 2020 14 Jan]; Available from: https://www.ephpp.ca/PDF/QADictionary_dec2009.pdf.
21. Viechtbauer, W. *Package 'metafor'*. 2020 [cited 2020 August 20]; Available from: <https://cran.r-project.org/web/packages/metafor/metafor.pdf>.
22. Schwarzer, G. *Package 'meta'*. 2021 [cited 2021 Feb 15]; Available from: <https://cran.r-project.org/web/packages/meta/meta.pdf>.
23. Viechtbauer, W. and M.W.L. Cheung, *Outlier and influence diagnostics for meta-analysis*. Research Synthesis Methods, 2010. **1**(2): p. 112-125.
24. Leys, C., et al., *Detecting multivariate outliers: Use a robust variant of the Mahalanobis distance*. Journal of Experimental Social Psychology, 2018. **74**: p. 150-156.
25. Ellington, E.H., et al., *Using multiple imputation to estimate missing data in meta-regression*. Methods in Ecology and Evolution, 2015. **6**(2): p. 153-163.
26. Tang, F. and H. Ishwaran, *Random forest missing data algorithms*. Statistical Analysis and Data Mining: The ASA Data Science Journal, 2017. **10**(6): p. 363-377.
27. van Buuren, S. *mice: Multivariate Imputation by Chained Equations. R package version: 4.0.1*. 2020 [cited 2020 1 Aug]; Available from: <https://cran.r-project.org/web/packages/mice/mice.pdf>.
28. White, I.R., P. Royston, and A.M. Wood, *Multiple imputation using chained equations: Issues and guidance for practice*. Stat Med, 2011. **30**(4): p. 377-99.
29. Li, X., et al. *Package 'metacart'*. 2020 [cited 2020 May 20]; Available from: <https://cran.r-project.org/web/packages/metacart/metacart.pdf>.
30. Langan, D., et al., *A comparison of heterogeneity variance estimators in simulated random-effects meta-analyses*. Res Synth Methods, 2019. **10**(1): p. 83-98.
31. Verbyla, A.P., *A note on model selection using information criteria for general linear models estimated using REML*. Australian & New Zealand Journal of Statistics, 2019. **61**(1): p. 39-50.
32. Gagnier, J.J., et al., *An empirical study using permutation-based resampling in meta-regression*. Systematic reviews, 2012. **1**(1): p. 1-9.
33. Nakagawa, S., et al., *The orchard plot: Cultivating a forest plot for use in ecology, evolution, and beyond*. Research Synthesis Methods, 2021. **12**(1): p. 4-12.
34. Shi, X., et al., *Effect comparison between Egger's test and Begg's test in publication bias diagnosis in meta-analyses: evidence from a pilot survey*. International Journal of Research Studies in Biosciences, 2017. **5**(5): p. 14-20.
35. Violán, C., et al., *Burden of multimorbidity, socioeconomic status and use of health services across stages of life in urban areas: a cross-sectional study*. BMC public health, 2014. **14**(530): p. 1-13.

36. Prazeres, F. and L. Santiago, *Prevalence of multimorbidity in the adult population attending primary care in Portugal: a cross-sectional study*. BMJ Open, 2015. **5**(9): p. 1-10.

37. Sullivan, P.W., V.H. Ghushchyan, and E.A. Bayliss, *The impact of co-morbidity burden on preference-based health-related quality of life in the United States*. Pharmacoeconomics, 2012. **30**(5): p. 431-42.

38. Fortin, M., et al., *A systematic review of prevalence studies on multimorbidity: toward a more uniform methodology*. Ann Fam Med, 2012. **10**(2): p. 142-51.

39. Nguyen, H., et al., *Prevalence of multimorbidity in community settings: A systematic review and meta-analysis of observational studies*. J Comorb, 2019. **9**: p. 1-15.


40. Violan, C., et al., *Prevalence, Determinants and Patterns of Multimorbidity in Primary Care: A Systematic Review of Observational Studies*. PLOS ONE, 2014. **9**(7): p. 1-9.

41. Hu, R.-H., et al., *Increasing age- and gender-specific burden and complexity of multimorbidity in Taiwan, 2003–2013: a cross-sectional study based on nationwide claims data*. BMJ Open, 2019. **9**(6): p. 1-10.



42. Kale, M.S. and D. Korenstein, *Overdiagnosis in primary care: framing the problem and finding solutions*. BMJ, 2018. **362**(k2820): p. 1-48.



43. Gouda, H.N., et al., *Burden of non-communicable diseases in sub-Saharan Africa, 1990–2017: results from the Global Burden of Disease Study 2017*. The Lancet Global Health, 2019. **7**(10): p. e1375-e1387.

Figure 1: Preferred reporting items for systematic reviews and meta-analyses (PRISMA) flow diagram





1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30





1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56



Supplementary appendix

Supplement to: Ho ISS, Azcoaga-Lorenzo A, Akbari A, et al. Variation in the estimated prevalence of multimorbidity: systematic review and meta-analysis of 193 studies.

For peer review only

Table of Contents

Table S1: Search strategy.....3

Table S2: Summary of the characteristics of outlying studies.....4

Table S3: Characteristics of 24 outlying studies.....5

Table S4: Characteristics of 193 included studies9

Table S5: Associations between predictors25

Table S6: Risk of bias assessment of included studies26

Table S7: Output of adjusted meta-analytic model based on 217 studies39

Table S8: Definition of variables40

Figure S1: Graphical display of study effect sizes and heterogeneity41

Figure S2: Process of examining and identifying outlying studies in meta-analysis42

Figure S3: Summary of risk of bias assessment43

Figure S4: Meta-regression trees for predicting the pooled estimated prevalence of multimorbidity (based on ‘mean age’ and ‘number of conditions’ predictors. unit: log(odds)).....44

References.....45

Table S1: Search strategy

| Database | Search strategy |
|---|--|
| Ovid Interface PsycINFO Embase Global Health Ovid MEDLINE | <ol style="list-style-type: none"> 1. (multimorbidity\$ or multi-morbidity\$ or comorbidity\$ or co-morbidity\$ or polymorbidity\$ or poly-morbidity\$ or multicondition\$ or multicondition\$ or "multiple chronic condition\$" or "morbidity burden" or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) adj2 (disease\$ or illness\$ or condition\$ or diagnos\$ or morbid\$))).m_titl. 2. (measure\$ or index or indices or instrument\$ or scale\$ or "disease count\$").mp. 3. 1 and 2 4. Limit 3 to human |
| EBSCO Interface CINAHL Plus | <ol style="list-style-type: none"> 1. MM (multimorbidity* or multi-morbidity* or comorbidity* or co-morbidity* or polymorbidity* or poly-morbidity* or multicondition* or multicondition* or "multiple chronic condition*" or "morbidity burden" or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) N2 (disease* or illness* or condition* or diagnos* or morbid*))) 2. AB (measure* or index or indices or instrument* or scale*) 3. 1 AND 2 <p>Limiters – Full Text; Human; Language: English</p> |
| Scopus | TITLE (multimorbidity* or multi-morbidity* or comorbidity* or co-morbidity* or polymorbidity* or poly-morbidity* or multicondition* or multicondition* or "multiple chronic condition*" or "morbidity burden" or ((multiple or coexisting or co-existing or concurrent or con-current or morbid or co-morbid) W/2 (disease* or illness* or condition* or diagnos?s or morbid*))) AND TITLE (measure* or index or indices or instrument* or scale* or "disease counts") |
| Web of Science | (TI=(measure* or index or indices or instrument* or scale*))AND (TI=(multimorbidity* or multi-morbidity* or comorbidity* or co-morbidity* or polymorbidity* or poly-morbidity* or multicondition* or multicondition* or 'multiple chronic condition*' or 'morbidity burden' or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) NEAR/2 (disease* or illness* or condition* or diagnos* or morbid*)))) AND LANGUAGE: (English) |
| Cochrane library | (multimorbidity or multi-morbidity or comorbidity or co-morbidity or polymorbidity or poly-morbidity or multicondition or multicondition or 'multiple chronic conditions' or 'morbidity burden' or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) NEAR/2 (disease or illness or condition or diagnosis or morbid))) AND (measure or index or indices or instrument or scale or "disease count*"):ti |
| ProQuest Dissertations & Theses Global | ti((multimorbidity* OR multi-morbidity* OR comorbidity* OR co-morbidity* OR polymorbidity* OR poly-morbidity* OR multicondition* OR multicondition* OR 'multiple chronic condition*' OR 'morbidity burden' OR ((multiple OR coexisting OR co-existing OR concurrent OR con-current OR morbid OR co-morbid) NEAR/2 (disease* OR illness* OR condition* OR diagnos?s OR morbid*)))) AND noft((measure* OR index OR indices OR instrument* OR scale*)) Limited by: Manuscript type: Doctoral dissertations, Master's theses Language: English |

Table S2: Summary of the characteristics of outlying studies

| Name of variable | Outlying studies (n=24) | All studies (n=217) |
|--|---|--|
| Prevalence of multimorbidity (%) | Range: 7.3 to 89.1 Pooled prevalence with the REML estimator: 31.0 (21.6-42.2) | Range: 2.7-95.6 Pooled prevalence with the REML estimator: 41.1 (37.7-44.6) |
| Mean age of study population (year) | Range of mean age: 39.6 to 82.2 Median of mean age: 56.6 (Q1, Q3: 52.3, 66.4) | Range of mean age: 32.2 to 83.8 Median of mean age: 62.4 (Q1, Q3: 50.2, 72.0) |
| No of conditions (count) | Range: 7 to 259 Median: 34 (Q1, Q3: 19.5, 54.5) | Range: 3 to 259 Median: 14.0 (Q1, Q3: 9, 21) |
| Country income (count, %) | | |
| High income | 21 (87.5%) | 166 (76.5%) |
| Low- or Middle-income | 3 (11.5%) | 51 (23.5%) |
| Continent (count, %) | | |
| Europe | 6 (25.0%) | 70 (32.3%) |
| North America | 7 (29.2%) | 54 (24.9%) |
| Asia | 7 (29.2%) | 51 (23.5%) |
| Australasia | 3 (12.5%) | 14 (6.5%) |
| Multiple continents | 1 (4.2%) | 12 (5.5%) |
| South America | | 12 (5.5%) |
| Africa | | 4 (1.8%) |
| Study population (count, %) | | |
| Only older people | 2 (8.3%) | 65 (30.0%) |
| Middle-aged and older | 1 (4.2%) | 47 (21.7%) |
| All adults | 15 (62.5%) | 99 (45.6%) |
| Only children | 1 (4.2%) | 1 (0.5%) |
| All age population | 5 (20.8%) | 5 (2.3%) |
| Setting (count, %) | | |
| Community | 12 (50.0%) | 159 (73.3%) |
| Primary care | 7 (29.2%) | 39 (18.0%) |
| Hospital | 4 (16.7%) | 18 (8.3%) |
| Care home | 1 (4.2%) | 1 (0.5%) |
| Source (count, %) | | |
| Self-report | 8 (33.3%) | 158 (72.8%) |
| Database | 16 (66.6%) | 59 (27.2%) |
| Risk of bias assessment (count, %) | | |
| Low | 4 (16.7%) | 13 (6.0%) |
| Moderate | 19 (79.2%) | 181 (83.4%) |
| High | 1 (4.2%) | 23 (10.6%) |

IQR: Interquartile range. SD: Standard deviation. The percentages were rounded so they do not add up to 100%.

Table S3: Characteristics of 24 outlying studies

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias | Rationale for exclusion |
|-----------------------------------|-------------|---------------|----------------|-----------|---------------------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|---|
| ¹ Stanley et al (2018) | New Zealand | Australasia | High | Hospitals | All adults | Not reported | 3489747 | Medical records and administrative database | 61 | 275706 | 0.08 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) and the studentized residual of this study is more than 2 standard deviations away from its expected value. |
| ² Lenzi et al (2016) | Italy | Europe | High | Hospitals | All adults | 66.4 | 3759836 | Medical records and administrative database | 26 | 574208 | 0.15 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ³ Hu et al (2019) | Taiwan | Asia | High | Community | All adults | Not reported | 1429527 | Medical records and administrative database | 20 | 939485 | 0.66 | Low | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ⁴ Gawron et al (2020) | USA | North America | High | Hospitals | All adults but not older people | Not reported | 741612 | Medical records and administrative database | Not reported | 53824 | 0.07 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) and the studentized residual of this study is more than 2 standard deviations away from its expected value. |
| ⁵ Low et al (2019) | Singapore | Asia | High | Community | All adults | 39.6 | 1181024 | Self-report | 48 | 309428 | 0.26 | Low | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ⁶ Wang et al (2014) | China | Asia | Low or middle | Community | Whole population | Not reported | 162464 | Self-report | 40 | 17987 | 0.11 | Low | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ⁷ Gaulin et al (2019) | Canada | North America | High | Hospitals | All adults | 51.2 | 1316832 | Medical records and administrative database | 34 | 416282 | 0.32 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias | Rationale for exclusion |
|-------------------------------------|----------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|--|
| ⁸ Violan et al (2014) | Spain | Europe | High | Primary care | All adults | 47.4 | 1356761 | Medical records and administrative database | 146 | 645818 | 0.48 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ⁹ Nicholson et al (2019) | Canada | North America | High | Primary care | All adults | 52.3 | 367743 | Medical records and administrative database | 20 | 195838 | 0.53 | High | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ¹⁰ Bao et al (2019) | China | Asia | Low or middle | Community | Middle aged and older | 61.36 | 18137 | Self-report | 19 | 3773 | 0.21 | Moderate | Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) |
| ¹¹ Fortin et al (2005) | Canada | North America | High | Primary care | All adults | 56.55 | 980 | Medical records and administrative database | 14 | 873 | 0.89 | Moderate | The studentized residual of this study is more than 2 standard deviations away from its expected value. |
| ¹² Prazeres et al (2015) | Portugal | Europe | High | Primary care | All adults | 56.3 | 1993 | Medical records and administrative database | 147 | 1449 | 0.73 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |
| ¹³ Lawson et al (2013) | UK | Europe | High | Community | All adults | 72.7 | 7054 | Medical records and administrative database | 40 | 1243 | 0.18 | Moderate | Irregular patterns found in compositional data (in scatter plot and Mahalanobis distance test)- low prevalence in studies with high mean participant age and a larger number of conditions |
| ¹⁴ Sullivan et al (2012) | USA | North America | High | Community | All adults | Not reported | 47178 | Medical records and administrative database | 259 | 19666 | 0.42 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias | Rationale for exclusion |
|--------------------------------------|-------------|---------------|----------------|--------------|-------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|---|
| ¹⁵ Peng et al (2020) | China | Asia | Low or middle | Community | Only older people | 71.6 | 1321 | Self-report | 15 | 589 | 0.45 | Moderate | Contributing to high levels of heterogeneity of effect sizes (in leave-one-out analysis) |
| ¹⁶ Excoffier et al (2018) | Switzerland | Europe | High | Primary care | All adults | 56.5 | 2904 | Medical records and administrative database | 75 | 1513 | 0.52 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |
| ¹⁷ Chung et al (2015) | Hong Kong | Asia | High | Community | All adults | Not reported | 25780 | Self-report | 46 | 3227 | 0.13 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |
| ¹⁸ Ki et al (2017) | South Korea | Asia | High | Community | All adults | 57.05 | 19942 | Medical records and administrative database | 66 | 5979 | 0.30 | Moderate | Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection) |
| ¹⁹ Bobo et al (2016) | USA | North America | High | Community | Whole population | Not reported | 138858 | Self-report | 19 | 33682 | 0.24 | Moderate | Infrequent values in compositional categorical data (few studies focused on whole population) |
| ²⁰ Randall et al (2018) | Australia | Australasia | High | Community | Whole population | Not reported | 5437018 | Self-report | 30 | 660449 | 0.12 | Moderate | Infrequent values in compositional categorical data (few studies focused on whole population) |
| ²¹ Russell et al (2020) | New Zealand | Australasia | High | Community | Only children | Not reported | 3838 | Self-report | 7 | 374 | 0.10 | Moderate | Infrequent values in compositional categorical data (only one study focused on children population) |
| ²² Barnett et al (2012) | UK | Europe | High | Primary care | Whole population | Not reported | 1751841 | Medical records and administrative database | 40 | 406427 | 0.23 | Low | Infrequent values in compositional categorical data (few studies focused on whole population) |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias | Rationale for exclusion |
|--------------------------------------|---|---------------------|----------------|--------------|-------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|---|
| ²³ St Sauver et al (2015) | USA | North America | High | Primary care | Whole population | Not reported | 106061 | Medical records and administrative database | 20 | 34592 | 0.33 | Moderate | Infrequent values in compositional categorical data (few studies focused on whole population) |
| ²⁴ Vetrano et al (2016) | Denmark, Finland, Iceland, Italy, the Netherlands, Norway, United Kingdom, Czech Republic, France, Sweden and Germany, Canada | Multiple continents | High | Care homes | Only older people | 82.2 | 6903 | Medical records and administrative database | 13 | 5098 | 0.74 | Moderate | Infrequent values in compositional categorical data (only one study focused on care home) |

MM: Multimorbidity. No of participants: The total number of participants in the denominator for estimating prevalence in a study (which could be a subset in some included studies)

Table S4: Characteristics of 193 included studies

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-----------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ²⁵ Aarts et al (2012) | The Netherlands | Europe | High | Primary care | All adults | 55.4 | 1184 | Medical records and administrative database | 23 | 420 | 0.35 | Moderate |
| ²⁶ Aarts et al (2011a) | The Netherlands | Europe | High | Community | Middle aged and older | 70 | 15188 | Self-report | Not reported | 7729 | 0.51 | Moderate |
| ²⁷ Aarts et al (2011b) | The Netherlands | Europe | High | Primary care | All adults | 55.4 | 1763 | Medical records and administrative database | 23 | 985 | 0.56 | Moderate |
| ²⁸ Abizanda et al (2014) | Spain | Europe | High | Primary care | Only older people | 78.6 | 842 | Medical records and administrative database | 14 | 580 | 0.69 | Moderate |
| ²⁹ Agborsangaya et al (2012) | Canada | North America | High | Community | All adults | 46.6 | 4003 | Self-report | 16 | 919 | 0.23 | Moderate |
| ³⁰ Agborsangaya et al (2013) | Canada | North America | High | Community | All adults | 47.8 | 4803 | Self-report | 16 | 1729 | 0.36 | Moderate |
| ³¹ Agborsangaya et al (2014) | Canada | North America | High | Community | All adults | 47.7 | 4752 | Self-report | 16 | 1597 | 0.34 | Moderate |
| ³² Ahrenfeldt et al (2019) | Europe | Europe | High | Community | Middle aged and older | 66.25 | 244258 | Self-report | 10 | 90652 | 0.37 | Moderate |
| ³³ Alimohammadian et al (2017) | Iran | Asia | Low or middle | Community | Middle aged and older | Not reported | 49946 | Self-report | 8 | 10035 | 0.20 | Moderate |
| ³⁴ Angst et al (2002) | Switzerland | Europe | High | Primary care | All adults | Not reported | 591 | Medical records and administrative database | 10 | 201 | 0.34 | High |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|---|---------------------|----------------|--------------|-----------------------|---------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ³⁵ Appa et al (2014) | USA | North America | High | Community | All adults | 60.2 | 1997 | Self-report | 16 | 1417 | 0.71 | Moderate |
| ³⁶ Adams et al (2017) | USA | North America | High | Community | All adults | Not reported | 400000 | Self-report | 12 | 191600 | 0.48 | Moderate |
| ³⁷ Ahmadi et al (2016) | Iran | Asia | Low or middle | Community | Middle aged and older | 52.1 | 49946 | Self-report | 8 | 10035 | 0.20 | Moderate |
| ³⁸ Amaral et al (2018) | Brazil | South America | Low or middle | Community | Only older people | Not reported | 264 | Self-report | 8 | 175 | 0.66 | Moderate |
| ³⁹ An et al (2016) | South Korea | Asia | High | Community | Middle aged and older | 54.8 | 10118 | Self-report | 8 | 3228 | 0.32 | Moderate |
| ⁴⁰ Araujo et al (2018) | Brazil | South America | Low or middle | Community | All adults | Not reported | 4001 | Self-report | 12 | 1160 | 0.29 | Moderate |
| ⁴¹ Arnold-Reed et al (2018) | Australia | Australasia | High | Primary care | All adults | 38.2 | 4285 | Medical records and administrative database | 43 | 2269 | 0.53 | Moderate |
| ⁴² Arokiasamy et al (2015) | 6 low middle income countries (China, Ghana, India, Mexico, Russia, South Africa) | Multiple continents | Low or middle | Community | All adults | Not reported | 42236 | Self-report | 8 | 9250 | 0.22 | Moderate |
| ⁴³ Sinnige et al (2015) | The Netherlands | Europe | High | Primary care | Middle aged and older | 66.9 | 120480 | Medical records and administrative database | 29 | 74733 | 0.62 | Moderate |
| ⁴⁴ Zemedikun et al (2018) | UK | Europe | High | Community | Middle aged and older | Median age 58 | 502643 | Medical records and administrative database | 36 | 95710 | 0.19 | Moderate |
| ⁴⁵ Wensing et al (2001) | The Netherlands | Europe | High | Primary care | All adults | Not reported | 3867 | Self-report | 25 | 626 | 0.16 | Moderate |
| ⁴⁶ Mounce et al (2018) | UK | Europe | High | Community | Middle aged and older | Not reported | 4564 | Self-report | 15 | 1553 | 0.34 | Moderate |
| ⁴⁷ Taylor et al (2010) | Australia | Australasia | High | Community | All adults | Not reported | 3206 | Self-report | 7 | 547 | 0.17 | Low |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|---|---------------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁴⁸ Vancampfort et al (2019) | Six low and middle income countries (China, Ghana, India, Mexico, Russia, and South Africa) | Multiple continents | Low or middle | Community | Middle aged and older | 62.4 | 34129 | Self-report | 11 | 15529 | 0.46 | Moderate |
| ⁴⁹ Vancampfort et al (2018) | Six low and middle income countries (China, Ghana, India, Mexico, Russia, and South Africa) | Multiple continents | Low or middle | Community | Only older people | 72.6 | 14585 | Self-report | 11 | 8780 | 0.60 | Moderate |
| ⁵⁰ Aubert et al (2016) | Switzerland | Europe | High | Primary care | Middle aged and older | 63.5 | 1002 | Medical records and administrative database | 17 | 676 | 0.67 | Moderate |
| ⁵¹ Autenrieth et al (2013) | Germany | Europe | High | Community | Only older people | 75.7 | 1007 | Self-report | 13 | 658 | 0.65 | Moderate |
| ⁵² Bahler et al (2015) | Switzerland | Europe | High | Community | Only older people | 74.9 | 229493 | Medical records and administrative database | 22 | 175752 | 0.77 | Moderate |
| ⁵³ Vancampfort et al (2017) | 44 low and middle income countries | Multiple continents | Low or middle | Community | All adults | 38.3 | 194431 | Self-report | 11 | 27518 | 0.14 | Moderate |
| ⁵⁴ Banjare et al (2014) | India | Asia | Low or middle | Community | Only older people | Not reported | 310 | Self-report | 20 | 176 | 0.57 | Moderate |
| ⁵⁵ Barra et al (2015) | USA | North America | High | Community | All adults | 45.36 | 43079 | Self-report | Not reported | 22412 | 0.52 | Moderate |
| ⁵⁶ Bernard et al (2016) | Australia | Australasia | High | Hospitals | Only older people | 81.8 | 306 | Medical records and administrative database | 19 | 125 | 0.41 | High |
| ⁵⁷ Biswas et al (2019) | Bangladesh | Asia | Low or middle | Community | All adults | Not reported | 8763 | Self-report | 3 | 1078 | 0.12 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|---|---------------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁵⁸ Blakemore et al (2016) | UK | Europe | High | Primary care | Only older people | 75 | 4377 | Self-report | 24 | 2631 | 0.60 | Moderate |
| ⁵⁹ Blyth et al (2008) | Australia | Australasia | High | Community | Only older people | 76.9 | 1685 | Self-report | 18 | 920 | 0.55 | Moderate |
| ⁶⁰ Bowling et al (2019) | USA | North America | High | Community | Middle aged and older | 56.7 | 4217 | Self-report | 12 | 3053 | 0.72 | Moderate |
| ⁶¹ Britt et al (2008) | Australia | Australasia | High | Primary care | All adults | Not reported | 9156 | Medical records and administrative database | 18 | 3398 | 0.37 | Moderate |
| ⁶² Broeiro-Goncalves et al (2019) | Portugal | Europe | High | Hospitals | All adults | 59.8 | 800376 | Medical records and administrative database | 22 | 335357 | 0.42 | Moderate |
| ⁶³ Bruce et al (2010) | Canada | North America | High | Community | All adults | 37.8 | 453 | Self-report | 4 | 163 | 0.36 | High |
| ⁶⁴ Burgers et al (2010) | France, Germany, Canada, Australia, Netherlands, New Zealand, UK, USA | Multiple continents | High | Community | All adults | Not reported | 8973 | Self-report | 7 | 4037 | 0.45 | Moderate |
| ⁶⁵ Burke et al (2017) | US, Europe, Asia | Multiple continents | High | Community | Only older people | Not reported | 4668 | Self-report | 9 | 2165 | 0.46 | Moderate |
| ⁶⁶ Buurman et al (2016) | The Netherlands | Europe | High | Hospitals | Only older people | 78.2 | 639 | Medical records and administrative database | 35 | 440 | 0.69 | Moderate |
| ⁶⁷ Calderon-Larranaga et al (2017) | Sweden | Europe | High | Primary care | Only older people | 74.6 | 3363 | Self-report | 60 | 2980 | 0.89 | Moderate |
| ⁶⁸ Camargo-Casas et al (2018) | Colombia | South America | Low or middle | Community | Only older people | 71.1 | 2000 | Self-report | 12 | 808 | 0.40 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|---|---------------------|----------------|--------------|-----------------------|----------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁶⁹ Canevelli et al (2019) | Italy | Europe | High | Primary care | Only older people | 75.1 | 185 | Medical records and administrative database | 18 | 162 | 0.88 | High |
| ⁷⁰ Chamberlain et al (2020) | USA | North America | High | Community | All adults | Not reported | 198941 | Self-report | 21 | 78527 | 0.39 | Low |
| ⁷¹ Chen et al (2018) | China | Asia | Low or middle | Community | Only older people | Not reported | 30774 | Medical records and administrative database | 33 | 25101 | 0.82 | Low |
| ⁷² Chen et al (2018) | China | Asia | Low or middle | Community | Middle aged and older | Not reported | 3737 | Self-report | 16 | 1722 | 0.46 | Moderate |
| ⁷³ Cheung et al (2013) | Hong Kong (SAR of China) | Asia | High | Community | Middle aged and older | 71.3 | 1145 | Self-report | 18 | 654 | 0.57 | Moderate |
| ⁷⁴ Chu et al (2018) | Hong Kong (SAR of China) | Asia | High | Primary care | Middle aged and older | Not reported | 382 | Medical records and administrative database | 40 | 206 | 0.54 | Moderate |
| ⁷⁵ Chudasama et al (2019) | UK | Europe | High | Community | Middle aged and older | Median age:58 | 491939 | Medical records and administrative database | 36 | 96622 | 0.20 | Moderate |
| ⁷⁶ Cimarras-Otal et al (2014) | Spain | Europe | High | Community | All adults | Not reported | 22190 | Self-report | 20 | 7830 | 0.35 | Moderate |
| ⁷⁷ Chin et al (2016) | Hong Kong (SAR of China) | Asia | High | Primary care | All adults | Median age: 48 | 9259 | Self-report | 8 | 2350 | 0.25 | Moderate |
| ⁷⁸ Agrawal et al (2016) | India, China, Russia, Mexico, South Africa, Ghana | Multiple continents | Low or middle | Community | All adults | 57.8 | 40166 | Self-report | 9 | 9238 | 0.23 | Moderate |
| ⁷⁹ Gu et al (2018) | China | Asia | Low or middle | Community | Only older people | Not reported | 411 | Self-report | 13 | 232 | 0.56 | Moderate |
| ⁸⁰ Gunn et al (2012) | Australia | Australasia | High | Primary care | All adults | 50.89 | 6864 | Self-report | 12 | 2154 | 0.31 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-----------|---------------|----------------|--------------|-----------------------|----------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁸¹ Han et al (2013) | USA | North America | High | Primary care | Only older people | 76 | 159 | Medical records and administrative database | 18 | 117 | 0.74 | High |
| ⁸² Hanlon et al (2018) | UK | Europe | High | Community | All adults | Not reported | 493737 | Medical records and administrative database | 42 | 161576 | 0.33 | Low |
| ⁸³ Jantsch et al (2018) | Brazil | South America | Low or middle | Community | All adults | 42 | 3092 | Self-report | 11 | 912 | 0.29 | Moderate |
| ⁸⁴ John et al (2003) | USA | North America | High | Community | Only older people | 71.3 | 992 | Self-report | 11 | 732 | 0.74 | High |
| ⁸⁵ Johnson-Lawrence et al (2017) | USA | North America | High | Community | All adults | 49.9 | 115097 | Self-report | 9 | 27278 | 0.24 | Moderate |
| ⁸⁶ Johnston et al (2019) | UK | Europe | High | Community | All adults | 48 | 7184 | Self-report | Not reported | 388 | 0.05 | Moderate |
| ⁸⁷ Jones et al (2016) | USA | North America | High | Community | Only older people | Not reported | 6964 | Self-report | 10 | 4951 | 0.71 | Moderate |
| ⁸⁸ Jovic et al (2016) | Serbia | Europe | Low or middle | Community | All adults | 49.4 | 13103 | Self-report | 13 | 3522 | 0.27 | Moderate |
| ⁸⁹ Juul-Larsen et al (2020) | Denmark | Europe | High | Hospitals | Only older people | Median age: 78 | 369 | Self-report | 34 | 311 | 0.84 | Moderate |
| ⁹⁰ Hudon et al (2008) | Canada | North America | High | Community | All adults | Not reported | 16782 | Self-report | 25 | 5343 | 0.32 | Low |
| ⁹¹ Hussain et al (2015) | Indonesia | Asia | Low or middle | Community | Middle aged and older | Not reported | 9438 | Self-report | 12 | 3369 | 0.36 | Moderate |
| ⁹² Ie et al (2017) | USA | North America | High | Hospitals | Only older people | Not reported | 1084 | Medical records and administrative database | 24 | 1036 | 0.96 | High |
| ⁹³ Ishizaki et al (2019) | Japan | Asia | High | Community | Only older people | 76.9 | 2525 | Self-report | 9 | 1121 | 0.44 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ⁹⁴ Danon-Hersch et al (2012) | Switzerland | Europe | High | Community | Only older people | Not reported | 1283 | Self-report | 12 | 448 | 0.35 | Moderate |
| ⁹⁵ de Heer et al (2013) | USA | North America | High | Community | All adults | 47.72 | 1002 | Self-report | 19 | 378 | 0.38 | Moderate |
| ⁹⁶ Demirchyan et al (2013) | Armenia | Asia | Low or middle | Community | All adults | 58.8 | 721 | Self-report | Not reported | 564 | 0.78 | High |
| ⁹⁷ Fabbri et al (2015) | Italy | Europe | High | Community | Only older people | 73.6 | 1018 | Self-report | 15 | 458 | 0.45 | Moderate |
| ⁹⁸ Fillenbaum et al (2000) | USA | North America | High | Community | Only older people | 73.44 | 4034 | Self-report | 5 | 1181 | 0.29 | Moderate |
| ⁹⁹ Kaneko et al (2019) | Japan | Asia | High | Community | Only older people | Not reported | 253 | Self-report | Not reported | 135 | 0.53 | Moderate |
| ¹⁰⁰ Kang et al (2017) | South Korea | Asia | High | Primary care | All adults | 32.2 | 590 | Medical records and administrative database | 14 | 153 | 0.26 | Moderate |
| ¹⁰¹ Gandhi et al (2020) | USA | North America | High | Community | All adults | Not reported | 9499 | Self-report | 8 | 3379 | 0.36 | Moderate |
| ¹⁰² Costa et al (2018) | Brazil | South America | Low or middle | Community | Only older people | Not reported | 1451 | Self-report | 29 | 1343 | 0.93 | Moderate |
| ¹⁰³ Rizzuto et al (2017) | Sweden | Europe | High | Community | Only older people | Not reported | 1099 | Self-report | 36 | 774 | 0.70 | Moderate |
| ¹⁰⁴ Dhalwani et al (2017) | UK | Europe | High | Community | Middle aged and older | Not reported | 5476 | Self-report | 18 | 1156 | 0.21 | Moderate |
| ¹⁰⁵ Elixhauser et al (1998) | USA | North America | High | Hospitals | All adults | 57.1 | 1779167 | Medical records and administrative database | 30 | 619150 | 0.35 | Low |
| ¹⁰⁶ Fabbri et al (2015) | USA | North America | High | Hospitals | Only older people | 72.3 | 695 | Self-report | 15 | 440 | 0.63 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---------------------------------------|---|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁰⁷ Fortin et al (2014) | Canada | North America | High | Community | Middle aged and older | 57.8 | 1196 | Self-report | 14 | 599 | 0.50 | Moderate |
| ¹⁰⁸ Fuchs et al (1998) | Israel | Asia | High | Community | Only older people | Not reported | 1820 | Self-report | 14 | 1174 | 0.65 | Moderate |
| ¹⁰⁹ Galenkamp et al (2011) | The Netherlands | Europe | High | Community | Middle aged and older | 69.2 | 2046 | Self-report | 7 | 876 | 0.43 | High |
| ¹¹⁰ Galenkamp et al (2016) | Germany, UK, Italy, The Netherlands, Spain and Sweden | Europe | High | Community | Only older people | 74.2 | 2792 | Self-report | 8 | 1358 | 0.49 | Moderate |
| ¹¹¹ Gamma et al (2001) | Switzerland | Europe | High | Community | All adults | Not reported | 407 | Self-report | 14 | 53 | 0.13 | High |
| ¹¹² Ge et al (2018) | Singapore | Asia | High | Community | All adults | 51.4 | 1940 | Self-report | 17 | 715 | 0.37 | Moderate |
| ¹¹³ Ge et al (2019) | Singapore | Asia | High | Community | All adults | 51.3 | 1932 | Self-report | 17 | 564 | 0.29 | Moderate |
| ¹¹⁴ Gould et al (2016) | USA | North America | High | Community | Only older people | 74.82 | 4184 | Self-report | 7 | 2932 | 0.70 | Moderate |
| ¹¹⁵ Habib et al (2014) | Lebanon | Asia | Low or middle | Community | All adults | 46.6 | 2501 | Self-report | Not reported | 665 | 0.27 | Moderate |
| ¹¹⁶ Harrison et al (2017) | Australia | Australasia | High | Primary care | All adults | Not reported | 8707 | Medical records and administrative database | 28 | 2838 | 0.33 | Moderate |
| ¹¹⁷ Hayek et al (2017) | Israel | Asia | High | Community | All adults | 47.2 | 4325 | Self-report | 10 | 1579 | 0.37 | Moderate |
| ¹¹⁸ Henninger et al (2012) | USA | North America | High | Community | Only older people | 76 | 3212 | Self-report | 9 | 1753 | 0.55 | Moderate |
| ¹¹⁹ Hernandez et al (2019) | Ireland | Europe | High | Community | Middle aged and older | Not reported | 6101 | Self-report | 31 | 4468 | 0.73 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|---|---------------------|----------------|-----------|-----------------------|--------------|--------------------|-------------|---------------------------|----------------|--------------------|--------------|
| ¹²⁰ Ho et al (2014) | Singapore | Asia | High | Community | Middle aged and older | 66.15 | 1844 | Self-report | 12 | 830 | 0.45 | Moderate |
| ¹²¹ Khan et al (2019) | Bangladesh | Asia | Low or middle | Community | All adults | 58.6 | 12338 | Self-report | 6 | 1031 | 0.08 | Low |
| ¹²² Kiliari et al (2013) | Cyprus | Europe | High | Community | All adults | 53 | 465 | Self-report | Not reported | 132 | 0.28 | Moderate |
| ¹²³ King et al (2018) | USA | North America | High | Community | All adults | Not reported | 5541 | Self-report | 11 | 3342 | 0.60 | Moderate |
| ¹²⁴ Kingston et al (2018) | UK | Europe | High | Community | All adults | Not reported | 9723900 | Self-report | 12 | 5250906 | 0.54 | High |
| ¹²⁵ Koyanagi et al (2018) | China, Ghana, India, Mexico, Russia, and South Africa | Multiple continents | Low or middle | Community | Middle aged and older | 62.1 | 32715 | Self-report | 10 | 16324 | 0.50 | Moderate |
| ¹²⁶ Kriegsman et al (2004) | The Netherlands | Europe | High | Community | Middle aged and older | 69.2 | 2489 | Self-report | 7 | 519 | 0.21 | Moderate |
| ¹²⁷ Kristensen et al (2019) | Germany | Europe | High | Community | Middle aged and older | 63.47 | 19605 | Self-report | 13 | 12600 | 0.64 | Moderate |
| ¹²⁸ Kristensen et al (2019) | Germany | Europe | High | Community | Middle aged and older | 64.37 | 7604 | Self-report | 13 | 5140 | 0.68 | Moderate |
| ¹²⁹ Kunna et al (2017) | China, Ghana | Multiple continents | Low or middle | Community | Middle aged and older | Not reported | 15864 | Self-report | 7 | 4731 | 0.30 | Low |
| ¹³⁰ Kuwornu et al (2014) | Canada | North America | High | Community | All adults | 51.05 | 3284 | Self-report | 15 | 1143 | 0.35 | Moderate |
| ¹³¹ Lai et al (2019) | Hong Kong (SAR of China) | Asia | High | Community | All adults | Not reported | 69636 | Self-report | 14 | 3898 | 0.06 | Moderate |
| ¹³² Lai et al (2018) | Hong Kong (SAR of China) | Asia | High | Community | All adults | Not reported | 300 | Self-report | 11 | 48 | 0.16 | Moderate |
| ¹³³ Laires et al (2019) | Portugal | Europe | High | Community | All adults | Not reported | 15196 | Self-report | 13 | 6671 | 0.44 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-----------------------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹³⁴ Lang et al (2015) | USA | North America | High | Community | Middle aged and older | 53.4 | 3058 | Self-report | 6 | 948 | 0.31 | Moderate |
| ¹³⁵ Le Cossec et al (2016) | France | Europe | High | Community | Middle aged and older | 70 | 15325 | Self-report | 4 | 3528 | 0.23 | Moderate |
| ¹³⁶ Lee et al (2007) | USA | North America | High | Hospitals | Middle aged and older | Not reported | 741847 | Medical records and administrative database | 11 | 302792 | 0.41 | Low |
| ¹³⁷ Lee et al (2018) | Taiwan | Asia | High | Community | Only older people | Not reported | 20898 | Medical records and administrative database | Not reported | 4234 | 0.20 | High |
| ¹³⁸ Li et al (2016) | UK | Europe | High | Primary care | All adults | Not reported | 27806 | Self-report | 12 | 10332 | 0.37 | Moderate |
| ¹³⁹ Li et al (2019) | USA | North America | High | Community | Middle aged and older | 67.4 | 14996 | Self-report | 8 | 9805 | 0.65 | Moderate |
| ¹⁴⁰ Lujic et al (2017) | Australia | Australasia | High | Community | Middle aged and older | 70.2 | 90352 | Self-report | 8 | 33792 | 0.37 | Moderate |
| ¹⁴¹ Lupianez-Villanueva et al (2018) | 14 European countries | Europe | High | Community | All adults | Not reported | 14000 | Self-report | 13 | 3416 | 0.24 | Moderate |
| ¹⁴² Zhou et al (2018) | Bangladesh, India and China | Asia | Low or middle | Community | All adults | Not reported | 18696 | Self-report | 9 | 3512 | 0.19 | Moderate |
| ¹⁴³ Zhang et al (2019) | China | Asia | Low or middle | Community | Only older people | 70.5 | 11707 | Self-report | 11 | 5104 | 0.44 | Moderate |
| ¹⁴⁴ Wong et al (2010) | Canada | North America | High | Community | Only older people | Not reported | 740 | Self-report | 7 | 489 | 0.66 | Moderate |
| ¹⁴⁵ Weimann et al (2016) | South Africa | Africa | Low or middle | Community | All adults | 34 | 18526 | Self-report | 4 | 506 | 0.027 | Moderate |
| ¹⁴⁶ Wang et al (2017) | Australia | Australasia | High | Community | All adults | 44 | 8820 | Self-report | 8 | 2539 | 0.29 | Moderate |
| ¹⁴⁷ Wang et al (2019) | South Africa | Africa | Low or middle | Community | Only older people | Not reported | 2627 | Self-report | 5 | 439 | 0.17 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|-----------------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁴⁸ Wade et al (2019) | New Zealand | Australasia | High | Community | All adults | 59.05 | 7654 | Self-report | 12 | 2786 | 0.36 | Moderate |
| ¹⁴⁹ Maciejewski et al (2019) | USA | North America | High | Community | Only older people | 77.1 | 20124230 | Medical records and administrative database | 19 | 14425446 | 0.72 | Moderate |
| ¹⁵⁰ Marengoni et al (2016) | Sweden | Europe | High | Community | Only older people | 74.4 | 3155 | Medical records and administrative database | 14 | 1654 | 0.52 | Moderate |
| ¹⁵¹ Marengoni et al (2009) | Sweden | Europe | High | Community | Only older people | Not reported | 1099 | Self-report | 22 | 575 | 0.52 | Moderate |
| ¹⁵² Marques et al (2018) | 13 European countries | Europe | High | Community | All adults | 50.2 | 32931 | Self-report | 6 | 7113 | 0.22 | Moderate |
| ¹⁵³ Mavaddat et al (2014) | UK | Europe | High | Primary care | Middle aged and older | 58.7 | 11439 | Self-report | 6 | 1006 | 0.09 | Moderate |
| ¹⁵⁴ McDaid et al (2013) | Ireland | Europe | High | Community | Middle aged and older | Not reported | 6018 | Self-report | 8 | 733 | 0.12 | High |
| ¹⁵⁵ Melis et al (2014) | Sweden | Europe | High | Hospitals | Only older people | 83.75 | 390 | Medical records and administrative database | 39 | 213 | 0.55 | Moderate |
| ¹⁵⁶ Min et al (2007) | USA | North America | High | Community | Only older people | 81 | 372 | Self-report | 9 | 230 | 0.62 | High |
| ¹⁵⁷ Momtaz et al (2010) | Malaysia | Asia | High | Community | Only older people | 69.26 | 385 | Self-report | 16 | 165 | 0.43 | Moderate |
| ¹⁵⁸ Mondor et al (2018) | Canada | North America | High | Community | All adults | Not reported | 27195 | Medical records and administrative database | 17 | 11390 | 0.42 | Moderate |
| ¹⁵⁹ Muggah et al (2012) | Canada | North America | High | Community | All adults | Not reported | 28450000 | Medical records and administrative database | 9 | 4523550 | 0.16 | Moderate |
| ¹⁶⁰ Nagel et al (2008) | Germany | Europe | High | Community | Middle aged and older | 56.5 | 13781 | Self-report | 15 | 9275 | 0.67 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|--------------|---------------|----------------|--------------|-----------------------|----------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁶¹ Niedzwiedz et al (2019) | USA | North America | High | Community | Middle aged and older | 67.2 | 2272 | Self-report | 8 | 1491 | 0.66 | Moderate |
| ¹⁶² Nunes et al (2016) | Brazil | South America | Low or middle | Community | All adults | 45.75 | 2927 | Self-report | 11 | 852 | 0.29 | Moderate |
| ¹⁶³ Nunes et al (2017) | Brazil | South America | Low or middle | Community | All adults | 43.7 | 60202 | Self-report | 22 | 13365 | 0.22 | Moderate |
| ¹⁶⁴ Nunes et al (2015) | Brazil | South America | Low or middle | Community | Only older people | Not reported | 1593 | Self-report | 17 | 1295 | 0.81 | Moderate |
| ¹⁶⁵ Olaya et al (2017) | Spain | Europe | High | Community | Only older people | 71.75 | 2113 | Self-report | 7 | 1088 | 0.51 | Moderate |
| ¹⁶⁶ Olivares et al (2017) | Argentina | South America | High | Community | All adults | 43 | 1044 | Self-report | Not reported | 346 | 0.33 | Moderate |
| ¹⁶⁷ Park et al (2018) | South Korea | Asia | High | Community | Middle aged and older | 62.7 | 5996 | Self-report | 25 | 1607 | 0.27 | Moderate |
| ¹⁶⁸ Patel et al (2006) | Mexico | South America | Low or middle | Community | Middle aged and older | 73 | 7852 | Self-report | 5 | 1833 | 0.23 | Moderate |
| ¹⁶⁹ Pati et al (2016) | India | Asia | Low or middle | Community | All adults | 44.96 | 103 | Self-report | 18 | 24 | 0.23 | Moderate |
| ¹⁷⁰ Pati et al (2019) | India | Asia | Low or middle | Primary care | All adults | 44 | 1649 | Self-report | 21 | 567 | 0.34 | Moderate |
| ¹⁷¹ Payne et al (2013) | UK | Europe | High | Primary care | All adults | 49 | 180815 | Medical records and administrative database | 40 | 54945 | 0.30 | Moderate |
| ¹⁷² Perez et al (2020) | Sweden | Europe | High | Community | Only older people | 72.8 | 2596 | Self-report | 60 | 2213 | 0.85 | Moderate |
| ¹⁷³ Petersen et al (2019) | South Africa | Africa | Low or middle | Primary care | All adults | Not reported | 2549 | Self-report | Not reported | 893 | 0.35 | Moderate |
| ¹⁷⁴ Pfortmueller et al (2013) | Switzerland | Europe | High | Hospitals | All adults | Median age: 28 | 3170 | Medical records and administrative database | 18 | 1183 | 0.37 | High |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---------------------------------------|--|---------------|----------------|--------------|-----------------------|---------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁷⁵ Pressley et al (1999) | USA | North America | High | Hospitals | Only older people | Not reported | 5934 | Medical records and administrative database | Not reported | 3534 | 0.60 | Moderate |
| ¹⁷⁶ Prior et al (2016) | Denmark | Europe | High | Community | All adults | Not reported | 118410 | Self-report | 39 | 33937 | 0.29 | Moderate |
| ¹⁷⁷ Ribeiro et al (2018) | Brazil | South America | High | Community | Only older people | 70 | 820 | Self-report | 8 | 270 | 0.33 | Moderate |
| ¹⁷⁸ Ruel et al (2014) | Australia | Australasia | High | Community | All adults | 50 | 1854 | Self-report | 8 | 585 | 0.32 | Moderate |
| ¹⁷⁹ Ruel et al (2014) | China | Asia | Low or middle | Community | All adults | 49 | 1020 | Self-report | 11 | 346 | 0.34 | Moderate |
| ¹⁸⁰ Ryan et al (2018) | Ireland | Europe | High | Community | Middle aged and older | Not reported | 4823 | Self-report | 16 | 2588 | 0.54 | Moderate |
| ¹⁸¹ Schmidt et al (2016) | Austria, Belgium, Denmark, France, Germany, Italy, the Netherlands, Spain, Sweden, and Switzerland | Europe | High | Community | Only older people | Not reported | 56609 | Self-report | 11 | 13794 | 0.24 | Moderate |
| ¹⁸² Schottker et al (2016) | Germany | Europe | High | Primary care | Middle aged and older | Median age:70 | 2547 | Medical records and administrative database | 14 | 251 | 0.10 | Moderate |
| ¹⁸³ Seo et al (2017) | South Korea | Asia | High | Community | Middle aged and older | Not reported | 156747 | Self-report | 15 | 42006 | 0.27 | Moderate |
| ¹⁸⁴ She et al (2019) | China | Asia | Low or middle | Hospitals | Only older people | 68.9 | 1497 | Self-report | 22 | 1255 | 0.84 | Moderate |
| ¹⁸⁵ Singh et al (2019) | India | Asia | Low or middle | Community | All adults | 41 | 16287 | Self-report | 5 | 1531 | 0.09 | Moderate |
| ¹⁸⁶ Stepanova et al (2015) | USA | North America | High | Community | All adults | 34.7 | 26225 | Self-report | 13 | 9992 | 0.38 | High |
| ¹⁸⁷ Stickley et al (2020) | USA | North America | High | Community | All adults | 44.9 | 15311 | Self-report | 9 | 3996 | 0.26 | High |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|---|---|---------------------|----------------|--------------|-----------------------|----------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ¹⁸⁸ Streit et al (2014) | Switzerland | Europe | High | Primary care | Middle aged and older | 63.5 | 1002 | Medical records and administrative database | 17 | 676 | 0.67 | Moderate |
| ¹⁸⁹ Stubbs et al (2018) | China, Ghana, India, Mexico, Russia, South Africa | Multiple continents | Low or middle | Community | Middle aged and older | 62.4 | 34129 | Self-report | 13 | 19317 | 0.57 | Moderate |
| ¹⁹⁰ Su et al (2016) | China | Asia | Low or middle | Community | Only older people | Not reported | 2058 | Self-report | 10 | 1012 | 0.49 | Moderate |
| ¹⁹¹ Sundstrup et al (2017) | USA | North America | High | Community | All adults | 43.5 | 10427 | Self-report | 8 | 2489 | 0.24 | High |
| ¹⁹² Takahashi et al (2016) | USA | North America | High | Hospitals | All adults | 57 | 6402 | Medical records and administrative database | Not reported | 3140 | 0.49 | High |
| ¹⁹³ Tinetti et al (2011) | USA | North America | High | Community | Only older people | 72.6 | 5298 | Self-report | 5 | 1200 | 0.23 | High |
| ¹⁹⁴ Troelstra et al (2020) | The Netherlands | Europe | High | Community | All adults | Not reported | 604 | Self-report | 26 | 321 | 0.53 | High |
| ¹⁹⁵ van Zon et al (2020) | USA | North America | High | Community | Middle aged and older | 53.8 | 10719 | Self-report | 8 | 2390 | 0.22 | Moderate |
| ¹⁹⁶ Vancampfort et al (2017) | China, Ghana, India, Mexico, Russia, and South Africa | Multiple continents | Low or middle | Community | All adults | Median age: 62 | 32585 | Self-report | 11 | 14524 | 0.45 | Moderate |
| ¹⁹⁷ Vassilaki et al (2015) | USA | North America | High | Primary care | Only older people | 78.5 | 2176 | Medical records and administrative database | 17 | 1884 | 0.87 | Moderate |
| ¹⁹⁸ Vassilaki et al (2016) | USA | North America | High | Primary care | Only older people | 79 | 1449 | Medical records and administrative database | 17 | 1237 | 0.85 | Moderate |
| ¹⁹⁹ Villarreal et al (2015) | Panama | South America | High | Primary care | Only older people | 78.2 | 304 | Self-report | 7 | 227 | 0.75 | Moderate |
| ²⁰⁰ Violan et al (2019) | Spain | Europe | High | Primary care | Only older people | 75.4 | 916619 | Medical records and | 60 | 853085 | 0.93 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|--|-----------------|---------------|----------------|--------------|-----------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| | | | | | | | | administrative database | | | | |
| ²⁰¹ von Strauss et al (2000) | Sweden | Europe | High | Community | Only older people | Not reported | 502 | Self-report | 15 | 155 | 0.31 | Moderate |
| ²⁰² Vos et al (2013) | The Netherlands | Europe | High | Community | Only older people | 71.9 | 315 | Self-report | 21 | 202 | 0.64 | Moderate |
| ²⁰³ Vu et al (2019) | Vietnam | Asia | Low or middle | Hospitals | Only older people | 71.9 | 405 | Medical records and administrative database | Not reported | 146 | 0.36 | High |
| ²⁰⁴ Wang et al (2018) | USA | North America | High | Community | All adults | 47 | 3086 | Self-report | 20 | 1109 | 0.36 | Moderate |
| ²⁰⁵ Wang et al (2017) | China | Asia | Low or middle | Community | Only older people | 69.24 | 2705 | Self-report | 17 | 1230 | 0.45 | Moderate |
| ²⁰⁶ Wijers et al (2019) | Spain | Europe | High | Community | Middle aged and older | 74.2 | 707 | Self-report | 21 | 491 | 0.69 | Moderate |
| ²⁰⁷ Williams et al (2016) | USA | North America | High | Community | All adults | Not reported | 23789 | Self-report | 9 | 9213 | 0.39 | Moderate |
| ²⁰⁸ Woldeamayrat et al (2018) | Ethiopia | Africa | Low or middle | Primary care | All adults | Not reported | 411 | Self-report | 18 | 73 | 0.18 | Moderate |
| ²⁰⁹ Yao et al (2020) | China | Asia | Low or middle | Community | Middle aged and older | 57.7 | 10084 | Self-report | 15 | 3243 | 0.32 | Moderate |
| ²¹⁰ Yorke et al (2017) | USA | North America | High | Community | Middle aged and older | 66.6 | 5877 | Self-report | 7 | 3391 | 0.58 | Moderate |
| ²¹¹ You et al (2019) | China | Asia | Low or middle | Community | Only older people | 72 | 5296 | Self-report | 27 | 2201 | 0.42 | Moderate |
| ²¹² Zhang et al (2020) | China | Asia | Low or middle | Community | Only older people | 74.14 | 4348 | Self-report | 15 | 2338 | 0.54 | Moderate |
| ²¹³ Khanam et al (2011) | Bangladesh | Asia | Low or middle | Community | Only older people | 69.5 | 452 | Medical records and administrative database | 9 | 243 | 0.54 | Moderate |

| Author | Country | Continent | Country income | Setting | Study population | Mean age | No of participants | Source | No of conditions measured | No of MM cases | Proportion with MM | Risk of bias |
|-------------------------------------|--------------------------|---------------|----------------|--------------|------------------|--------------|--------------------|---|---------------------------|----------------|--------------------|--------------|
| ²¹⁴ Cornell et al (2009) | USA | North America | High | Primary care | All adults | 62.4 | 1645314 | Medical records and administrative database | 45 | 1327382 | 0.81 | Moderate |
| ²¹⁵ Cassell et al (2018) | UK | Europe | High | Primary care | All adults | Not reported | 403985 | Medical records and administrative database | 36 | 109884 | 0.27 | Moderate |
| ²¹⁶ Wong et al (2019) | Hong Kong (SAR of China) | Asia | High | Community | All adults | 45.67 | 1014 | Self-report | 5 | 124 | 0.12 | Moderate |
| ²¹⁷ Puth et al (2017) | Germany | Europe | High | Community | All adults | Not reported | 19294 | Self-report | 17 | 7640 | 0.40 | Moderate |

MM: Multimorbidity. No of participants is the total number of participants in the denominator for estimating prevalence in a study (which could be a subset in some included studies)

Table S5: Associations between predictors

| | Mean age (lm) Unadjusted coefficient estimates | No of conditions (nb) Unadjusted incident rate ratio |
|-------------------------|--|--|
| Mean age | | 1.0 (1.0-1.0) |
| Source | | |
| Self-report | 59.7 (57.1-62.3) (intercept) | Ref |
| Database | 7.0 (1.5-12.5)* | 1.8 (1.5-2.2)*** |
| Continent | | |
| Europe | 66.8 (62.8-70.9) (intercept) | Ref |
| North America | -7.0 (-12.8 to -1.1)* | 0.6 (0.5-0.8)*** |
| Australasia | -8.0 (-17.5-1.6) | 0.8 (0.6-1.1) |
| Asia | -8.4 (-14.6 to -2.2)** | 0.6 (0.5-0.8)*** |
| South America | -8.5 (-18.0-1.1) | 0.6 (0.4-0.9)** |
| Africa | -32.8 (-57.8 to -8.0)** | 0.4 (0.2-0.8)* |
| Multiple continents | -7.6 (-18.3-3.2) | 0.5 (0.3-0.7)*** |
| Setting | | |
| Community | 59.8 (57.2-62.5) (intercept) | Ref |
| Primary care | 3.5 (-2.5-9.6) | 1.7 (1.4-2.1)*** |
| Hospitals | 10.2 (1.5-19.0)* | 1.8 (1.3-2.4)*** |
| Study population | | |
| All adults | 48.3 (46.6-50.0) (intercept) | Ref |
| Middle-aged and older | 15.4 (12.7-18.0)*** | 0.9 (0.7-1.1) |
| Only older people | 26.2 (23.7-28.7)*** | 1.2 (0.9-1.4) |

* <0.05 ** <0.01 *** <0.001

Ref: Reference category. lm: Linear regression. nb: Negative binomial regression

Table S6: Risk of bias assessment of included studies

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{25.} Aarts et al (2012) | Moderate | Moderate | Moderate | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{26.} Aarts et al (2011) | Low | High | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | No |
| ^{27.} Aarts et al (2011) | Moderate | Moderate | Moderate | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{28.} Abizanda et al (2014) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{29.} Agborsangaya et al (2012) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{30.} Agborsangaya et al (2013) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{31.} Agborsangaya et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{32.} Ahrenfeldt et al (2019) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | No |
| ^{33.} Alimohammadian et al (2017) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | Yes |
| ^{34.} Angst et al (2002) | Moderate | Moderate | Moderate | High | Low | High | High | Unclear | High | No |
| ^{35.} Appa et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{36.} Adams et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{37.} Ahmadi et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{38.} Amaral et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{39.} An et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{40.} Araujo et al (2018) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{41.} Arnold-Reed et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{42.} Arokiasamy et al (2015) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{43.} Sinnige et al (2015) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{44.} Zemedikun et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{45.} Wensing et al (2001) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Unclear | Moderate | Yes |
| ^{46.} Mounce et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{47.} Taylor et al (2010) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Low | Yes |
| ^{48.} Vancampfort et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{49.} Vancampfort et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{50.} Aubert et al (2016) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{51.} Autenrieth et al (2013) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{52.} Bahler et al (2015) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{53.} Vancampfort et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{54.} Banjare et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{55.} Barra et al (2015) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{56.} Bernard et al (2016) | High | Moderate | High | High | Moderate | Low | Moderate | Low | High | No |
| ^{57.} Biswas et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{58.} Blakemore et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{59.} Blyth et al (2008) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{60.} Bowling et al (2019) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{61.} Britt et al (2008) | Low | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{62.} Broeiro-Goncalves (2019) | Low | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{63.} Bruce et al (2010) | High | Moderate | Moderate | High | Low | High | Moderate | Unclear | High | No |
| ^{64.} Burgers et al (2010) | Low | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{65.} Burke et al (2017) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{66.} Buurman et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{67.} Calderon-Larranaga et al (2017) | Moderate | Moderate | Moderate | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{68.} Camargo-Casas et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{69.} Canevelli et al (2019) | High | High | High | High | Moderate | High | Moderate | Low | High | Yes |
| ^{70.} Chamberlain et al (2020) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Low | Yes |
| ^{71.} Chen et al (2018) | Low | Moderate | High | High | Low | Low | Moderate | Low | Low | Yes |
| ^{72.} Chen et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{73.} Cheung et al (2013) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{74.} Chu et al (2018) | Low | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{75.} Chudasama et al (2019) | Moderate | Moderate | Low | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{76.} Cimarras-Otal et al (2014) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{77.} Chin et al (2016) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{78.} Agrawal et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{79.} Gu et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{80.} Gunn et al (2012) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{81.} Han et al (2013) | High | High | Moderate | High | Moderate | High | Moderate | Unclear | High | No |
| ^{82.} Hanlon et al (2018) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Low | Yes |
| ^{83.} Jantsch et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{84.} John et al (2003) | Moderate | High | Moderate | High | Low | High | Moderate | Low | High | No |
| ^{85.} Johnson-Lawrence et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{86.} Johnston et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{87.} Jones et al (2016) | Low | Moderate | Moderate | High | Low | Low | Moderate | Unclear | Moderate | Yes |
| ^{88.} Jovic et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{89.} Juul-Larsen et al (2020) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{90.} Hudon et al (2008) | Low | Moderate | Moderate | High | Low | Low | Moderate | Low | Low | Yes |
| ^{91.} Hussain et al (2015) | Moderate | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{92.} Ie et al (2017) | High | High | Moderate | High | Moderate | Low | Moderate | Low | High | Yes |
| ^{93.} Ishizaki et al (2019) | Moderate | Moderate | Low | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{94.} Danon-Hersch et al (2012) | Moderate | High | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{95.} de Heer et al (2013) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{96.} Demirchyan et al (2013) | High | Moderate | Low | High | Moderate | High | Moderate | Low | High | No |
| ^{97.} Fabbri et al (2015) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{98.} Fillenbaum et al (2000) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{99.} Kaneko et al (2019) | Moderate | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | No |
| ^{100.} Kang et al (2017) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{101.} Gandhi et al (2020) | Moderate | Moderate | Moderate | High | High | High | Moderate | Low | Moderate | Yes |
| ^{102.} Costa et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{103.} Rizzuto et al (2017) | High | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{104.} Dhalwani et al (2017) | Moderate | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{105.} Elixhauser et al (1998) | Low | Moderate | High | High | Low | Low | Moderate | Unclear | Low | Yes |
| ^{106.} Fabbri et al (2015) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{107.} Fortin et al (2014) | Low | Moderate | Low | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{108.} Fuchs et al (1998) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | No |
| ^{109.} Galenkamp et al (2011) | Low | Moderate | High | High | Moderate | High | Moderate | Unclear | High | No |
| ^{110.} Galenkamp et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{111.} Gamma et al (2001) | High | Moderate | High | High | Moderate | High | Moderate | Unclear | High | No |
| ^{112.} Ge et al (2018) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{113.} Ge et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{114.} Gould et al (2016) | Moderate | Moderate | Moderate | High | High | High | Moderate | Unclear | Moderate | Yes |
| ^{115.} Habib et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{116.} Harrison et al (2017) | Low | Moderate | High | High | Moderate | Low | Moderate | Unclear | Moderate | No |
| ^{117.} Hayek et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{118.} Henninger et al (2012) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{119.} Hernandez et al (2019) | Moderate | Moderate | Moderate | High | High | High | Moderate | Unclear | Moderate | Yes |
| ^{120.} Ho et al (2014) | Moderate | Moderate | High | High | Low | Low | Moderate | Low | Moderate | Yes |
| ^{121.} Khan et al (2019) | Low | Moderate | Low | High | Low | High | Moderate | Low | Low | Yes |
| ^{122.} Kiliari et al (2013) | High | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | No |
| ^{123.} King et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{124.} Kingston et al (2018) | Low | Moderate | High | High | Moderate | High | Moderate | Unclear | High | Yes |
| ^{125.} Koyanagi et al (2018) | Low | Moderate | Moderate | High | Moderate | Low | High | Low | Moderate | Yes |
| ^{126.} Kriegsman et al (2004) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{127.} Kristensen et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{128.} Kristensen et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{129.} Kunna et al (2017) | Low | Moderate | Low | High | Moderate | Low | High | Low | Low | Yes |
| ^{130.} Kuwornu et al (2014) | Moderate | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{131.} Lai et al (2019) | Low | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{132.} Lai et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{133.} Laires et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{134.} Lang et al (2015) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{135.} Le Cossec et al (2016) | Low | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{136.} Lee et al (2007) | Low | Moderate | High | High | Low | Low | Moderate | Low | Low | Yes |
| ^{137.} Lee et al (2018) | Low | Moderate | High | High | High | Low | Moderate | Unclear | High | No |
| ^{138.} Li et al (2016) | Low | Low | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{139.} Li et al (2019) | Low | Moderate | Low | High | Moderate | Moderate | Moderate | Low | Moderate | No |
| ^{140.} Lujic et al (2017) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | Yes |
| ^{141.} LupianezUnclearVillanueva et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{142.} Zhou et al (2018) | Moderate | Moderate | Moderate | High | Moderate | Low | High | Low | Moderate | Yes |
| ^{143.} Zhang et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{144.} Wong et al (2010) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{145.} Weimann et al (2016) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{146.} Wang et al (2017) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{147.} Wang et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{148.} Wade et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{149.} Maciejewski et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{150.} Marengoni et al (2016) | Moderate | Moderate | High | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{151.} Marengoni et al (2009) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{152.} Marques et al (2018) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{153.} Mavaddat et al (2014) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{154.} McDaid et al (2013) | Low | Moderate | High | High | Moderate | High | Moderate | Low | High | Yes |
| ^{155.} Melis et al (2014) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{156.} Min et al (2007) | High | Moderate | High | High | Moderate | High | Moderate | Unclear | High | Yes |
| ^{157.} Momtaz et al (2010) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{158.} Mondor et al (2018) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{159.} Muggah et al (2012) | Low | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | No |
| ^{160.} Nagel et al (2008) | Low | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{161.} Niedzwiedz et al (2019) | Moderate | Moderate | High | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{162.} Nunes et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{163.} Nunes et al (2017) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | Yes |
| ^{164.} Nunes et al (2015) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{165.} Olaya et al (2017) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{166.} Olivares et al (2017) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{167.} Park et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{168.} Patel et al (2006) | Moderate | Moderate | High | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{169.} Pati et al (2016) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{170.} Pati et al (2019) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{171.} Payne et al (2013) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{172.} Perez et al (2020) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{173.} Petersen et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{174.} Pfortmueller et al (2013) | Moderate | Moderate | High | High | High | High | Moderate | Unclear | High | No |
| ^{175.} Pressley et al (1999) | Low | Moderate | High | High | Moderate | Low | Moderate | Unclear | Moderate | No |
| ^{176.} Prior et al (2016) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{177.} Ribeiro et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{178.} Ruel et al (2014) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{179.} Ruel et al (2014) | Moderate | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | Yes |
| ^{180.} Ryan et al (2018) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{181.} Schmidt et al (2016) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{182.} Schottker et al (2016) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |
| ^{183.} Seo et al (2017) | Low | Moderate | Moderate | High | Low | High | Moderate | Low | Moderate | No |
| ^{184.} She et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{185.} Singh et al (2019) | Low | Moderate | Moderate | High | Low | Low | Moderate | Unclear | Moderate | Yes |
| ^{186.} Stepanova et al (2015) | Low | High | High | High | High | High | High | Unclear | High | Yes |
| ^{187.} Stickley et al (2020) | Low | Moderate | High | High | Moderate | High | Moderate | Low | High | Yes |
| ^{188.} Streit et al (2014) | Moderate | Moderate | Moderate | High | High | High | Moderate | Unclear | Moderate | Yes |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|--|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{189.} Stubbs et al (2018) | Low | Moderate | High | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{190.} Su et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{191.} Sundstrup et al (2017) | Low | Moderate | High | High | Moderate | High | Moderate | Unclear | High | Yes |
| ^{192.} Takahashi et al (2016) | Moderate | Moderate | High | High | High | Low | Moderate | Low | High | No |
| ^{193.} Tinetti et al (2011) | Low | Moderate | High | High | High | High | Moderate | Unclear | High | No |
| ^{194.} Troelstra et al (2020) | High | Moderate | High | High | Moderate | Low | Moderate | Unclear | High | Yes |
| ^{195.} van Zon et al (2020) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{196.} Vancampfort et al (2017) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{197.} Vassilaki et al (2015) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{198.} Vassilaki et al (2016) | Low | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{199.} Villarreal et al (2015) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{200.} Violan et al (2019) | Low | Moderate | Moderate | High | High | Low | Moderate | Low | Moderate | Yes |
| ^{201.} von Strauss et al (2000) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Low | Moderate | No |
| ^{202.} Vos et al (2013) | Moderate | Moderate | High | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{203.} Vu et al (2019) | High | Moderate | High | High | Moderate | High | Moderate | Low | High | No |

| Author | Selection bias | Study design | Confounders | Blinding | Data collection method | Withdrawals and dropouts | Publication bias | Conflict of interest | Overall rating | Reporting of MM measure and definition |
|---|----------------|--------------|-------------|----------|------------------------|--------------------------|------------------|----------------------|----------------|--|
| ^{204.} Wang et al (2018) | Moderate | Moderate | Low | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{205.} Wang et al (2017) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{206.} Wijers et al (2019) | Low | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | No |
| ^{207.} Williams et al (2016) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | No |
| ^{208.} Woldeamayrat et al (2018) | High | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{209.} Yao et al (2020) | Moderate | Moderate | Low | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{210.} Yorke et al (2017) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{211.} You et al (2019) | Moderate | Moderate | Moderate | High | Moderate | High | Moderate | Low | Moderate | Yes |
| ^{212.} Zhang et al (2020) | Moderate | Moderate | Low | High | Moderate | Low | Moderate | Low | Moderate | Yes |
| ^{213.} Khanam et al (2011) | Moderate | Moderate | Moderate | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{214.} Cornell et al (2009) | Low | Moderate | High | High | Moderate | Low | Moderate | Unclear | Moderate | Yes |
| ^{215.} Cassell et al (2018) | Low | Moderate | High | High | Moderate | Moderate | Moderate | Low | Moderate | No |
| ^{216.} Wong et al (2019) | High | Moderate | Moderate | High | Moderate | High | Moderate | Unclear | Moderate | Yes |
| ^{217.} Puth et al (2017) | Moderate | Moderate | Moderate | High | Moderate | Moderate | Moderate | Low | Moderate | Yes |

Table S7: Output of adjusted meta-analytic model based on 217 studies

| | Pooled prevalence of multimorbidity of each subgroup (%; 95% CI) | Meta-regression Unadjusted Odds Ratio (95% CI) | Meta-regression Adjusted Odds Ratio (95% CI) R ² 42.4% | FMI |
|--------------------------|--|--|--|------|
| Group of mean age | | R ² 27.0% | | |
| <59 | 30.4 (27.0-33.9) | Ref | Ref | Ref |
| 59-73 | 43.5 (38.0-49.1) | 1.8 (1.3-2.3)*** | 2.0 (1.6-2.6)*** | 0.3 |
| ≥74 | 67.8 (61.3-73.7) | 6.4 (4.6-8.9)*** | 4.7 (3.4-6.5)*** | 0.2 |
| No of conditions | | R ² 6.9% | | |
| <9 | 29.9 (24.9-35.4) | Ref | Ref | Ref |
| 9-19 | 43.5 (39.1-47.9) | 1.8 (1.3-2.5)*** | 1.7 (1.3-2.2)*** | 0.1 |
| 20-43 | 46.7 (38.4-55.2) | 2.1 (1.4-3.1)*** | 2.2 (1.5-3.3)*** | 0.2 |
| ≥44 | 54.5 (32.6-74.8) | 2.8 (1.5-5.4)** | 2.8 (1.6-4.8)*** | 0.1 |
| Setting | | R ² 3.7% | | |
| Community | 37.8 (34.4-41.4) | Ref | Ref | Ref |
| Primary care | 51.2 (41.6-60.7) | 1.7 (1.2-2.5)** | 1.8 (1.2-2.6)** | 0.1 |
| Hospital | 47.1 (31.9-63.0) | 1.5 (0.9-2.4) | 0.8 (0.5-1.3) | 0.1 |
| Care home | 73.9 (72.8-74.9) | 4.6 (0.6-36.6) | 1.5 (0.3-8.4) | 0.04 |
| Source | | R ² 2.8% | | |
| Self-report | 38.3 (34.4-42.2) | Ref | Ref | Ref |
| Database | 48.9 (42.2-55.6) | 1.5 (1.1-2.1)** | 0.8 (0.6-1.1) | 0.1 |
| Continent | | R ² 7.4% | | |
| North America | 48.9 (42.1-55.7) | Ref | Ref | Ref |
| Europe | 44.0 (37.7-50.4) | 0.8 (0.6-1.2) | 0.5 (0.4-0.7)*** | 0.1 |
| Australasia | 28.2 (20.3-37.6) | 0.4 (0.2-0.8)** | 0.4 (0.2-0.6)*** | 0.08 |
| Asia | 34.3 (28.6-40.5) | 0.5 (0.4-0.8)** | 0.5 (0.3-0.7)*** | 0.1 |
| South America | 47.5 (31.2-64.4) | 0.9 (0.5-1.8) | 0.8 (0.5-1.3) | 0.1 |
| Africa | 13.8 (4.5-35.2) | 0.2 (0.06-0.5)*** | 0.2 (0.1-0.5)*** | 0.1 |
| Multiple continents | 41.4 (31.0-52.6) | 0.7 (0.4-1.4) | 0.7 (0.4-1.2) | 0.1 |


*<0.05 **<0.01 ***<0.001

Ref: Reference category. FMI: Fraction of missing information.

Table S8: Definition of variables

| Variable name | Definition |
|--|---|
| Study setting | |
| Community | Studies that used population surveys, insurance claims databases, or research databases |
| Primary care | Studies that were carried out in primary care settings |
| Hospital | Studies that were carried out in hospital settings |
| Data source | |
| Self-report | Studies that collected data using self-report or interviews |
| Medical records and administrative databases | Studies that collected data using electronic medical records, medical chart reviews, insurance claims databases, pharmacy databases, or research databases |
| Study population | |
| All adults | Studies with a sample of population aged 18 and older (n=45), aged 20 and older (n=8), aged 21 and older (n=3), aged 25 and older (n=2), or others (n=27) (e.g. aged 16 and older, or aged 17 and older) |
| Middle-aged and older | Studies with a sample of population aged 50 and older (n=25), aged 40 and older (n=5), aged 40 and older (n=10), or others (n=6) (e.g. aged 57 and older, or aged 45 and older) |
| Only older people | Studies with a sample of population aged 65 and older (n=22), aged 60 and older (n=25), aged 70 and older (n=5) or others (n=11) (e.g. aged 68 and older, aged 77 and older, aged 78 and older, or aged 80 and older) |

Figure S1: Graphical display of study effect sizes and heterogeneity



No obvious subgroup effects were identified

Figure S2: Process of examining and identifying outlying studies in meta-analysis






Figure S3: Summary of risk of bias assessment

Figure S4: Meta-regression trees for predicting the pooled estimated prevalence of multimorbidity (based on ‘mean age’ and ‘number of conditions’ predictors. unit: log(odds))



References

- 1 Stanley J, Semper K, Millar E, Sarfati D. Epidemiology of multimorbidity in New Zealand: A cross-sectional study using national-level hospital and pharmaceutical data. *BMJ Open* 2018; **8**:1-12.
- 2 Lenzi J, Avaldi VM, Rucci P, Pieri G, Fantini MP. Burden of multimorbidity in relation to age, gender and immigrant status: A cross-sectional study based on administrative data. *BMJ Open* 2016; **6**:1-12.
- 3 Hu RH, Hsiao FY, Chen LJ, Huang PT, Hsu WW. Increasing age- and gender-specific burden and complexity of multimorbidity in taiwan, 2003-2013: A cross-sectional study based on nationwide claims data. *BMJ Open* 2019; **9**:1-10.
- 4 Gawron LM, Sanders JN, Sward K, Poursaid AE, Simmons R, Turok DK. Multimorbidity and highly effective contraception in reproductive-age women in the us intermountain west: A retrospective cohort study. *J Gen Intern Med* 2020; **35**:637-42.
- 5 Low LL, Kwan YH, Ko MSM, et al. Epidemiologic characteristics of multimorbidity and sociodemographic factors associated with multimorbidity in a rapidly aging asian country. *JAMA Network Open* 2019; **2**:1-14.
- 6 Wang HHX, Wang JJ, Wong SYS, et al. Epidemiology of multimorbidity in china and implications for the healthcare system: Cross-sectional survey among 162,464 community household residents in southern china. *BMC Medicine* 2014; **12**:1-12.
- 7 Gaulin M, Simard M, Candas B, Lesage A, Sirois C. Combined impacts of multimorbidity and mental disorders on frequent emergency department visits: A retrospective cohort study in quebec, canada. *Canadian Medical Association Journal* 2019; **191**:E724-E32.
- 8 Violan C, Foguet-Boreu Q, Flores-Mateo G, et al. Prevalence, determinants and patterns of multimorbidity in primary care: A systematic review of observational studies. *PLOS One* 2014; **9**:1-9.
- 9 Nicholson K, Terry AL, Fortin M, Williamson T, Bauer M, Thind A. Prevalence, characteristics, and patterns of patients with multimorbidity in primary care: A retrospective cohort analysis in canada. *British Journal of General Practice* 2019; **69**:1-10.
- 10 Bao X-Y, Xie Y-X, Zhang X-X, et al. The association between multimorbidity and health-related quality of life: A cross-sectional survey among community middle-aged and elderly residents in southern china. *Health and quality of life outcomes* 2019; **17**:107-09.
- 11 Fortin M, Bravo G, Hudon C, Vanasse A, Lapointe L. Prevalence of multimorbidity among adults seen in family practice. *Ann Fam Med* 2005; **3**:223-8.
- 12 Prazeres F, Santiago L. Prevalence of multimorbidity in the adult population attending primary care in portugal: A cross-sectional study. *BMJ Open* 2015; **5**:1-10.
- 13 Lawson KD, Mercer SW, Wyke S, et al. Double trouble: The impact of multimorbidity and deprivation on preference-weighted health related quality of life a cross sectional analysis of the scottish health survey. *Int J Equity Health* 2013; **12**:1-9.
- 14 Sullivan PW, Ghushchyan VH, Bayliss EA. The impact of co-morbidity burden on preference-based health-related quality of life in the United States. *Pharmacoeconomics* 2012; **30**:431-42.
- 15 Peng X, Bao X, Xie Y, et al. The mediating effect of pain on the association between multimorbidity and disability and impaired physical performance among community-dwelling older adults in southern china. *Aging Clin Exp Res* 2020; **32**:1327-34.

- 16 Excoffier S, Herzig L, N'Goran AA, Déruaz-Luyet A, Haller DM. Prevalence of multimorbidity in general practice: A cross-sectional study within the Swiss sentinel surveillance system (sentinella). *BMJ Open* 2018; **8**:1-7.
- 17 Chung RY, Mercer S, Lai FTT, Yip BHK, Wong MCS, Wong SYS. Socioeconomic determinants of multimorbidity: A population-based household survey of Hong Kong Chinese. *PLOS ONE* 2015; **10**:1-15.
- 18 Ki M, Lee YH, Kim Y-S, Shin J-Y, Lim J, Nazroo J. Socioeconomic inequalities in health in the context of multimorbidity: A Korean panel study. *PLOS ONE* 2017; **12**:1-14.
- 19 Bobo WV, Yawn BP, St Sauver JL, Grossardt BR, Boyd CM, Rocca WA. Prevalence of combined somatic and mental health multimorbidity: Patterns by age, sex, and race/ethnicity. *J Gerontol - A Biol Sci Med Sci* 2016; **71**:1483-91.
- 20 Randall DA, Lujic S, Havard A, Eades SJ, Jorm L. Multimorbidity among aboriginal people in New South Wales contributes significantly to their higher mortality. *Med J Aust* 2018; **209**:19-23.
- 21 Russell J, Grant CC, Morton SMB. Multimorbidity in early childhood and socioeconomic disadvantage: Findings from a large New Zealand child cohort. *Acad Pediatr* 2020; **20**:619-27.
- 22 Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: A cross-sectional study. *The Lancet* 2012; **380**:37-43.
- 23 St Sauver JL, Boyd CM, Grossardt BR, et al. Risk of developing multimorbidity across all ages in an historical cohort study: Differences by sex and ethnicity. *BMJ Open* 2015; **5**:1-13.
- 24 Vetrano DL, Foebel AD, Marengoni A, et al. Chronic diseases and geriatric syndromes: The different weight of comorbidity. *Eur J Intern Med* 2016; **27**:62-7.
- 25 Aarts S, den Akker Mv, Bosma H, et al. The effect of multimorbidity on health related functioning: Temporary or persistent? Results from a longitudinal cohort study. *J Psychosom Res* 2012; **73**:211-17.
- 26 Aarts S, van den Akker M, Hajema KJ, et al. Multimorbidity and its relation to subjective memory complaints in a large general population of older adults. *Int Psychogeriatr* 2011; **23**:616-24.
- 27 Aarts S, van den Akker M, Tan FES, Verhey FRJ, Metsemakers JFM, van Boxtel MPJ. Influence of multimorbidity on cognition in a normal aging population: A 12-year follow-up in the maastricht aging study. *Int J Geriatr Psychiatry* 2011; **26**:1046-53.
- 28 Abizanda P, Romero L, Sanchez-Jurado PM, Martinez-Reig M, Alfonso-Silguero SA, Rodriguez-Manas L. Age, frailty, disability, institutionalization, multimorbidity or comorbidity. Which are the main targets in older adults? *J Nutr; Health Aging* 2014; **18**:622-27.
- 29 Agborsangaya CB, Lau D, Lahtinen M, Cooke T, Johnson JA. Health-related quality of life and healthcare utilization in multimorbidity: Results of a cross-sectional survey. *Qual Life Res* 2012; **22**:791-99.
- 30 Agborsangaya CB, Ngwakongnwi E, Lahtinen M, Cooke T, Johnson JA. Multimorbidity prevalence in the general population: The role of obesity in chronic disease clustering. *BMC Public Health* 2013; **13**:1161-61.
- 31 Agborsangaya CB, Lahtinen M, Cooke T, Johnson JA. Comparing the eq-5d 3l and 5l: Measurement properties and association with chronic conditions and multimorbidity in the general population. *Health Qual Life Outcomes* 2014; **12**:1-7.

- 32 Ahrenfeldt LJ, Möller S, Thinggaard M, Christensen K, Lindahl-Jacobsen R. Sex differences in comorbidity and frailty in Europe. *Int J Public Health* 2019; **64**:1025-36.
- 33 Alimohammadian M, Majidi A, Yaseri M, et al. Multimorbidity as an important issue among women: Results of a gender difference investigation in a large population-based cross-sectional study in west asia. *BMJ Open* 2017; **7**:1-8.
- 34 Angst J, Sellaro R, Ries Merikangas K. Multimorbidity of psychiatric disorders as an indicator of clinical severity. *Eur Arch Psychiatry Clin Neurosci* 2002; **252**:147-54.
- 35 Appa AA, Creasman J, Brown JS, et al. The impact of multimorbidity on sexual function in middle-aged and older women: Beyond the single disease perspective. *J Sex Med* 2014; **11**:2744-55.
- 36 Adams ML, Grandpre J, Katz DL, Shenson D. Linear association between number of modifiable risk factors and multiple chronic conditions: Results from the behavioral risk factor surveillance system. *Prev Med* 2017; **105**:169-75.
- 37 Ahmadi B, Alimohammadian M, Yaseri M, et al. Multimorbidity: Epidemiology and risk factors in the golesan cohort study, iran a cross-sectional analysis. *Medicine* 2016; **95**:1-7.
- 38 Amaral TLM, Amaral CdA, de Lima NS, Herculano PV, Do Prado PR, Monteiro GTR. Multimorbidity, depression and quality of life among elderly people assisted in the family health strategy in senador guiomard, acre, Brazil. *Ciencia e Saude Coletiva* 2018; **23**:3077-84.
- 39 An KO Kim J. Association of sarcopenia and obesity with multimorbidity in Korean adults: A nationwide cross-sectional study. *J Am Med Dir Assoc* 2016; **17**:e1-7.
- 40 Araujo MEA, Silva MT, Galvao TF, Nunes BP, Pereira MG. Prevalence and patterns of multimorbidity in amazon region of Brazil and associated determinants: A cross-sectional study. *BMJ Open* 2018; **8**:1-11.
- 41 Arnold-Reed D, Troeung L, Brett T, et al. Increasing multimorbidity in an Australian street health service: A 10-year retrospective cohort study. *Aust J Gen Pract* 2018; **47**:181-89.
- 42 Arokiasamy P, Uttamacharya U, Jain K, et al. The impact of multimorbidity on adult physical and mental health in low- and middle-income countries: What does the study on global ageing and adult health (sage) reveal? *BMC Med* 2015; **13**:1-16.
- 43 Sinnige J, Korevaar JC, Westert GP, Spreeuwenberg P, Schellevis FG, Braspenning JC. Multimorbidity patterns in a primary care population aged 55 years and over. *Fam Pract* 2015; **32**:505-13.
- 44 Zemedikun DT, Gray LJ, Khunti K, Davies MJ, Dhalwani NN. Patterns of multimorbidity in middle-aged and older adults: An analysis of the uk biobank data. *Mayo Clin Proc* 2018; **93**:857-66.
- 45 Wensing M, Vingerhoets E, Grol R. Functional status, health problems, age and comorbidity in primary care patients. *Qual Life Res* 2001; **10**:141-48.
- 46 Mounce LTAP, Campbell JLMD, Henley WEP, Arreal MCTPMPHLCP, Porter IP, Valderas JMBPMPH. Predicting incident multimorbidity. *Ann Fam Med* 2018; **16**:322-29.
- 47 Taylor AW, Price K, Gill TK, et al. Multimorbidity - not just an older person's issue. Results from an Australian biomedical study. *BMC Public Health* 2010; **10**:1-10.
- 48 Vancampfort D, Stubbs B, Firth J, Koyanagi A. Handgrip strength, chronic physical conditions and physical multimorbidity in middle-aged and older adults in six low- and middle income countries. *Eur J Intern Med* 2019; **61**:96-102.

49 Vancampfort D, Smith L, Stubbs B, et al. Associations between active travel and
physical multi-morbidity in six low- and middle-income countries among community-
dwelling older adults: A cross-sectional study. *PloS one* 2018; **13**:1-15.

50 Aubert CE, Streit S, Da Costa BR, et al. Polypharmacy and specific comorbidities in
university primary care settings. *Eur J Intern Med* 2016; **35**:35-42.

51 Autenrieth CS, Kirchberger I, Heier M, et al. Physical activity is inversely associated
with multimorbidity in elderly men: Results from the kora-age augsburg study. *Prev
Med* 2013; **57**:17-19.

52 Bähler C, Huber CA, Brüngger B, Reich O. Multimorbidity, health care utilization
and costs in an elderly community-dwelling population: A claims data based
observational study. *BMC Health Services Res* 2015; **15**:1-12.

53 Vancampfort DK, A.: Ward, P. B.: Veronese, N.: Carvalho, A. F.: Solmi, M.: Mugisha,
J.: Rosenbaum, S.: De Hert, M.: Stubbs, B. Perceived stress and its relationship with
chronic medical conditions and multimorbidity among 229,293 community-dwelling
adults in 44 low- and middle-income countries. *Am J Epidemiol* 2017; **186**:979-89.

54 Banjare P Pradhan J. Socio-economic inequalities in the prevalence of multi-
morbidity among the rural elderly in Bargarh district of Odisha (India). *PLoS ONE*
2014; **9**:1-10.

55 Barra M, Augestad LA, Whitehurst DGT, Rand-Hendriksen K. Examining the
relationship between health-related quality of life and increasing numbers of
diagnoses. *Qual life Res* 2015; **24**:2823-32.

56 Bernard S, Inderjeeth C, Raymond W. Higher charlson comorbidity index scores do
not influence functional independence measure score gains in older rehabilitation
patients. *Aust J Ageing* 2016; **35**:236-41.

57 Biswas T, Townsend N, Islam MS, et al. Association between socioeconomic status
and prevalence of non-communicable diseases risk factors and comorbidities in
bangladesh: Findings from a nationwide cross-sectional survey. *BMJ Open* 2019; **9**:1-
9.

58 Blakemore A, Hann M, Howells K, et al. Patient activation in older people with long-
term conditions and multimorbidity: Correlates and change in a cohort study in the
united kingdom. *BMC Health Services Res* 2016; **16**:582-82.

59 Blyth FM, Rochat S, Cumming RG, et al. Pain, frailty and comorbidity on older men:
The champ study. *Pain* 2008; **140**:224-30.

60 Bowling CB, Deng L, Sakhuja S, Morey MC, Jaeger BC, Muntner P. Prevalence of
activity limitations and association with multimorbidity among us adults 50 to
64 years old. *J Gen Intern Med* 2019; **34**:2390-96.

61 Britt HC, Harrison CM, Miller GC, Knox SA. Prevalence and patterns of
multimorbidity in australia. *Med J Aust* 2008; **189**:72-77.

62 Broeiro-Gonçalves P, Nogueira P, Aguiar P. Multimorbidity and disease severity by
age groups, in inpatients: Cross-sectional study. *Portuguese J Public Health* 2019;
37:1-9.

63 Bruce SG, Riediger ND, Zacharias JM, Young TK. Obesity and obesity-related
comorbidities in a canadian first nation population. *Chronic Dis Can* 2010; **31**:27-32.

64 Burgers JS, Voerman GE, Grol R, Faber MJ, Schneider EC. Quality and coordination
of care for patients with multiple conditions: Results from an international survey of
patient experience. *Eval Health Prof* 2010; **33**:343-64.

65 Burke BT. *Connections that count: A network analysis of multimorbidity and gait
speed in older adults*, in *Department of Epidemiology (dissertation)*. 2017, Johns
Hopkins University: Baltimore, Maryland.1-224.

- 66 Buurman BM, Frenkel WJ, Abu-Hanna A, Parlevliet JL, de Rooij SE. Acute and chronic diseases as part of multimorbidity in acutely hospitalized older patients. *Eur J Intern Med* 2016; **27**:68-75.
- 67 Calderón-Larrañaga A, Vetrano DL, Onder G, et al. Assessing and measuring chronic multimorbidity in the older population: A proposal for its operationalization. *J Gerontol A Biol Sci Med Sci* 2017; **72**:1417-23.
- 68 Camargo-Casas S, Suarez-Monsalve S, Zepeda MUP, García-Peña C, Cano-Gutierrez CA. Multimorbidity, depressive symptoms, and self-reported health in older adults: A secondary analysis of the sabe bogota study. *Rev Invest Clin* 2018; **70**:192-97.
- 69 Canevelli M, Raganato R, Remiddi F, et al. Counting deficits or diseases? The agreement between frailty and multimorbidity in subjects with cognitive disturbances. *Aging Clin Exp Res* 2019; **32**:179-82.
- 70 Chamberlain AM, Rutten LJF, Wilson PM, et al. Neighborhood socioeconomic disadvantage is associated with multimorbidity in a geographically-defined community. *BMC Public Health* 2020; **20**:1-10.
- 71 Chen H, Chen Y, Cui B. The association of multimorbidity with healthcare expenditure among the elderly patients in Beijing, China. *Arch Gerontol Geriatr* 2018; **79**:32-38.
- 72 Chen H, Cheng M, Zhuang Y, Broad JBJG, International G. Multimorbidity among middle-aged and older persons in urban china: Prevalence, characteristics and health service utilization. *Geriatr Gerontol Int* 2018; **18**:1447-52.
- 73 Cheung CL, Nguyen US, Au E, Tan KC, Kung AW. Association of handgrip strength with chronic diseases and multimorbidity: A cross-sectional study. *Age (Dordr)* 2013; **35**:929-41.
- 74 Chu T, Lau P, Cheng R, Chan M, Liang J. A study on the prevalence of multimorbidities of diseases and utilisation of public healthcare services in the new territories west area of Hong Kong. *Hong Kong Practitioner* 2018; **40**:43-50.
- 75 Chudasama YV, Khunti KK, Zaccardi F, et al. Physical activity, multimorbidity, and life expectancy: A UK biobank longitudinal study. *BMC Med* 2019; **17**:1-13.
- 76 Cimarras-Otal C, Calderón-Larrañaga A, Poblador-Plou B, et al. Association between physical activity, multimorbidity, self-rated health and functional limitation in the Spanish population. *BMC Public Health* 2014; **14**:1-10.
- 77 Chin WY, Choi EP, Wan EY, Lam CL. Health-related quality of life mediates associations between multi-morbidity and depressive symptoms in Chinese primary care patients. *Fam Pract* 2016; **33**:61-8.
- 78 Agrawal S, Agrawal PK. Association between body mass index and prevalence of multimorbidity in low-and middle-income countries: A cross-sectional study. *Int J Med Public Health* 2016; **6**:73-83.
- 79 Gu J, Chao J, Chen W, et al. Multimorbidity and health-related quality of life among the community-dwelling elderly: A longitudinal study. *Arch Gerontol Geriatr* 2018; **74**:133-40.
- 80 Gunn JM, Ayton DR, Densley K, et al. The association between chronic illness, multimorbidity and depressive symptoms in an australian primary care cohort. *Soc Psychiatry Psychiatr Epidemiol* 2010; **47**:175-84.
- 81 Han MA, Tinetti ME, Agostini JV, Han L, Lee H, Walke LM. The impact of medication use on symptom burden in older patients with multiple medical morbidities. *J Clin Gerontol and Geriatr* 2013; **4**:84-88.
- 82 Hanlon P, Nicholl BI, Jani BD, Lee D, McQueenie R, Mair FS. Frailty and pre-frailty in middle-aged and older adults and its association with multimorbidity and mortality:

- A prospective analysis of 493 737 UK biobank participants. *Lancet Public Health* 2018; **3**:e323-e32.
- 83 Jantsch AG, Alves RFS, Faerstein E. Educational inequality in Rio de Janeiro and its impact on multimorbidity: Evidence from the pró-saúde study. A cross-sectional analysis. *Sao Paulo Med J* 2018; **136**:51-58.
- 84 John R, Kerby DS, Hagan Hennessy C. Patterns and impact of comorbidity and multimorbidity among community-resident American Indian elders. *Gerontol* 2003; **43**:649-60.
- 85 Johnson-Lawrence V, Zajacova A, Sneed R. Education, race/ethnicity, and multimorbidity among adults aged 30-64 in the national health interview survey. *SSM Popul Health* 2017; **3**:366-72.
- 86 Johnston MC, Black C, Mercer SW, Prescott GJ, Crilly MA. Impact of educational attainment on the association between social class at birth and multimorbidity in middle age in the Aberdeen children of the 1950s cohort study. *BMJ Open* 2019; **9**:1-9.
- 87 Jones SM, Amtmann D, Gell NM. A psychometric examination of multimorbidity and mental health in older adults. *Aging Ment Health* 2016; **20**:309-17.
- 88 Jovic D, Marinkovic J, Vukovic D. Association between body mass index and prevalence of multimorbidity: A cross-sectional study. *Public Health* 2016; **139**:103-11.
- 89 Juul-Larsen HG, Andersen O, Bandholm T, et al. Differences in function and recovery profiles between patterns of multimorbidity among older medical patients the first year after an acute admission—an exploratory latent class analysis. *Arch Gerontol Geriatr* 2020; **86**:1-11.
- 90 Hudon C, Soubhi H, Fortin M. Relationship between multimorbidity and physical activity: Secondary analysis from the Quebec health survey. *BMC Public Health* 2008; **8**:1-8.
- 91 Hussain MA, Huxley RR, Al Mamun A. Multimorbidity prevalence and pattern in Indonesian adults: An exploratory study using national survey data. *BMJ Open* 2015; **5**:1-10.
- 92 Ie K, Felton M, Springer S, Wilson SA, Albert SM. Multimorbidity and polypharmacy in family medicine residency practices. *J Pharm Technol* 2017; **33**:219-24.
- 93 Ishizaki T, Kobayashi E, Fukaya T, Takahashi Y, Shinkai S, Liang J. Association of physical performance and self-rated health with multimorbidity among older adults: Results from a nationwide survey in japan. *Arch Gerontol Geriatr* 2019; **84**:1-9.
- 94 Danon-Hersch N, Rodondi N, Spagnoli J, Santos-Eggimann B. Prefrailty and chronic morbidity in the youngest old: An insight from the Lausanne cohort Lc65+. *J Am Geriatr Soc* 2012; **60**:1687-94.
- 95 de Heer HD, Balcázar HG, Morera OF, et al. Barriers to care and comorbidities along the u.S.-mexico border. *Public Health Rep* 2013; **128**:480-8.
- 96 Demirchyan A, Khachadourian V, Armenian HK, Petrosyan V. Short and long term determinants of incident multimorbidity in a cohort of 1988 earthquake survivors in Armenia. *Int. J. Equity Health* 2013; **12**:68-68.
- 97 Fabbri E, An Y, Zoli M, et al. Aging and the burden of multimorbidity: Associations with inflammatory and anabolic hormonal biomarkers. *J Gerontol A Biol Sci Med Sci* 2015; **70**:63-70.
- 98 Fillenbaum GGP, C. F.: Cohen, H. J.: Cornoni-Huntley, J. C.: Guralnik, J. M. Comorbidity of five chronic health conditions in elderly community residents: Determinants and impact on mortality. *J Gerontol: Med Sci* 2000; **55**:M84-89.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
- 99 Kaneko M, Boven Kv, Takayanagi H, Kusaba T, Yamada T, Matsushima MJFp. Multicentre descriptive cross-sectional study of japanese home visit patients: Reasons for encounter, health problems and multimorbidity. *Fam Pract* 2019; **37**:227-33.
- 100 Kang J Song Y-M. The association between submarine service and multimorbidity: A cross-sectional study of korean naval personnel. *BMJ Open* 2017; **7**:1-6.
- 101 Gandhi K, Lim E, Davis J, Chen JJ. Racial-ethnic disparities in self-reported health status among us adults adjusted for sociodemographics and multimorbidities, national health and nutrition examination survey 2011-2014. *Ethn Health* 2020; **25**:65-78.
- 102 Costa CDS, Flores TR, Wendt A, et al. Inequalities in multimorbidity among elderly: A population-based study in a city in southern brazil. *Cad Saude Publica* 2018; **34**:1-12.
- 103 Rizzuto DM, R. J. F.: Angleman, S.: Qiu, C.: Marengoni, A. Effect of chronic diseases and multimorbidity on survival and functioning in elderly adults. *J Am Geriatr Soc* 2017; **65**:1056-60.
- 104 Dhalwani NN, Zaccardi F, O'Donovan G, et al. Association between lifestyle factors and the incidence of multimorbidity in an older english population. *J Gerontol A Biol Sci Med Sci* 2017; **72**:528-34.
- 105 Elixhauser A, Steiner C, Harris DR, Coffey RM. Comorbidity measures for use with administrative data. *Med Care* 1998; **36**:8-27.
- 106 Fabbri E, An Y, Schrack JA, et al. Energy metabolism and the burden of multimorbidity in older adults: Results from the baltimore longitudinal study of aging. *J Gerontol A Biol Sci Med Sci* 2015; **70**:1297-303.
- 107 Fortin M, Haggerty J, Almirall J, Bouhali T, Sasseville M, Lemieux M. Lifestyle factors and multimorbidity: A cross sectional study. *BMC Public Health* 2014; **14**:1-6.
- 108 Fuchs ZB, T.: Novikov, I.: Walter-Ginzburg, A.: Lyanders, M.: Gindin, J.: Habet, B.: Modan, B. Morbidity, comorbidity, and their association with disability among community-dwelling oldest-old in israel. *J Gerontol: Med Sci* 1998; **53**:M447-55.
- 109 Galenkamp H, Braam AW, Huisman M, Deeg DJ. Somatic multimorbidity and self-rated health in the older population. *J Gerontol B Psychol Sci Soc Sci* 2011; **66**:380-6.
- 110 Galenkamp H, Gagliardi C, Principi A, et al. Predictors of social leisure activities in older europeans with and without multimorbidity. *Eur J Ageing* 2016; **13**:129-43.
- 111 Gamma A Angst J. Concurrent psychiatric comorbidity and multimorbidity in a community study: Gender differences and quality of life. *Eur Arch Psychiatry Clin Neurosci* 2001; **251**:43-46.
- 112 Ge L, Yap CW, Heng BH. Sex differences in associations between multimorbidity and physical function domains among community-dwelling adults in Singapore. *PLoS One* 2018; **13**:1-14.
- 113 Ge L, Ong R, Yap CW, Heng BH. Effects of chronic diseases on health-related quality of life and self-rated health among three adult age groups. *Nurs Health Sci* 2019; **21**:214-22.
- 114 Gould CE, O'Hara R, Goldstein MK, Beaudreau SA. Multimorbidity is associated with anxiety in older adults in the health and retirement study. *Int J Geriatr Psychiatry* 2016; **31**:1105-15.
- 115 Habib RR, Hojeij S, Elzein K, Chaaban J, Seyfert K. Associations between life conditions and multi-morbidity in marginalized populations: The case of palestinian refugees. *Eur J Public Health* 2014; **24**:727-33.
- 116 Harrison C, Henderson J, Miller G, Britt H. The prevalence of diagnosed chronic conditions and multimorbidity in australia: A method for estimating population prevalence from general practice patient encounter data. *PLOS ONE* 2017; **12**:1-13.

- 117 Hayek S, Ifrah A, Enav T, Shohat T. Prevalence, correlates, and time trends of multiple chronic conditions among israeli adults: Estimates from the Israeli national health interview survey, 2014-2015. *Prev Chronic Dis* 2017; **14**:1-14.
- 118 Henninger DE, Whitson HE, Cohen HJ, Ariely D. Higher medical morbidity burden is associated with external locus of control. *J Am Geriatr Soc* 2012; **60**:751-55.
- 119 Hernández B, Reilly RB, Kenny RA. Investigation of multimorbidity and prevalent disease combinations in older irish adults using network analysis and association rules. *Sci Rep* 2019; **9**:1-12.
- 120 Ho C, Feng L, Fam J, Mahendran R, Kua EH, Ng TP. Coexisting medical comorbidity and depression: Multiplicative effects on health outcomes in older adults. *Int Psychogeriatr* 2014; **26**:1221-9.
- 121 Khan N, Rahman M, Mitra D, Afsana K. Prevalence of multimorbidity among bangladeshi adult population: A nationwide cross-sectional study. *BMJ Open* 2019; **9**:1-8.
- 122 Kiliari N, Theodosopoulou E, Papanastasiou E. Multimorbidity and unmet citizens' needs and expectations urge for reforms in the health system of cyprus: A questionnaire survey. *JRSM Open* 2014; **5**:1-11.
- 123 King DE, Xiang J, Pilkerton CS. Multimorbidity trends in united states adults, 1988-2014. *J Am Board Fam Med* 2018; **31**:503-13.
- 124 Kingston AR, L.: Booth, H.: Knapp, M.: Jagger, C.: Modem project. Projections of multi-morbidity in the older population in england to 2035: Estimates from the population ageing and care simulation (pacsim) model. *Age Ageing* 2018; **47**:374-80.
- 125 Koyanagi A, Lara E, Stubbs B, et al. Chronic physical conditions, multimorbidity, and mild cognitive impairment in low- and middle-income countries. *J Am Geriatr Soc* 2018; **66**:721-27.
- 126 Kriegsman DMD, D. J.: Stalman, W. A. Comorbidity of somatic chronic diseases and decline in physical functioning;; the longitudinal aging study amsterdam. *J Clin Epidemiol* 2004; **57**:55-65.
- 127 Kristensen K, König HH, Hajek A. The longitudinal association of multimorbidity on loneliness and network size: Findings from a population-based study. *Int J Geriatr Psychiatry* 2019; **34**:1490-97.
- 128 Kristensen K, König HH, Hajek A. The association of multimorbidity, loneliness, social exclusion and network size: Findings from the population-based german ageing survey. *BMC Public Health* 2019; **19**:1-10.
- 129 Kunna R, San Sebastian M, Stewart Williams J. Measurement and decomposition of socioeconomic inequality in single and multimorbidity in older adults in china and ghana: Results from the who study on global ageing and adult health (sage). *Int J Equity Health* 2017; **16**:2-17.
- 130 Kusumastuti S, Gerds TA, Lund R, Mortensen EL, Westendorp RGJ. Discrimination ability of comorbidity, frailty, and subjective health to predict mortality in community-dwelling older people: Population based prospective cohort study. *Eur J Intern Med* 2017; **42**:29-38.
- 131 Lai FTT, Guthrie B, Wong SYS, et al. Sex-specific intergenerational trends in morbidity burden and multimorbidity status in hong kong community: An age-period-cohort analysis of repeated population surveys. *BMJ Open* 2019; **9**:1-10.
- 132 Lai FTT, Ma TW, Hou WK. Multimorbidity is associated with more subsequent depressive symptoms in three months: A prospective study of community-dwelling adults in hong kong. *Int Psychogeriatr* 2018; **31**:1367-71.
- 133 Laires PA Perelman J. The current and projected burden of multimorbidity: A cross-sectional study in a southern europe population. *Eur J Ageing* 2019; **16**:181-92.

- 134 Lang K, Alexander IM, Simon J, et al. The impact of multimorbidity on quality of life among midlife women: Findings from a u.S. Nationally representative survey. *J Women's Health* 2015; **24**:374-83.
- 135 Le Cossec C, Perrine AL, Beltzer N, Fuhrman C, Carcaillon-Bentata L. Pre-frailty, frailty, and multimorbidity: Prevalences and associated characteristics from two french national surveys. *J Nutr Health Aging* 2016; **20**:860-69.
- 136 Lee TA, Shields AE, Vogeli C, et al. Mortality rate in veterans with multiple chronic conditions. *J Gen Intern Med* 2007; **22** 403-7.
- 137 Lee WJ, Peng LN, Lin CH, Lin HP, Loh CH, Chen LK. The synergic effects of frailty on disability associated with urbanization, multimorbidity, and mental health: Implications for public health and medical care. *Sci Rep* 2018; **8**:1-7.
- 138 Li J, Green M, Kearns B, et al. Patterns of multimorbidity and their association with health outcomes within yorkshire, england: Baseline results from the yorkshire health study. *BMC Public Health* 2016; **16**:1-9.
- 139 Li Y, White K, O'Shields KR, McLain AC, Merchant AT. Light-intensity physical activity and cardiometabolic risk among older adults with multiple chronic conditions. *Am J Health Promot* 2019; **33**:507-15.
- 140 Lujic S, Simpson JM, Zwar N, Hosseinzadeh H, Jorm L. Multimorbidity in australia: Comparing estimates derived using administrative data sources and survey data. *PLoS One* 2017; **12**:1-18.
- 141 Lupiáñez-Villanueva F, Anastasiadou D, Codagnone C, Nuño-Solinís R, Garcia-Zapirain Soto MB. Electronic health use in the european union and the effect of multimorbidity: Cross-sectional survey. *J Med Internet Res* 2018; **20**:1-14.
- 142 Zhou C-h, Tang S-f, Wang X-h, et al. Satisfaction about patient-centeredness and healthcare system among patients with chronic multimorbidity. *Current Med Sci* 2018; **38**:184-90.
- 143 Zhang R, Lu Y, Shi L, Zhang S, Chang F. Prevalence and patterns of multimorbidity among the elderly in china: A cross-sectional study using national survey data. *BMJ Open* 2019; **9**:1-9.
- 144 Wong CH, Weiss D, Sourial N, et al. Frailty and its association with disability and comorbidity in a community-dwelling sample of seniors in montreal: A cross-sectional study. *Aging Clin Exp Res* 2010; **22**:54-62.
- 145 Weimann A, Dai D, Oni T. A cross-sectional and spatial analysis of the prevalence of multimorbidity and its association with socioeconomic disadvantage in south africa: A comparison between 2008 and 2012. *Soc Sci Med* 2016; **163**:144-56.
- 146 Wang L, Palmer AJ, Cocker F, Sanderson K. Multimorbidity and health-related quality of life (hrqol) in a nationally representative population sample: Implications of count versus cluster method for defining multimorbidity on hrqol. *Health Qual life Outcomes* 2017; **15**:1-12.
- 147 Wang C, Pu R, Li Z, et al. Subjective health and quality of life among elderly people living with chronic multimorbidity and difficulty in activities of daily living in rural South Africa. *Clin Interv Aging* 2019; **14**:1285-96.
- 148 Wade A, Hobbs M, Green MA. Investigating the relationship between multimorbidity and dental attendance: A cross-sectional study of uk adults. *British Dental Journal* 2019; **226**:138-43.
- 149 Maciejewski ML Hammill BG. Measuring the burden of multimorbidity among medicare beneficiaries via condition counts and cumulative duration. *Health Serv Res* 2019; **54**:484-91.
- 150 Marengoni A, Angleman S, Meinow B, et al. Coexisting chronic conditions in the older population: Variation by health indicators. *Eur J Intern Med* 2016; **31**:29-34.

- 151 Marengoni A, von Strauss E, Rizzuto D, Winblad B, Fratiglioni L. The impact of chronic multimorbidity and disability on functional decline and survival in elderly persons. A community-based, longitudinal study. *J Intern Med* 2009; **265**:288-95.
- 152 Marques A, Santos DA, Peralta M, Sardinha LB, González Valeiro M. Regular physical activity eliminates the harmful association of television watching with multimorbidity. A cross-sectional study from the european social survey. *Prev Med* 2018; **109**:28-33.
- 153 Mavaddat N, Valderas JM, van der Linde R, Khaw KT, Kinmonth AL. Association of self-rated health with multimorbidity, chronic disease and psychosocial factors in a large middle-aged and older cohort from general practice: A cross-sectional study. *BMC Fam Pract* 2014; **15**:1-11.
- 154 McDaid O, Hanly MJ, Richardson K, Kee F, Kenny RA, Savva GM. The effect of multiple chronic conditions on self-rated health, disability and quality of life among the older populations of northern ireland and the republic of ireland: A comparison of two nationally representative cross-sectional surveys. *BMJ Open* 2013; **3**:1-9.
- 155 Melis R, Marengoni A, Angleman S, Fratiglioni L. Incidence and predictors of multimorbidity in the elderly: A population-based longitudinal study. *PLoS One* 2014; **9**:1-8.
- 156 Min LC, Wenger NS, Fung C, et al. Multimorbidity is associated with better quality of care among vulnerable elders. *Med Care* 2007; **45**:480-8.
- 157 Momtaz YA, Hamid TA, Yahaya N, Ibrahim R. Effects of chronic comorbidity on psychological well-being among older persons in northern peninsular Malaysia. *Appl Res Qual Life* 2010; **5**:133-46.
- 158 Mondor L, Cohen D, Khan AI, Wodchis WP. Income inequalities in multimorbidity prevalence in ontario, canada: A decomposition analysis of linked survey and health administrative data. *Int J Equity Health* 2018; **17**:1-13.
- 159 Muggah E, Graves E, Bennett C, Manuel DG. The impact of multiple chronic diseases on ambulatory care use; a population based study in ontario, canada. *BMC Health Serv Res* 2012; **12**:1-6.
- 160 Nagel A, Jungert A, Spinneker A, Neuhäuser-Berthold M. The impact of multimorbidity on resting metabolic rate in community-dwelling women over a ten-year period: A cross-sectional and longitudinal study. *J Nutr Health Aging* 2017; **21**:781-86.
- 161 Niedzwiedz CL, Katikireddi SV, Pell JP, Smith DJ. Sex differences in the association between salivary telomere length and multimorbidity within the us health & retirement study. *Age Ageing* 2019; **48**:703-10.
- 162 Nunes BP, Camargo-Figuera FA, Guttier M, et al. Multimorbidity in adults from a southern brazilian city: Occurrence and patterns. *Int J Public Health* 2016; **61**:1013-20.
- 163 Nunes BP, Chiavegatto Filho ADP, Pati S, et al. Contextual and individual inequalities of multimorbidity in brazilian adults: A cross-sectional national-based study. *BMJ Open* 2017; **7**:1-12.
- 164 Nunes BP, Thumé E, Facchini LA. Multimorbidity in older adults: Magnitude and challenges for the brazilian health system. *BMC Public Health* 2015; **15**:1-11.
- 165 Olaya B, Domènech-Abella J, Moneta MV, et al. All-cause mortality and multimorbidity in older adults: The role of social support and loneliness. *Exp Gerontol* 2017; **99**:120-26.
- 166 Olivares DEV, Chambi FRV, Chañi EMM, Craig WJ, Pacheco SOS, Pacheco FJ. Risk factors for chronic diseases and multimorbidity in a primary care context of

- central argentina: A web-based interactive and cross-sectional study. *Int J Environ Res Public Health* 2017; **14**:1-22.
- 167 Park B, Ock M, Lee HA, et al. Multimorbidity and health-related quality of life in
168 koreans aged 50 or older using knhanes 2013-2014. *Health Qual Life Outcomes* 2018;
169 **16**:1-10.
- 168 Patel KV, Peek MK, Wong R, Markides KS. Comorbidity and disability in elderly
169 mexican and mexican american adults: Findings from mexico and the southwestern
170 united states. *J Aging Health* 2006; **18**:315-29.
- 169 Pati S, Hussain MA, Swain S, et al. Development and validation of a questionnaire to
170 assess multimorbidity in primary care: An indian experience. *BioMed Res Int* 2016;
171 **2016**:1-9.
- 170 Pati SS, Subhashisa: Knottnerus, J.: Metsemakers, Job F.: van den Akker, Marjan.
171 Health related quality of life in multimorbidity: A primary-care based study from
172 odisha, India. *Health Qual Life Outcomes* 2019; **17**:1-11.
- 171 Payne RA, Abel GA, Guthrie B, Mercer SW. The effect of physical multimorbidity,
172 mental health conditions and socioeconomic deprivation on unplanned admissions to
173 hospital: A retrospective cohort study. *Can Med Assoc J* 2013; **185**:E221-8.
- 172 Pérez LM, Hooshmand B, Mangialasche F, et al. Glutathione serum levels and rate of
173 multimorbidity development in older adults. *J Gerontol A Biol Sci Med Sci* 2020;
174 **75**:1089-94.
- 173 Petersen I, Rathod S, Kathree T, Selohilwe O, Bhana A. Risk correlates for physical-
174 mental multimorbidities in south africa: A cross-sectional study. *Epidemiol Psychiatr
175 Sci* 2019; **28**:418-26.
- 174 Pfortmueller CA, Stotz M, Lindner G, Müller T, Rodondi N, Exadaktylos AK.
175 Multimorbidity in adult asylum seekers: A first overview. *PloS one* 2013; **8**:1-7.
- 175 Pressley JC Patrick CH. Frailty bias in comorbidity risk adjustments of community-
176 dwelling elderly populations. *J Clin Epidemiol* 1999; **52**:753-60.
- 176 Prior A, Fenger-Grøn M, Larsen KK, et al. The association between perceived stress
177 and mortality among people with multimorbidity: A prospective population-based
178 cohort study. *Am J Epidemiol* 2016; **184**:199-210.
- 177 Ribeiro AS, Pereira LC, Silva DRP, et al. Physical activity and sitting time are
178 specifically associated with multiple chronic diseases and medicine intake in brazilian
179 older adults. *J Aging Phys Act* 2018; **26**:608-13.
- 178 Ruel G, Lévesque J-F, Stocks N, et al. Understanding the evolution of multimorbidity:
179 Evidences from the north west adelaide health longitudinal study (nwahs). *PLOS ONE*
180 2014; **9**:1-11.
- 179 Ruel G, Shi Z, Zhen S, et al. Association between nutrition and the evolution of
180 multimorbidity: The importance of fruits and vegetables and whole grain products.
181 *Clin Nutr* 2014; **33**:513-20.
- 180 Ryan A, Murphy C, Boland F, Galvin R, Smith SM. What is the impact of physical
181 activity and physical function on the development of multimorbidity in older adults
182 over time? A population-based cohort study. *J Gerontol A Biol Sci Med Sci* 2018;
183 **73**:1538-44.
- 181 Schmidt AE, Ilinca S, Schulmann K, et al. Fit for caring: Factors associated with
182 informal care provision by older caregivers with and without multimorbidity. *Eur J
183 Ageing* 2016; **13**:103-13.
- 182 Schöttker B, Saum KU, Jansen EH, Holleczeck B, Brenner H. Associations of
183 metabolic, inflammatory and oxidative stress markers with total morbidity and multi-
184 morbidity in a large cohort of older german adults. *Age Ageing* 2016; **45**:127-35.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
- 183 Seo JC, B.: Kim, S.: Lee, H.: Oh, D. The relationship between multiple chronic diseases and depressive symptoms among middle-aged and elderly populations: Results of a 2009 Korean community health survey of 156,747 participants. *BMC Public Health* 2017; **17**:1-10.
- 184 She R, Yan Z, Jiang H, Vetrano DL, Lau JTF, Qiu C. Multimorbidity and health-related quality of life in old age: Role of functional dependence and depressive symptoms. *J Am Med Dir Assoc* 2019; **20**:1143-49.
- 185 Singh K, Patel SA, Biswas S, et al. Multimorbidity in south Asian adults: Prevalence, risk factors and mortality. *J Public Health (Oxf)* 2019; **41**:80-89.
- 186 Spangenberg L, Forkmann T, Brähler E, Glaesmer H. The association of depression and multimorbidity in the elderly: Implications for the assessment of depression. *Psychogeriatr* 2011; **11**:227-34.
- 187 Stickley A, Koyanagi A, Ueda M, Inoue Y, Waldman K, Oh H. Physical multimorbidity and suicidal behavior in the general population in the United States. *J Affect Disord* 2020; **260**:604-09.
- 188 Streit S, da Costa BR, Bauer DC, et al. Multimorbidity and quality of preventive care in Swiss university primary care cohorts. *PLOS ONE* 2014; **9**:1-7.
- 189 Stubbs B, Vancampfort D, Veronese N, et al. Multimorbidity and perceived stress: A population-based cross-sectional study among older adults across six low- and middle-income countries. *Maturitas* 2018; **107**:84-91.
- 190 Su P, Ding H, Zhang W, et al. The association of multimorbidity and disability in a community-based sample of elderly aged 80 or older in Shanghai, China. *BMC Geriatr* 2016; **16**:1-7.
- 191 Sundstrup E, Jakobsen MD, Mortensen OS, Andersen LL. Joint association of multimorbidity and work ability with risk of long-term sickness absence: A prospective cohort study with register follow-up. *Scand J Work Environ Health* 2017; **43**:146-54.
- 192 Takahashi PY, Ryu E, Hathcock MA, et al. A novel housing-based socioeconomic measure predicts hospitalisation and multiple chronic conditions in a community population. *J Epidemiol Community Health* 2016; **70**:286-91.
- 193 Tinetti MEM, G. J.: Chang, S. S.: Newman, A. B.: Fitzpatrick, A. L.: Fried, T. R.: Peduzzi, P. N. Contribution of multiple chronic conditions to universal health outcomes. *J Am Geriatr Soc* 2011; **59**:1686-91.
- 194 Troelstra SA, Straker L, Harris M, Brown S, van der Beek AJ, Coenen P. Multimorbidity is common among young workers and related to increased work absenteeism and presenteeism: Results from the population-based Raine study cohort. *Scand J Work Environ Health* 2020; **46**:218-27.
- 195 van Zon SKR, Reijneveld SA, Galaurchi A, Mendes de Leon CF, Almansa J, Bültmann U. Multimorbidity and the transition out of full-time paid employment: A longitudinal analysis of the Health and Retirement Study. *J Gerontol B Psychol Sci Soc Sci* 2020; **75**:705-15.
- 196 Vancampfort D, Stubbs B, Koyanagi A. Physical chronic conditions, multimorbidity and sedentary behavior amongst middle-aged and older adults in six low- and middle-income countries. *Int J Behav Nutr Phys Act* 2017; **14**:1-13.
- 197 Vassilaki M, Aakre JA, Cha RH, et al. Multimorbidity and risk of mild cognitive impairment. *J Am Geriatr Soc* 2015; **63**:1783-90.
- 198 Vassilaki M, Aakre JA, Mielke MM, et al. Multimorbidity and neuroimaging biomarkers among cognitively normal persons. *Neurology* 2016; **86**:2077-84.
- 199 Villarreal AEG, S.: Lopez, L.: Britton, G. B.: Initiative, P. A. R. Cognitive impairment, depression, and cooccurrence of both among the elderly in Panama:

- Differential associations with multimorbidity and functional limitations. *BioMed Res Int* 2015; **2015**:1-7.
- 200 Violán C, Foguet-Boreu Q, Fernández-Bertolín S, et al. Soft clustering using real-world data for the identification of multimorbidity patterns in an elderly population: Cross-sectional study in a mediterranean population. *BMJ Open* 2019; **9**:1-14.
- 201 von Strauss E, Fratiglioni L, Viitanen M, Forsell Y, Winblad B. Morbidity and comorbidity in relation to functional status: A community-based study of the oldest old (90+ years). *J Am Geriatr Soc* 2000; **48**:1462-9.
- 202 Vos HM, Bor HH, Rangelrooij-Minkels MJ, Schellevis FG, Lagro-Janssen AL. Multimorbidity in older women: The negative impact of specific combinations of chronic conditions on self-rated health. *Eur J Gen Pract* 2013; **19**:117-22.
- 203 Vu HMN, L. H.: Tran, T. H.: Pham, K. T. H.: Phan, H. T.: Nguyen, H. N.: Tran, B. X.: Latkin, C. A.: Ho, C. S. H.: Ho, R. C. M. Effects of chronic comorbidities on the health-related quality of life among older patients after falls in vietnamese hospitals. *Int J Environ Res Public Health* 2019; **16**:1-9.
- 204 Wang L, Cocker F, Kilpatrick M, et al. The associations of multimorbidity with health-related productivity loss in a large and diverse public sector setting: A cross-sectional survey. *J Occup Environ Med* 2018; **60**:528-35.
- 205 Wang X-X, Lin W-Q, Chen X-J, et al. Multimorbidity associated with functional independence among community-dwelling older people: A cross-sectional study in southern china. *Health Qual Life Outcomes* 2017; **15**:73-9.
- 206 Wijers IGM, Ayala A, Rodriguez-Blazquez C, et al. The disease burden morbidity assessment in older adults and its association with mortality and other health outcomes. *Eur J Ageing* 2019; **16**:193-203.
- 207 Williams JSE, L. E. The association between multimorbidity and quality of life, health status and functional disability. *Am J Med Sci* 2016; **352**:45-52.
- 208 Woldeamayem EM, Kassa A, Gari T, Dangisso MH. Chronic diseases multimorbidity among adult patients at hawassa university comprehensive specialized hospital. *BMC Public Health* 2018; **18**:1-8.
- 209 Yao SS, Cao GY, Han L, et al. Associations between somatic multimorbidity patterns and depression in a longitudinal cohort of middle-aged and older chinese. *J Am Med Dir Assoc* 2020; **21**:1282-87.
- 210 Yorke AM, Curtis AB, Shoemaker M, Vangsnes E. The impact of multimorbidity on grip strength in adults age 50 and older: Data from the health and retirement survey (hrs). *Arch Gerontol Geriatr* 2017; **72**:164-68.
- 211 You L, Yu Z, Zhang X, et al. Association between multimorbidity and depressive symptom among community-dwelling elders in eastern china. *Clin Interv Aging* 2019; **14**:2273-80.
- 212 Zhang L, Ma L, Sun F, Tang Z, Chan P. A multicenter study of multimorbidity in older adult inpatients in china. *J Nutr Health Aging* 2020; **24**:269-76.
- 213 Khanam MA, Streatfield PK, Kabir ZN, Qiu C, Cornelius C, Wahlin Å. Prevalence and patterns of multimorbidity among elderly people in rural bangladesh: A cross-sectional study. *J Health, Popul Nutr* 2011; **29**:406-14.
- 214 Cornell J, Pugh J, Williams, Jr., et al. Multimorbidity clusters: Clustering binary data from multimorbidity clusters: Clustering binary data from a large administrative medical database. *Appl Multivar Res* 2009; **12**:1-6.
- 215 Cassell A, Edwards D, Harshfield A, et al. The epidemiology of multimorbidity in primary care: A retrospective cohort study. *Br J GenPract* 2018; **68**:e245-e51.

216 Wong ELY, Xu RH, Cheung AWL. Measuring the impact of chronic conditions and
associated multimorbidity on health-related quality of life in the general population in
hong kong sar, china: A cross-sectional study. *PLOS ONE* 2019; **14**:1-16.

217 Puth M-T, Weckbecker K, Schmid M, Münster E. Prevalence of multimorbidity in
germany: Impact of age and educational level in a cross-sectional study on 19,294
adults. *BMC Public Health* 2017; **17**:1-7.

For peer review only



PRISMA Checklist

| Section/topic | # | Checklist item | Reported on page # |
|------------------------------------|----|---|-----------------------------|
| TITLE | | | |
| Title | 1 | Identify the report as a systematic review, meta-analysis, or both. | Page 1 |
| ABSTRACT | | | |
| Structured summary | 2 | Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number. | Page 2 |
| INTRODUCTION | | | |
| Rationale | 3 | Describe the rationale for the review in the context of what is already known. | Page 4 |
| Objectives | 4 | Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS). | Page 5 |
| METHODS | | | |
| Protocol and registration | 5 | Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number. | Page 2 |
| Eligibility criteria | 6 | Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. | Page 5 |
| Information sources | 7 | Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched. | Page 6 |
| Search | 8 | Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated. | Supplementary Table S1 |
| Study selection | 9 | State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis). | Page 6, Figure 1 |
| Data collection process | 10 | Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators. | Page 6 |
| Data items | 11 | List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made. | Supplementary Table S8 |
| Risk of bias in individual studies | 12 | Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis. | Page 6 Supplementary p26 |
| Summary measures | 13 | State the principal summary measures (e.g., risk ratio, difference in means). | Page 7-8 |
| Synthesis of results | 14 | Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis. | Page 7-8 |

| Section/topic | # | Checklist item | Reported on page # |
|---------------|---|----------------|--------------------|
|---------------|---|----------------|--------------------|



PRISMA Checklist

| | | | |
|-------------------------------|----|--|---|
| Risk of bias across studies | 15 | Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies). | Page 10 and Table 2 |
| Additional analyses | 16 | Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified. | Page 7-8 |
| RESULTS | | | |
| Study selection | 17 | Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram. | Figure 1 |
| Study characteristics | 18 | For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations. | Page 8-9, Table 1; Supplementary Table S4 |
| Risk of bias within studies | 19 | Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12). | Supplementary Table S6 |
| Results of individual studies | 20 | For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot. | Page 9-10 Figure 2-4 |
| Synthesis of results | 21 | Present results of each meta-analysis done, including confidence intervals and measures of consistency. | Page 9-10 Table 2 |
| Risk of bias across studies | 22 | Present results of any assessment of risk of bias across studies (see Item 15). | Page 9-10, Table 1 and Table 2 |
| Additional analysis | 23 | Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]). | Page 10-11 |
| DISCUSSION | | | |
| Summary of evidence | 24 | Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers). | Page 11,12 |
| Limitations | 25 | Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias). | Page 13 |
| Conclusions | 26 | Provide a general interpretation of the results in the context of other evidence, and implications for future research. | Page 14 |
| FUNDING | | | |
| Funding | 27 | Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. | Page 15 |

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.